

# Measurement of b hadron fractions in 13 TeV pp collisions

LHCb Collaboration

DOI:

[10.1103/PhysRevD.100.031102](https://doi.org/10.1103/PhysRevD.100.031102)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

LHCb Collaboration 2019, 'Measurement of b hadron fractions in 13 TeV pp collisions', *Physical Review D*, vol. 100, no. 3, 031102. <https://doi.org/10.1103/PhysRevD.100.031102>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Measurement of  $b$  hadron fractions in 13 TeV  $pp$  collisionsR. Aaij *et al.*\*  
(LHCb Collaboration)

(Received 18 February 2019; published 27 August 2019)

The production fractions of  $\bar{B}_s^0$  and  $\Lambda_b^0$  hadrons, normalized to the sum of  $B^-$  and  $\bar{B}^0$  fractions, are measured in 13 TeV  $pp$  collisions using data collected by the LHCb experiment, corresponding to an integrated luminosity of  $1.67 \text{ fb}^{-1}$ . These ratios, averaged over the  $b$  hadron transverse momenta from 4 to 25 GeV and pseudorapidity from 2 to 5, are  $0.122 \pm 0.006$  for  $\bar{B}_s^0$ , and  $0.259 \pm 0.018$  for  $\Lambda_b^0$ , where the uncertainties arise from both statistical and systematic sources. The  $\Lambda_b^0$  ratio depends strongly on transverse momentum, while the  $\bar{B}_s^0$  ratio shows a mild dependence. Neither ratio shows variations with pseudorapidity. The measurements are made using semileptonic decays to minimize theoretical uncertainties. In addition, the ratio of  $D^+$  to  $D^0$  mesons produced in the sum of  $\bar{B}^0$  and  $B^-$  semileptonic decays is determined as  $0.359 \pm 0.006 \pm 0.009$ , where the uncertainties are statistical and systematic.

DOI: [10.1103/PhysRevD.100.031102](https://doi.org/10.1103/PhysRevD.100.031102)

Knowledge of the fragmentation fractions of  $\bar{B}_s^0$  ( $f_s$ ) and  $\Lambda_b^0$  ( $f_{\Lambda_b^0}$ ) hadrons is essential for determining absolute branching fractions ( $\mathcal{B}$ ) of decays of these hadrons at the LHC, allowing measurements, e.g., of  $\mathcal{B}(\bar{B}_s^0 \rightarrow \mu^+\mu^-)$  [1] and the future evaluation of  $|V_{cb}|$  from  $\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu$  decays [2].<sup>1</sup> Once these fractions are determined, measurements of absolute branching fractions of  $B^-$  and  $\bar{B}^0$  mesons performed at  $e^+e^-$  colliders operating at the  $\Upsilon(4S)$  resonance can be used to determine the  $\bar{B}_s^0$  and  $\Lambda_b^0$  branching fractions [3].

In this paper we measure the ratios  $f_s/(f_u + f_d)$  and  $f_{\Lambda_b^0}/(f_u + f_d)$ , where the denominator is the sum of  $B^-$  and  $\bar{B}^0$  contributions, in the LHCb acceptance of pseudorapidity  $2 < \eta < 5$  and transverse momentum  $4 < p_T < 25 \text{ GeV}$ ,<sup>2</sup> in 13 TeV  $pp$  collisions. These ratios can depend on  $p_T$  and  $\eta$ ; therefore, we perform the analysis using two-dimensional binning.

Much of the analysis method adopted in this study is an evolution of our previous  $b$  hadron fraction measurements for 7 TeV  $pp$  collisions [4]. We use the inclusive semileptonic decays  $H_b \rightarrow H_c X \mu^- \bar{\nu}_\mu$ , where  $H_b$  indicates a  $b$  hadron,  $H_c$  a charm hadron, and  $X$  possible additional particles. Each of the different  $H_c$  plus muon final states can originate from the decay of different  $b$  hadrons.

Semileptonic decays of  $\bar{B}^0$  mesons usually result in a mixture of  $D^0$  and  $D^+$  mesons, while  $B^-$  mesons decay predominantly into  $D^0$  mesons with a smaller admixture of  $D^+$  mesons. Both include a tiny component of  $D_s^+ \bar{K}$  meson pairs. Similarly,  $\bar{B}_s^0$  mesons decay predominantly into  $D_s^+$  mesons, but can also decay into  $D^0 K^+$  and  $D^+ K^0$  meson pairs; this is expected if the  $\bar{B}_s^0$  meson decays into an excited  $D_s^+$  state that is heavy enough to decay into a  $DK$  pair. We measure this contribution using  $D^0 K^+ X \mu^- \bar{\nu}_\mu$  events. Finally,  $\Lambda_b^0$  baryons decay semileptonically mostly into  $\Lambda_c^+$  final states, but can also decay into  $D^0 p$  and  $D^+ n$  pairs. We ignore the contributions of  $b \rightarrow u$  decays that comprise approximately 1% of semileptonic  $b$  hadron decays and contribute almost equally to all  $b$  hadron species. The detailed equations relating these yields to the final results are given in Ref. [4] and in the Supplemental Material [5].

The theoretical basis for this measurement is the near equality of semileptonic widths,  $\Gamma_{\text{SL}}$ , for all  $b$  hadron species [6] whose differences are predicted to precisions of about 1%. The values we use for the individual  $H_b$  semileptonic branching fractions ( $\mathcal{B}_{\text{SL}}$ ) are listed in Table I. The  $H_c$  decay modes used and their branching fractions are given in Table II.

The ratio of  $D^+$  to  $D^0$  meson production in the sum of semileptonic  $\bar{B}^0$  and  $B^-$  decays,  $f_+/f_0$ , is used to check the analysis method. This result can be related to models of the hadronic final states in  $B^-$  and  $\bar{B}^0$  semileptonic decays [11].

The data sample corresponds to  $1.67 \text{ fb}^{-1}$  of integrated luminosity obtained with the LHCb detector in 13 TeV  $pp$  collisions during 2016. The LHCb detector [12,13] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks.

\*Full author list given at end of the article.

<sup>1</sup>Mention of a particular decay mode implies the use of the charge-conjugate one as well.<sup>2</sup>We use natural units where  $c = \hbar = 1$ .

TABLE I. Branching fractions of semileptonic  $b$  hadron decays from direct measurements for  $\bar{B}^0$  and  $B^-$  mesons, ( $\langle B \rangle \equiv \langle \bar{B}^0 + B^- \rangle$ ), and derived for  $\bar{B}_s^0$  and  $\Lambda_b^0$  hadrons based on the equality of semileptonic widths and the lifetime ratios [3,6]. Corrections to  $\Gamma_{\text{SL}}$  for  $\bar{B}_s^0$  ( $-1.0 \pm 0.5\%$ ) and  $\Lambda_b^0$  ( $3.0 \pm 1.5\%$ ) are applied [6]. Correlations in the  $\bar{B}^0$  and  $B^-$  branching fraction measurements have been taken into account. See Ref. [7] for more information.

Particle	$\tau$ (ps)	$\mathcal{B}_{\text{SL}}$ (%)	$\mathcal{B}_{\text{SL}}$ (%)
	measured	measured	used
$\bar{B}^0$	$1.520 \pm 0.004$	$10.30 \pm 0.19$	$10.30 \pm 0.19$
$B^-$	$1.638 \pm 0.004$	$11.08 \pm 0.20$	$11.08 \pm 0.20$
$\langle B \rangle$		$10.70 \pm 0.19$	$10.70 \pm 0.19$
$\bar{B}_s^0$	$1.526 \pm 0.015$		$10.24 \pm 0.21$
$\Lambda_b^0$	$1.470 \pm 0.010$		$10.26 \pm 0.25$

TABLE II. Charm-hadron branching fractions for the decay modes used in this analysis. Note that the  $\Lambda_c^+$  branching fraction has been significantly improved since the previous analysis.

Decay	$\mathcal{B}$ (%)	Source
$D^0 \rightarrow K^- \pi^+$	$3.93 \pm 0.05$	PDG average [3]
$D^+ \rightarrow K^- \pi^+ \pi^+$	$9.22 \pm 0.17$	CLEO-c [8]
$D_s^+ \rightarrow K^- K^+ \pi^+$	$5.44 \pm 0.18$	PDG average [3]
$\Lambda_c^+ \rightarrow p K^- \pi^+$	$6.23 \pm 0.33$	From Refs. [9,10]

The online event selection is performed by a trigger [14] which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. At the hardware trigger stage, events are required to have a muon with large  $p_T$  or a hadron, photon or electron with high transverse energy in the calorimeters. For hadrons, the transverse energy threshold is 3.5 GeV. The software trigger requires a two-, three- or four-track secondary vertex with a significant displacement from any primary  $pp$  interaction vertex (PV). At least one charged particle must have  $p_T > 1.6$  GeV and be inconsistent with originating from a PV. A multivariate algorithm [15] is used for the identification of secondary vertices consistent with the decay of a  $b$  hadron.

Simulation is required to model the effects of the detector acceptance and the imposed selection requirements. Here  $pp$  collisions are generated using PYTHIA [16] with a specific LHCb configuration [17]. Decays of unstable particles are described by EVTGEN [18], in which final-state radiation is generated using PHOTOS [19]. The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [20] as described in Ref. [21].

Selection criteria are applied to muons and  $H_c$  decay particles. The transverse momentum of each hadron must

be greater than 0.3 GeV, and that of the muon larger than 1.3 GeV. Each track cannot point to any PV, implemented by requiring  $\chi_{\text{IP}}^2 > 9$  with respect to any PV, where  $\chi_{\text{IP}}^2$  is defined as the difference in the vertex-fit  $\chi^2$  of a given PV reconstructed with and without the track under consideration being included. All final-state particles are required to be positively identified using information from the Ring Imaging CHerenkov detectors particle identification (PID). Particles from  $H_c$  decay candidates must have a good fit to a common vertex with  $\chi^2/\text{ndof} < 9$ , where ndof is the number of degrees of freedom. They must also be well separated from the nearest PV, with the flight distance divided by its uncertainty greater than 5.

Candidate  $b$  hadrons are formed by combining  $H_c$  and muon candidates originating from a common vertex with  $\chi^2/\text{ndof} < 9$  and an  $H_c \mu^-$  invariant mass,  $m_{H_c \mu^-}$ , in the range 3.0–5.0 GeV for  $D^0$  and  $D^+$ , 3.1–5.1 GeV for  $D_s^+$  and 3.3–5.3 GeV for  $\Lambda_c^+$  candidates. In addition, we define  $m_{\text{corr}} \equiv \sqrt{m_{H_c \mu}^2 + p_{\perp}^2 + p_{\perp}^2}$ , where  $p_{\perp}$  is the magnitude of the combination's momentum component transverse to the  $b$  hadron flight direction; we require that  $m_{\text{corr}} > 4.2$  or 4.5 GeV for  $\bar{B}_s^0$  or  $\Lambda_b^0$  candidates, respectively. For the  $D_s^+ \rightarrow K^+ K^- \pi^+$  decay mode, vetoes are employed to remove backgrounds from real  $D^+$  or  $\Lambda_c^+$  decays where the particle assignments are incorrect.

Background from prompt  $H_c$  production at the PV needs to be considered. We use the natural logarithm of the  $H_c$  impact parameter, IP, with respect to the PV in units of mm. Requiring  $\ln(\text{IP}/\text{mm}) > -3$  is found to reduce the prompt component to be below 0.1%, while preserving 97% of all signals. This restriction allows us to perform fits only to the  $H_c$  candidate mass spectra to find the  $b$  hadron decay yields.

The  $H_c$  candidates' mass distributions integrated over  $p_T(H_b)$  and  $\eta$  are shown in Fig. 1. They consist of a prominent peak resulting from signal and a small contribution due to combinatorial background from random combinations of particles that pass the selection. They are fit with a signal component comprised of two Gaussian functions and a combinatorial background component modeled as a linear function. The total signal yields for  $D^0 X \mu^- \bar{\nu}_\mu$ ,  $D^+ X \mu^- \bar{\nu}_\mu$ ,  $D_s^+ X \mu^- \bar{\nu}_\mu$  and  $\Lambda_c^+ \mu^- X \bar{\nu}_\mu$  are 13 775 000, 4 282 700, 845 300, and 1 753 600, respectively.

Background contributions to the  $b$  hadron candidates include hadrons faking muons, false combinations of charm hadrons and muons from the two  $b$  hadrons in the event, as well as real muons and charm hadrons from  $B \rightarrow D\bar{D}X$  decays, where one of the  $D$  mesons decays into a muon. All the backgrounds are evaluated in two-dimensional  $\eta$  and  $p_T$  intervals. The first two backgrounds are evaluated using events where the  $H_c$  is combined with a muon of the wrong sign (e.g.,  $D^0 \mu^+$ ), forbidden in a semileptonic  $b$  hadron decay. The wrong-sign backgrounds are  $< 1\%$  for each  $H_c$  species. The background from  $B \rightarrow D\bar{D}X$  decays is




FIG. 1. Fit to the mass spectra of the  $H_c$  candidates of the selected  $H_b$  decays: (a)  $D^0$ , (b)  $D^+$ , (c)  $D_s^+$  mesons, and (d) the  $\Lambda_c^+$  baryon. The data are shown as black points with error bars. The signal component is shown as the dashed (green) line and the combinatorial background component is shown as the dashed (red) line. The solid (blue) line shows all components added together.

determined by simulating a mixture of these decays using their measured branching fractions [3]. The only decay mode significantly affected is  $\bar{B}_s^0 \rightarrow D_s^+ X \mu^- \bar{\nu}_\mu$  with contributions varying from 0.1% for  $D^0 D_s^- X$  to 1.8% for  $D_s^+ D_s^- X$  due to the large  $D_s^+ \rightarrow \mu^+ \nu$  decay rate. The total  $B \rightarrow D\bar{D}X$  background is  $(5.8 \pm 0.9)\%$ .

The dominant component in  $\bar{B}_s^0$  semileptonic decays is  $D_s^+ X \mu^- \bar{\nu}_\mu$ , where  $X$  contains possible additional hadrons. However, the  $\bar{B}_s^0$  meson also can decay into  $D^0 K^+$  or  $D^+ K^0$  instead of  $D_s^+$ , so we must add this component to the  $\bar{B}_s^0$  rate and subtract it from the  $f_u + f_d$  fraction. Similarly, in  $\Lambda_b^0$  semileptonic decays we find a  $D^0 p X$  component. The selection criteria for these final states are similar to those for the  $D^0 X \mu^- \bar{\nu}_\mu$  and  $\Lambda_c^+ X \mu^- \bar{\nu}_\mu$  final states described above with the addition of a kaon or proton with  $p_T > 300$  MeV that has been positively identified. A veto is also applied to reject  $D^{*+} \rightarrow \pi^+ D^0$  decays where the pion mimics a kaon or a proton.

These samples contain background, resonant and non-resonant decays. Separation of these components is achieved by using both right-sign ( $H_c$  with  $\mu^-$ ) and wrong-sign ( $H_c$  with  $\mu^+$ ) candidates. In addition, the logarithm of the difference between the vertex  $\chi^2$  formed by the added hadron track and the  $D\mu$  system and the vertex  $\chi^2$  of the  $D\mu$  system,  $\ln(\Delta\chi_V^2)$ , provides separation between combinatorial background and nonresonant semileptonic decays. True resonant and nonresonant  $\bar{B}_s^0 \rightarrow D^0 K^+ \mu^- \bar{\nu}_\mu$  or  $\Lambda_b^0 \rightarrow D^0 p \mu^- \bar{\nu}_\mu$  decays peak in the  $\ln(\Delta\chi_V^2)$  distribution at a value of unity

while the background is smooth and rises at higher values as the added track is generally not associated with the  $D^0 \mu^-$  vertex. To distinguish signal from background we define  $m(D^0 h)_C \equiv m(D^0 h) - m(D^0) + m(D^0)_{\text{PDG}}$  and perform two-dimensional fits to the  $m(D^0 h)_C$  and  $\ln(\Delta\chi_V^2)$  distributions, where  $h = K^+(p)$  for right-sign  $\bar{B}_s^0$  ( $\Lambda_b^0$ ) decays.

The wrong-sign shapes are used to model the backgrounds. The resonant structures are modeled with relativistic Breit-Wigner functions convoluted with Gaussians to take into account the experimental resolution, except for the narrow  $D_{s1}(2536)^+$  which is modeled with the sum of two Gaussians with a fixed mean. The nonresonant shape for the  $\ln(\Delta\chi_V^2)$  distribution is taken as the same as the resonant one. Figure 2 shows the data and result of the fits for  $\bar{B}_s^0$  and  $\Lambda_b^0$  candidates.

For the  $\bar{B}_s^0$  case, we find  $22\,610 \pm 210$   $D_{s1}(2536)^+$ ,  $14\,290 \pm 260$   $D_{s2}^*(2573)^+$ , and  $38\,140 \pm 460$  nonresonant decays, confirming the existence of both the  $D_{s1}^+$  [22,23] and  $D_{s2}^{*+}$  [23] particles in semileptonic  $\bar{B}_s^0$  decays with substantially more data, and showing the existence of the nonresonant component. To account for the unmeasured  $D^+ K^0$  channel we take different mixtures of  $D^*$  and  $D$  final states for the different resonant and nonresonant components. The  $D_{s1}^+$  decays dominantly into  $D^*$ , while the  $D_{s2}^{*+}$  decays dominantly into  $D$  mesons [3]. For the nonresonant part we assume equal  $D^*$  and  $D$  yields.

In the  $\Lambda_b^0$  case, we find  $6120 \pm 460$   $\Lambda_c^+(2860)$ ,  $2200 \pm 200$   $\Lambda_c^+(2880)$ ,  $1200 \pm 260$   $\Lambda_c^+(2940)$ , and  $29\,770 \pm 690$




FIG. 2. Projections of the two-dimensional fits to the (a)  $m(D^0 K^\pm)_C$  and (c)  $m(D^0 \bar{p})_C$  mass distributions and (b),(d)  $\ln(\Delta\chi^2_V)$  for (top)  $D^0 K^\pm X \mu^- \bar{\nu}_\mu$  candidates and for (bottom)  $D^0 \bar{p} X \mu^- \bar{\nu}_\mu$  candidates. The curves show projections of the 2D fit. The dashed (red) curves show the  $D_{s1}^+$  and  $D_{s2}^+$  resonant components in (a) and (b), and  $\Lambda_c^+(2860)$ ,  $\Lambda_c^+(2880)$  and  $\Lambda_c^+(2940)$  resonant components in (c) and (d). The long-dashed-dotted (green) curves show the nonresonant component, the dotted (black) curves are the background components, whose shapes are determined from wrong-sign combinations, and the solid (blue) curves shows all components added together.

nonresonant events. The decay rate into  $D^0 p$  is assumed to be equal to that into  $D^+ n$  using isospin conservation. All decays with an extra hadron have lower detection efficiencies than the sample without.

Efficiencies for all the samples are determined using data in two-dimensional  $p_T$  and  $\eta$  bins. Trigger efficiencies are determined using a sample of  $B^- \rightarrow J/\psi K^-$ , with  $J/\psi \rightarrow \mu^+ \mu^-$  decays where only one muon track is positively identified, in conjunction with viewing the effects of combinations of different triggers [24]. This sample is also used to determine muon identification efficiencies. Decays of  $J/\psi$  mesons to muons reconstructed using partial information from the tracking system, e.g., eliminating the vertex locator information, are also used to determine tracking efficiencies using data and to correct the simulation. Finally, the PID efficiencies are evaluated using kaons and pions from  $D^{*+} \rightarrow \pi^+ D^0$  decays, with  $D^0 \rightarrow K^- \pi^+$ , and protons from  $\Lambda \rightarrow p \pi^-$  and  $\Lambda_c^+ \rightarrow p K^- \pi^+$  decays [25]. In the measurement of  $b$  hadron fraction ratios many of the efficiencies cancel and we are left with only residual effects to which we assign systematic uncertainties.

The  $b$  hadron  $\eta$  and  $p_T$ ,  $p_T(H_b)$ , must be known because the  $b$  fractions can depend on production kinematics. While  $\eta$  can be evaluated directly using the measured primary and

secondary  $b$  vertices, the value of  $p_T(H_b)$  must be determined to account for the missing neutrino plus extra particles. The correction factor  $k$  is given by the ratio of the average reconstructed to true  $p_T(H_b)$  as a function of  $m(H_c \mu^-)$  and is determined using simulation. It varies from 0.75 for  $m(H_c \mu^-) = 3$  GeV to unity at  $m(H_c \mu^-) = m(H_b)$ .

The distribution of  $f_s/(f_u + f_d)$  as a function of  $p_T(H_b)$  is shown in Fig. 3. We perform a linear  $\chi^2$  fit incorporating a full covariance matrix which takes into account the bin-by-bin correlations introduced from the kaon kinematics, and PID and tracking systematic uncertainties. The factor  $A$  in Eq. (1) incorporates the global systematic uncertainties described later, which are independent of  $p_T(H_b)$ . The resulting function is

$$\frac{f_s}{f_u + f_d}(p_T) = A[p_1 + p_2 \times (p_T - \langle p_T \rangle)], \quad (1)$$

where  $p_T$  here refers to  $p_T(H_b)$ ,  $A = 1 \pm 0.043$ ,  $p_1 = 0.119 \pm 0.001$ ,  $p_2 = (-0.91 \pm 0.25) \times 10^{-3} \text{ GeV}^{-1}$ , and  $\langle p_T \rangle = 10.1$  GeV. The correlation coefficient between the fit parameters is 0.20. After integrating over  $p_T(H_b)$ , no  $\eta$  dependence is observed (see the Supplemental Material [5]).




FIG. 3. The ratios  $f_s/(f_u + f_d)$  and  $f_{\Lambda_b^0}/(f_u + f_d)$  in bins of  $p_T(H_b)$ . The  $\bar{B}_s^0$  data are indicated by solid circles, while the  $\Lambda_b^0$  by triangles. The smaller (black) error bars show the combined bin-by-bin statistical and systematic uncertainties, and the larger (blue) ones show the global systematics added in quadrature. The fits to the data are shown as the solid (green) bands, whose widths represents the  $\pm 1\sigma$  uncertainty limits on the fit shapes, and the dashed (black) lines give the total uncertainty on the fit results including the global scale uncertainty. In the highest two  $p_T$  bins the points have been displaced from the center of the bin.

We determine an average value for  $f_s/(f_u + f_d)$  by dividing the yields of  $\bar{B}_s^0$  semileptonic decays by the sum of  $\bar{B}^0$  and  $B^-$  semileptonic yields, which are all efficiency-corrected, between the limits of  $p_T(H_b)$  of 4 and 25 GeV and  $\eta$  of 2 and 5, resulting in

$$\frac{f_s}{f_u + f_d} = 0.122 \pm 0.006,$$

where the uncertainty contains both statistical and systematic components, with the latter being dominant, and discussed subsequently. The total relative uncertainty is 4.8%.

Figure 3 also shows the  $\Lambda_b^0$  fraction as a function of  $p_T(H_b)$  demonstrating a large  $p_T$  dependence. The distribution in  $\eta$  is flat. We perform a similar fit as in the  $\bar{B}_s^0$  fraction case, using

$$\frac{f_{\Lambda_b^0}}{f_u + f_d}(p_T) = A[p_1 + \exp(p_2 + p_3 \times p_T)], \quad (2)$$

where  $p_T$  here refers to  $p_T(H_b)$ ,  $A = 1 \pm 0.061$ ,  $p_1 = (7.93 \pm 1.41) \times 10^{-2}$ ,  $p_2 = -1.022 \pm 0.047$ , and  $p_3 = -0.107 \pm 0.002 \text{ GeV}^{-1}$ . The correlation coefficients among the fit parameters are 0.40 ( $\rho_{12}$ ),  $-0.95$  ( $\rho_{13}$ ), and  $-0.63$  ( $\rho_{23}$ ).

The average value for  $f_{\Lambda_b^0}/(f_u + f_d)$  is determined using the same method as in the  $\bar{B}_s^0$  case. The result is

$$\frac{f_{\Lambda_b^0}}{f_u + f_d} = 0.259 \pm 0.018,$$

where the dominant uncertainty is systematic, and the statistical uncertainty is included. The overall uncertainty is 6.9%.

As a systematic check of the analysis method, and a useful measurement to test the knowledge of known semileptonic branching fractions and extrapolations used to saturate the unknown portion of the inclusive hadron spectrum, we measure the ratio of the  $D^0 X \mu^- \bar{\nu}_\mu$  to  $D^+ X \mu^- \bar{\nu}_\mu$  corrected yields  $f_+/f_0$ . We subtract the small contributions from  $\bar{B}_s^0$  and  $\Lambda_b^0$  decays, and a very small contribution from  $B \rightarrow D_s^+ \bar{K} \mu^- X$  decays has been taken into account [26], as in all the fractions measured above.

Assuming  $f_u$  equals  $f_d$ , Ref. [11] estimates the fraction of  $D^+ \mu^-$  with respect to  $D^0 \mu^-$  modes in the sum of  $B^-$  and  $\bar{B}^0$  decays as  $0.387 \pm 0.012 \pm 0.026$ . The first uncertainty comes from the uncertainties on known measurements. The second uncertainty comes from the different extrapolations from excited  $D$  mesons used to saturate the remaining portion of the inclusive rate.

The  $f_+/f_0$  ratio must be independent of  $\eta$  and  $p_T$ . To derive an overall value for  $f_+/f_0$ , the  $p_T(H_b)$  distribution is fit to a constant. Only the PID and tracking systematic uncertainties on the second pion in the  $D^+$  decay need be considered. Performing a  $\chi^2$  fit using the full covariance matrix we find  $f_+/f_0 = 0.359 \pm 0.006 \pm 0.009$ , where the first uncertainty is from bin-by-bin statistical and systematic uncertainties, including correlations, and the second is systematic. The  $\chi^2/\text{ndof}$  is 0.63, in agreement with a flat spectrum. The measurement is consistent with the prediction and places some constraints on the  $D^{**}$  content of semileptonic  $B$  decays [11].

The dominant global systematic uncertainties are listed in Table III. Simulation uncertainties are due to the

TABLE III. Global systematic uncertainties. The  $D^0$  and  $D^+$  branching fraction uncertainties are scaled by the fraction of each decay,  $f_0$  and  $f_+$  for  $f_s/(f_u + f_d)$  and  $f_{\Lambda_b^0}/(f_u + f_d)$  uncertainties.

Source	Value (%)		
	$f_s/(f_u + f_d)$	$f_{\Lambda_b^0}/(f_u + f_d)$	$f_+/f_0$
Simulation	1.7	2.4	...
Backgrounds	0.9	0.3	...
Cross feeds	1.2	0.4	0.2
$\mathcal{B}(D^0 \rightarrow K^- \pi^+)$	1.0	1.0	1.3
$\mathcal{B}(D^+ \rightarrow K^+ \pi^- \pi^-)$	0.6	0.6	1.8
$\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+)$	3.3	...	...
$\mathcal{B}(\Lambda_c^+ \rightarrow p K^+ \pi^-)$	...	5.3	...
Measured lifetime ratio	1.2	0.7	...
$\Gamma_{SL}$ correction	0.5	1.5	...
Total	4.3	6.1	2.2

modeling of excited charm states for the  $f_s/(f_u + f_d)$  determination and the weighting required for the  $f_{\Lambda_b^0}/(f_u + f_d)$  ratio, due to differences between the simulated and measured  $p_T$  spectra. Background uncertainties arise from  $D\bar{D}X$  final states with uncertain branching fractions. Cross-feed uncertainties come from errors on efficiency estimates and the assumed  $D^*$  to  $D$  mixtures. Other smaller uncertainties depend on  $p_T(H_b)$  and include tracking 0.2%–1.8%, particle identification 0.4%–3.0%, trigger 0.3%–3.9% and  $k$ -factor 0.2%–1.8%.

In conclusion, we measure the ratios of  $\bar{B}_s^0$  and  $\Lambda_b^0$  production to the sum of  $B^-$  and  $\bar{B}^0$  to be  $p_T(H_b)$  dependent [see Eqs. (1) and (2)]. The averages in the ranges  $4 < p_T(H_b) < 25$  GeV and  $2 < \eta < 5$  are  $f_s/(f_u + f_d) = 0.122 \pm 0.006$  and  $f_{\Lambda_b^0}/(f_u + f_d) = 0.259 \pm 0.018$ , respectively. Using 7 TeV data, LHCb determined  $f_s/(f_u + f_d) = 0.1295 \pm 0.0075$  with a  $p_T(H_b)$  slope larger than, but consistent with, these 13 TeV results [27]; no dependence on  $\eta$  was observed. For the  $\Lambda_b^0$  baryon, the fraction ratio is consistent with the 7 TeV measurements after taking into account the different  $p_T(H_b)$  ranges used [4,28,29]. We observe no rapidity dependence over a similar  $p_T(H_b)$  range as in Ref. [29].

These results are crucial for determining absolute branching fractions of  $\bar{B}_s^0$  and  $\Lambda_b^0$  hadron decays in LHC experiments. We also determine the ratio of  $D^0$  to  $D^+$  mesons produced in the sum of  $\bar{B}^0$  and  $B^-$  semileptonic decays as  $f_+/f_0 = 0.359 \pm 0.006 \pm 0.009$ .

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from the AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom); and Laboratory Directed Research and Development program of LANL (USA).

- 
- [1] V. Khachatryan *et al.* (CMS and LHCb collaborations), Observation of the rare  $B_s^0 \rightarrow \mu^+ \mu^-$  decay from the combined analysis of CMS and LHCb data, *Nature (London)* **522**, 68 (2015).
- [2] R. Aaij *et al.* (LHCb Collaboration), Measurement of the shape of the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- \bar{\nu}_\mu$  differential decay rate, *Phys. Rev. D* **96**, 112005 (2017).
- [3] M. Tanabashi *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **98**, 030001 (2018).
- [4] R. Aaij *et al.* (LHCb Collaboration), Measurement of  $b$  hadron production fractions in 7 TeV pp collisions, *Phys. Rev. D* **85**, 032008 (2012).
- [5] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevD.100.031102> for equations relating the measured yields to the final results, tables of fractions as functions of  $p_T$ , plots of fractions versus  $\eta$ , and correlation matrices.
- [6] I. I. Bigi, T. Mannel, and N. Uraltsev, Semileptonic width ratios among beauty hadrons, *J. High Energy Phys.* **09** (2011) 012.
- [7] R. Aaij *et al.* (LHCb Collaboration), Measurement of the  $b$ -Quark Production Cross-Section in 7 and 13 TeV pp Collisions, *Phys. Rev. Lett.* **118**, 052002 (2017); Erratum, **119**, 169901(E) (2017).
- [8] G. Bonvicini *et al.* (CLEO Collaboration), Updated measurements of absolute  $D^+$  and  $D^0$  hadronic branching fractions and  $\sigma(e^+e^- \rightarrow D\bar{D})$  at  $E_{\text{cm}} = 3774$  MeV, *Phys. Rev. D* **89**, 072002 (2014); Erratum, **91**, 019903(E) (2015).
- [9] M. Ablikim *et al.* (BESIII Collaboration), Measurements of Absolute Hadronic Branching Fractions of the  $\Lambda_c^+$  Baryon, *Phys. Rev. Lett.* **116**, 052001 (2016).
- [10] A. Zupanc *et al.* (Belle Collaboration), Measurement of the Branching Fraction  $\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)$ , *Phys. Rev. Lett.* **113**, 042002 (2014).
- [11] M. Rudolph, An experimentalists guide to the semileptonic bottom to charm branching fractions, *Int. J. Mod. Phys. A* **33**, 1850176 (2018).
- [12] A. A. Alves, Jr. *et al.* (LHCb Collaboration), The LHCb detector at the LHC, *J. Instrum.* **3**, S08005 (2008).
- [13] R. Aaij *et al.* (LHCb Collaboration), LHCb detector performance, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [14] R. Aaij *et al.*, The LHCb trigger and its performance in 2011, *J. Instrum.* **8**, P04022 (2013).

- [15] V. V. Gligorov and M. Williams, Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree, *J. Instrum.* **8**, P02013 (2013).
- [16] T. Sjöstrand, S. Mrenna, and P. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* **178**, 852 (2008); T. Sjöstrand, S. Mrenna, and P. Skands, PYTHIA 6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [17] I. Belyaev *et al.*, Handling of the generation of primary events in Gauss, the LHCb simulation framework, *J. Phys. Conf. Ser.* **331**, 032047 (2011).
- [18] D. J. Lange, The EvtGen particle decay simulation package, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [19] P. Golonka and Z. Was, PHOTOS Monte Carlo: A precision tool for QED corrections in  $Z$  and  $W$  decays, *Eur. Phys. J. C* **45**, 97 (2006).
- [20] J. Allison *et al.* (Geant4 Collaboration), Geant4 developments and applications, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006); S. Agostinelli *et al.* (Geant4 Collaboration), Geant4: A simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [21] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranza, M. Pappagallo, and P. Robbe, The LHCb simulation application, Gauss: Design, evolution and experience, *J. Phys. Conf. Ser.* **331**, 032023 (2011).
- [22] V. M. Abazov *et al.* (D0 Collaboration), Measurement of the  $B_s^0$  Semileptonic Branching Ratio to an Orbitally Excited  $D_s^{*}$  State:  $\mathcal{B}(B_s^0 \rightarrow D_{s1}^-(2536)\mu^+\nu X)$ , *Phys. Rev. Lett.* **102**, 051801 (2009).
- [23] R. Aaij *et al.* (LHCb Collaboration), First observation of  $\bar{B}_s^0 \rightarrow D_{s2}^{*+} X \mu^- \bar{\nu}$  decays, *Phys. Lett. B* **698**, 14 (2011).
- [24] S. Tolk, J. Albrecht, F. Dettori, and A. Pellegrino, Data driven trigger efficiency determination at LHCb, CERN Reports No. CERN-LHCb-PUB-2014-039 and No. LHCb-PUB-2014-039, <https://cds.cern.ch/record/1701134/files/LHCb-PUB-2014-039.pdf>.
- [25] R. Aaij *et al.*, Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2, *EPJ Tech. Instrum.* **6**, 1 (2019).
- [26] P. del Amo Sanchez *et al.* (BABAR Collaboration), Observation of the Decay  $B^- \rightarrow D_s^{(*)+} K^- \ell^- \bar{\nu}_\ell$ , *Phys. Rev. Lett.* **107**, 041804 (2011).
- [27] R. Aaij *et al.* (LHCb Collaboration), Measurement of the fragmentation fraction ratio  $f_s/f_d$  and its dependence on  $B$  meson kinematics, *J. High Energy Phys.* **04** (2013) 001;  $f_s/f_d$  value updated in LHCb-CONF-2013-011.
- [28] R. Aaij *et al.* (LHCb Collaboration), Study of the kinematic dependences of  $\Lambda_b^0$  production in  $pp$  collisions and a measurement of the  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$  branching fraction, *J. High Energy Phys.* **08** (2014) 143.
- [29] R. Aaij *et al.* (LHCb Collaboration), Study of the production of  $\Lambda_b^0$  and  $\bar{B}^0$  hadrons in  $pp$  collisions and first measurement of the  $\Lambda_b^0 \rightarrow J/\psi p K^-$  branching fraction, *Chin. Phys. C* **40**, 011001 (2016).

R. Aaij,<sup>29</sup> C. Abellán Beteta,<sup>46</sup> B. Adeva,<sup>43</sup> M. Adinolfi,<sup>50</sup> C. A. Aidala,<sup>77</sup> Z. Ajaltouni,<sup>7</sup> S. Akar,<sup>61</sup> P. Albicocco,<sup>20</sup> J. Albrecht,<sup>12</sup> F. Alessio,<sup>44</sup> M. Alexander,<sup>55</sup> A. Alfonso Albergo,<sup>42</sup> G. Alkhazov,<sup>35</sup> P. Alvarez Cartelle,<sup>57</sup> A. A. Alves Jr.,<sup>43</sup> S. Amato,<sup>2</sup> S. Amerio,<sup>25</sup> Y. Amhis,<sup>9</sup> L. An,<sup>19</sup> L. Anderlini,<sup>19</sup> G. Andreassi,<sup>45</sup> M. Andreotti,<sup>18</sup> J. E. Andrews,<sup>62</sup> F. Archilli,<sup>29</sup> J. Arnau Romeu,<sup>8</sup> A. Artamonov,<sup>41</sup> M. Artuso,<sup>63</sup> K. Arzymatov,<sup>39</sup> E. Aslanides,<sup>8</sup> M. Atzeni,<sup>46</sup> B. Audurier,<sup>24</sup> S. Bachmann,<sup>14</sup> J. J. Back,<sup>52</sup> S. Baker,<sup>57</sup> V. Balagura,<sup>9,b</sup> W. Baldini,<sup>18</sup> A. Baranov,<sup>39</sup> R. J. Barlow,<sup>58</sup> G. C. Barrand,<sup>9</sup> S. Barsuk,<sup>9</sup> W. Barter,<sup>57</sup> M. Bartolini,<sup>21</sup> F. Baryshnikov,<sup>73</sup> V. Batozskaya,<sup>33</sup> B. Batsukh,<sup>63</sup> A. Battig,<sup>12</sup> V. Battista,<sup>45</sup> A. Bay,<sup>45</sup> J. Beddow,<sup>55</sup> F. Bedeschi,<sup>26</sup> I. Bediaga,<sup>1</sup> A. Beiter,<sup>63</sup> L. J. Bel,<sup>29</sup> S. Belin,<sup>24</sup> N. Belyi,<sup>4</sup> V. Bellee,<sup>45</sup> N. Belloli,<sup>22,i</sup> K. Belous,<sup>41</sup> I. Belyaev,<sup>36</sup> G. Bencivenni,<sup>20</sup> E. Ben-Haim,<sup>10</sup> S. Benson,<sup>29</sup> S. Beranek,<sup>11</sup> A. Berezhnoy,<sup>37</sup> R. Bernet,<sup>46</sup> D. Berninghoff,<sup>14</sup> E. Bertholet,<sup>10</sup> A. Bertolin,<sup>25</sup> C. Betancourt,<sup>46</sup> F. Betti,<sup>17,44</sup> M. O. Bettler,<sup>51</sup> I. A. Bezshyiko,<sup>46</sup> S. Bhasin,<sup>50</sup> J. Bhom,<sup>31</sup> M. S. Bieker,<sup>12</sup> S. Bifani,<sup>49</sup> P. Billoir,<sup>10</sup> A. Birnkraut,<sup>12</sup> A. Bizzeti,<sup>19,u</sup> M. Bjørn,<sup>59</sup> M. P. Blago,<sup>44</sup> T. Blake,<sup>52</sup> F. Blanc,<sup>45</sup> S. Blusk,<sup>63</sup> D. Bobulska,<sup>55</sup> V. Bocci,<sup>28</sup> O. Boente Garcia,<sup>43</sup> T. Boettcher,<sup>60</sup> A. Bondar,<sup>40,x</sup> N. Bondar,<sup>35</sup> S. Borghi,<sup>58,44</sup> M. Borisyak,<sup>39</sup> M. Borsato,<sup>14</sup> M. Boubdir,<sup>11</sup> T. J. V. Bowcock,<sup>56</sup> C. Bozzi,<sup>18,44</sup> S. Braun,<sup>14</sup> M. Brodski,<sup>44</sup> J. Brodzicka,<sup>31</sup> A. Brossa Gonzalo,<sup>52</sup> D. Brundu,<sup>24,44</sup> E. Buchanan,<sup>50</sup> A. Buonauro,<sup>46</sup> C. Burr,<sup>58</sup> A. Bursche,<sup>24</sup> J. Buytaert,<sup>44</sup> W. Byczynski,<sup>44</sup> S. Cadeddu,<sup>24</sup> H. Cai,<sup>67</sup> R. Calabrese,<sup>18,g</sup> R. Calladine,<sup>49</sup> M. Calvi,<sup>22,i</sup> M. Calvo Gomez,<sup>42,m</sup> A. Camboni,<sup>42,m</sup> P. Campana,<sup>20</sup> D. H. Campora Perez,<sup>44</sup> L. Capriotti,<sup>17,e</sup> A. Carbone,<sup>17,e</sup> G. Carboni,<sup>27</sup> R. Cardinale,<sup>21</sup> A. Cardini,<sup>24</sup> P. Carniti,<sup>22,i</sup> K. Carvalho Akiba,<sup>2</sup> G. Casse,<sup>56</sup> M. Cattaneo,<sup>44</sup> G. Cavallero,<sup>21</sup> R. Cenci,<sup>26,p</sup> D. Chamont,<sup>9</sup> M. G. Chapman,<sup>50</sup> M. Charles,<sup>10</sup> Ph. Charpentier,<sup>44</sup> G. Chatzikonstantinidis,<sup>49</sup> M. Chefdeville,<sup>6</sup> V. Chekalina,<sup>39</sup> C. Chen,<sup>3</sup> S. Chen,<sup>24</sup> S.-G. Chitic,<sup>44</sup> V. Chobanova,<sup>43</sup> M. Chrzaszcz,<sup>44</sup> A. Chubykin,<sup>35</sup> P. Ciambone,<sup>20</sup> X. Cid Vidal,<sup>43</sup> G. Ciezarek,<sup>44</sup> F. Cindolo,<sup>17</sup> P. E. L. Clarke,<sup>54</sup> M. Clemencic,<sup>44</sup> H. V. Cliff,<sup>51</sup> J. Closier,<sup>44</sup> V. Coco,<sup>44</sup> J. A. B. Coelho,<sup>9</sup> J. Cogan,<sup>8</sup> E. Cogneras,<sup>7</sup> L. Cojocariu,<sup>34</sup> P. Collins,<sup>44</sup> T. Colombo,<sup>44</sup> A. Comerma-Montells,<sup>14</sup> A. Contu,<sup>24</sup> G. Coombs,<sup>44</sup> S. Coquereau,<sup>42</sup> G. Corti,<sup>44</sup> M. Corvo,<sup>18,g</sup> C. M. Costa Sobral,<sup>52</sup> B. Couturier,<sup>44</sup> G. A. Cowan,<sup>54</sup> D. C. Craik,<sup>60</sup> A. Crocombe,<sup>52</sup> M. Cruz Torres,<sup>1</sup> R. Currie,<sup>54</sup> F. Da Cunha Marinho,<sup>2</sup> C. L. Da Silva,<sup>78</sup> E. Dall'Occo,<sup>29</sup> J. Dalseno,<sup>43,v</sup> C. D'Ambrosio,<sup>44</sup> A. Danilina,<sup>36</sup> P. d'Argent,<sup>14</sup> A. Davis,<sup>58</sup> O. De Aguiar Francisco,<sup>44</sup> K. De Bruyn,<sup>44</sup> S. De Capua,<sup>58</sup> M. De Cian,<sup>45</sup> J. M. De Miranda,<sup>1</sup>



L. De Paula,<sup>2</sup> M. De Serio,<sup>16,d</sup> P. De Simone,<sup>20</sup> J. A. de Vries,<sup>29</sup> C. T. Dean,<sup>55</sup> W. Dean,<sup>77</sup> D. Decamp,<sup>6</sup> L. Del Buono,<sup>10</sup> B. Delaney,<sup>51</sup> H.-P. Dembinski,<sup>13</sup> M. Demmer,<sup>12</sup> A. Dendek,<sup>32</sup> D. Derkach,<sup>74</sup> O. Deschamps,<sup>7</sup> F. Desse,<sup>9</sup> F. Dettori,<sup>56</sup> B. Dey,<sup>68</sup> A. Di Canto,<sup>44</sup> P. Di Nezza,<sup>20</sup> S. Didenko,<sup>73</sup> H. Dijkstra,<sup>44</sup> F. Dordei,<sup>24</sup> M. Dorigo,<sup>44,y</sup> A. C. dos Reis,<sup>1</sup> A. Dosit Suárez,<sup>43</sup> L. Douglas,<sup>55</sup> A. Dovbnya,<sup>47</sup> K. Dreimanis,<sup>56</sup> L. Dufour,<sup>29</sup> G. Dujany,<sup>10</sup> P. Durante,<sup>44</sup> J. M. Durham,<sup>78</sup> D. Dutta,<sup>58</sup> R. Dzhelyadin,<sup>41,t</sup> M. Dziewiecki,<sup>14</sup> A. Dziurda,<sup>31</sup> A. Dzyuba,<sup>35</sup> S. Easo,<sup>53</sup> U. Egede,<sup>57</sup> V. Egorychev,<sup>36</sup> S. Eidelman,<sup>40,x</sup> S. Eisenhardt,<sup>54</sup> U. Eitschberger,<sup>12</sup> R. Ekelhof,<sup>12</sup> L. Eklund,<sup>55</sup> S. Ely,<sup>63</sup> A. Ene,<sup>34</sup> S. Escher,<sup>11</sup> S. Esen,<sup>29</sup> T. Evans,<sup>61</sup> A. Falabella,<sup>17</sup> C. Färber,<sup>44</sup> N. Farley,<sup>49</sup> S. Farry,<sup>56</sup> D. Fazzini,<sup>22,44,i</sup> M. Féo,<sup>44</sup> P. Fernandez Declara,<sup>44</sup> A. Fernandez Prieto,<sup>43</sup> F. Ferrari,<sup>17,e</sup> L. Ferreira Lopes,<sup>45</sup> F. Ferreira Rodrigues,<sup>2</sup> M. Ferro-Luzzi,<sup>44</sup> S. Filippov,<sup>38</sup> R. A. Fini,<sup>16</sup> M. Fiorini,<sup>18,g</sup> M. Firlej,<sup>32</sup> C. Fitzpatrick,<sup>45</sup> T. Fiutowski,<sup>32</sup> F. Fleuret,<sup>9,b</sup> M. Fontana,<sup>44</sup> F. Fontanelli,<sup>21,h</sup> R. Forty,<sup>44</sup> V. Franco Lima,<sup>56</sup> M. Frank,<sup>44</sup> C. Frei,<sup>44</sup> J. Fu,<sup>23,q</sup> W. Funk,<sup>44</sup> E. Gabriel,<sup>54</sup> A. Gallas Torreira,<sup>43</sup> D. Galli,<sup>17,e</sup> S. Gallorini,<sup>25</sup> S. Gambetta,<sup>54</sup> Y. Gan,<sup>3</sup> M. Gandelman,<sup>2</sup> P. Gandini,<sup>23</sup> Y. Gao,<sup>3</sup> L. M. Garcia Martin,<sup>76</sup> J. García Pardiñas,<sup>46</sup> B. Garcia Plana,<sup>43</sup> J. Garra Tico,<sup>51</sup> L. Garrido,<sup>42</sup> D. Gascon,<sup>42</sup> C. Gaspar,<sup>44</sup> G. Gazzoni,<sup>7</sup> D. Gerick,<sup>14</sup> E. Gersabeck,<sup>58</sup> M. Gersabeck,<sup>58</sup> T. Gershon,<sup>52</sup> D. Gerstel,<sup>8</sup> Ph. Ghez,<sup>6</sup> V. Gibson,<sup>51</sup> O. G. Girard,<sup>45</sup> P. Gironella Gironell,<sup>42</sup> L. Giubega,<sup>34</sup> K. Gizdov,<sup>54</sup> V. V. Gligorov,<sup>10</sup> C. Göbel,<sup>65</sup> D. Golubkov,<sup>36</sup> A. Golutvin,<sup>57,73</sup> A. Gomes,<sup>1,a</sup> I. V. Gorelov,<sup>37</sup> C. Gotti,<sup>22,i</sup> E. Govorkova,<sup>29</sup> J. P. Grabowski,<sup>14</sup> R. Graciani Diaz,<sup>42</sup> L. A. Granado Cardoso,<sup>44</sup> E. Graugés,<sup>42</sup> E. Graverini,<sup>46</sup> G. Graziani,<sup>19</sup> A. Grecu,<sup>34</sup> R. Greim,<sup>29</sup> P. Griffith,<sup>24</sup> L. Grillo,<sup>58</sup> L. Gruber,<sup>44</sup> B. R. Gruberg Cazon,<sup>59</sup> O. Grünberg,<sup>70</sup> C. Gu,<sup>3</sup> E. Gushchin,<sup>38</sup> A. Guth,<sup>11</sup> Yu. Guz,<sup>41,44</sup> T. Gys,<sup>44</sup> T. Hadavizadeh,<sup>59</sup> C. Hadjivasiliou,<sup>7</sup> G. Haefeli,<sup>45</sup> C. Haen,<sup>44</sup> S. C. Haines,<sup>51</sup> B. Hamilton,<sup>62</sup> X. Han,<sup>14</sup> T. H. Hancock,<sup>59</sup> S. Hansmann-Menzemer,<sup>14</sup> N. Harnew,<sup>59</sup> T. Harrison,<sup>56</sup> C. Hasse,<sup>44</sup> M. Hatch,<sup>44</sup> J. He,<sup>4</sup> M. Hecker,<sup>57</sup> K. Heinicke,<sup>12</sup> A. Heister,<sup>12</sup> K. Hennessy,<sup>56</sup> L. Henry,<sup>76</sup> M. Heß,<sup>70</sup> J. Heuel,<sup>11</sup> A. Hicheur,<sup>64</sup> R. Hidalgo Charman,<sup>58</sup> D. Hill,<sup>59</sup> M. Hilton,<sup>58</sup> P. H. Hopchev,<sup>45</sup> J. Hu,<sup>14</sup> W. Hu,<sup>68</sup> W. Huang,<sup>4</sup> Z. C. Huard,<sup>61</sup> W. Hulsbergen,<sup>29</sup> T. Humair,<sup>57</sup> M. Hushchyn,<sup>74</sup> D. Hutchcroft,<sup>56</sup> D. Hynds,<sup>29</sup> P. Ibis,<sup>12</sup> M. Idzik,<sup>32</sup> P. Ilten,<sup>49</sup> A. Inglessi,<sup>35</sup> A. Inyakin,<sup>41</sup> K. Ivshin,<sup>35</sup> R. Jacobsson,<sup>44</sup> S. Jakobsen,<sup>44</sup> J. Jalocha,<sup>59</sup> E. Jans,<sup>29</sup> B. K. Jashal,<sup>76</sup> A. Jawahery,<sup>62</sup> F. Jiang,<sup>3</sup> M. John,<sup>59</sup> D. Johnson,<sup>44</sup> C. R. Jones,<sup>51</sup> C. Joram,<sup>44</sup> B. Jost,<sup>44</sup> N. Jurik,<sup>59</sup> S. Kandybei,<sup>47</sup> M. Karacson,<sup>44</sup> J. M. Kariuki,<sup>50</sup> S. Karodia,<sup>55</sup> N. Kazeev,<sup>74</sup> M. Kecke,<sup>14</sup> F. Keizer,<sup>51</sup> M. Kelsey,<sup>63</sup> M. Kenzie,<sup>51</sup> T. Ketel,<sup>30</sup> E. Khairullin,<sup>39</sup> B. Khanji,<sup>44</sup> C. Khurewathanakul,<sup>45</sup> K. E. Kim,<sup>63</sup> T. Kim,<sup>11</sup> V. S. Kirsabom,<sup>45</sup> S. Klaver,<sup>20</sup> K. Klimaszewski,<sup>33</sup> T. Klimovich,<sup>13</sup> S. Koliiev,<sup>48</sup> M. Kolpin,<sup>14</sup> R. Kopečna,<sup>14</sup> P. Koppenburg,<sup>29</sup> I. Kostiuik,<sup>29,48</sup> S. Kotriakhova,<sup>35</sup> M. Kozeiha,<sup>7</sup> L. Kravchuk,<sup>38</sup> M. Krepis,<sup>52</sup> F. Kress,<sup>57</sup> P. Krokovny,<sup>40,x</sup> W. Krupa,<sup>32</sup> W. Krzemien,<sup>33</sup> W. Kucewicz,<sup>31,l</sup> M. Kucharczyk,<sup>31</sup> V. Kudryavtsev,<sup>40,x</sup> A. K. Kuonen,<sup>45</sup> T. Kvaratskheliya,<sup>36,44</sup> D. Lacarrere,<sup>44</sup> G. Lafferty,<sup>58</sup> A. Lai,<sup>24</sup> D. Lancierini,<sup>46</sup> G. Lanfranchi,<sup>20</sup> C. Langenbruch,<sup>11</sup> T. Latham,<sup>52</sup> C. Lazzeroni,<sup>49</sup> R. Le Gac,<sup>8</sup> R. Lefèvre,<sup>7</sup> A. Leflat,<sup>37</sup> F. Lemaître,<sup>44</sup> O. Leroy,<sup>8</sup> T. Lesiak,<sup>31</sup> B. Leverington,<sup>14</sup> P.-R. Li,<sup>4,ab</sup> Y. Li,<sup>5</sup> Z. Li,<sup>63</sup> X. Liang,<sup>63</sup> T. Likhomanenko,<sup>72</sup> R. Lindner,<sup>44</sup> F. Lionetto,<sup>46</sup> V. Lisovskyi,<sup>9</sup> G. Liu,<sup>66</sup> X. Liu,<sup>3</sup> D. Loh,<sup>52</sup> A. Loi,<sup>24</sup> I. Longstaff,<sup>55</sup> J. H. Lopes,<sup>2</sup> G. Loustau,<sup>46</sup> G. H. Lovell,<sup>51</sup> D. Lucchesi,<sup>25,o</sup> M. Lucio Martinez,<sup>43</sup> Y. Luo,<sup>3</sup> A. Lupato,<sup>25</sup> E. Luppi,<sup>18,g</sup> O. Lupton,<sup>44</sup> A. Lusiani,<sup>26</sup> X. Lyu,<sup>4</sup> F. Machefert,<sup>9</sup> F. Maciuc,<sup>34</sup> V. Macko,<sup>45</sup> P. Mackowiak,<sup>12</sup> S. Maddrell-Mander,<sup>50</sup> O. Maev,<sup>35,44</sup> K. Maguire,<sup>58</sup> D. Maisuzenko,<sup>35</sup> M. W. Majewski,<sup>32</sup> S. Malde,<sup>59</sup> B. Malecki,<sup>44</sup> A. Malinin,<sup>72</sup> T. Maltsev,<sup>40,x</sup> H. Malygina,<sup>14</sup> G. Manca,<sup>24,f</sup> G. Mancinelli,<sup>8</sup> D. Marangotto,<sup>23,q</sup> J. Maratas,<sup>7,w</sup> J. F. Marchand,<sup>6</sup> U. Marconi,<sup>17</sup> C. Marin Benito,<sup>9</sup> M. Marinangeli,<sup>45</sup> P. Marino,<sup>45</sup> J. Marks,<sup>14</sup> P. J. Marshall,<sup>56</sup> G. Martellotti,<sup>28</sup> M. Martinelli,<sup>44</sup> D. Martinez Santos,<sup>43</sup> F. Martinez Vidal,<sup>76</sup> A. Massafferri,<sup>1</sup> M. Materok,<sup>11</sup> R. Matev,<sup>44</sup> A. Mathad,<sup>52</sup> Z. Mathe,<sup>44</sup> C. Matteuzzi,<sup>22</sup> K. R. Mattioli,<sup>77</sup> A. Mauri,<sup>46</sup> E. Maurice,<sup>9,b</sup> B. Maurin,<sup>45</sup> M. McCann,<sup>57,44</sup> A. McNab,<sup>58</sup> R. McNulty,<sup>15</sup> J. V. Mead,<sup>56</sup> B. Meadows,<sup>61</sup> C. Meaux,<sup>8</sup> N. Meinert,<sup>70</sup> D. Melnychuk,<sup>33</sup> M. Merk,<sup>29</sup> A. Merli,<sup>23,q</sup> E. Michielin,<sup>25</sup> D. A. Milanés,<sup>69</sup> E. Millard,<sup>52</sup> M.-N. Minard,<sup>6</sup> L. Minzoni,<sup>18,g</sup> D. S. Mitzel,<sup>14</sup> A. Mödden,<sup>12</sup> A. Mogini,<sup>10</sup> R. D. Moise,<sup>57</sup> T. Mombächer,<sup>12</sup> I. A. Monroy,<sup>69</sup> S. Monteil,<sup>7</sup> M. Morandin,<sup>25</sup> G. Morello,<sup>20</sup> M. J. Morello,<sup>26,t</sup> O. Morgunova,<sup>72</sup> J. Moron,<sup>32</sup> A. B. Morris,<sup>8</sup> R. Mountain,<sup>63</sup> F. Muheim,<sup>54</sup> M. Mukherjee,<sup>68</sup> M. Mulder,<sup>29</sup> D. Müller,<sup>44</sup> J. Müller,<sup>12</sup> K. Müller,<sup>46</sup> V. Müller,<sup>12</sup> C. H. Murphy,<sup>59</sup> D. Murray,<sup>58</sup> P. Naik,<sup>50</sup> T. Nakada,<sup>45</sup> R. Nandakumar,<sup>53</sup> A. Nandi,<sup>59</sup> T. Nanut,<sup>45</sup> I. Nasteva,<sup>2</sup> M. Needham,<sup>54</sup> N. Neri,<sup>23,q</sup> S. Neubert,<sup>14</sup> N. Neufeld,<sup>44</sup> R. Newcombe,<sup>57</sup> T. D. Nguyen,<sup>45</sup> C. Nguyen-Mau,<sup>45,n</sup> S. Nieswand,<sup>11</sup> R. Niet,<sup>12</sup> N. Nikitin,<sup>37</sup> A. Nogay,<sup>72</sup> N. S. Nolte,<sup>44</sup> A. Oblakowska-Mucha,<sup>32</sup> V. Obraztsov,<sup>41</sup> S. Ogilvy,<sup>55</sup> D. P. O'Hanlon,<sup>17</sup> R. Oldeman,<sup>24,f</sup> C. J. G. Onderwater,<sup>71</sup> A. Ossowska,<sup>31</sup> J. M. Otalora Goicochea,<sup>2</sup> T. Ovsianikova,<sup>36</sup> P. Owen,<sup>46</sup> A. Oyanguren,<sup>76</sup> P. R. Pais,<sup>45</sup> T. Pajero,<sup>26,t</sup> A. Palano,<sup>16</sup> M. Palutan,<sup>20</sup> G. Panshin,<sup>75</sup> A. Papanestis,<sup>53</sup> M. Pappagallo,<sup>54</sup> L. L. Pappalardo,<sup>18,g</sup> W. Parker,<sup>62</sup> C. Parkes,<sup>58,44</sup> G. Passaleva,<sup>19,44</sup> A. Pastore,<sup>16</sup> M. Patel,<sup>57</sup> C. Patrignani,<sup>17,e</sup> A. Pearce,<sup>44</sup> A. Pellegrino,<sup>29</sup> G. Penso,<sup>28</sup> M. Pepe Altarelli,<sup>44</sup> S. Perazzini,<sup>44</sup> D. Pereima,<sup>36</sup> P. Perret,<sup>7</sup> L. Pescatore,<sup>45</sup> K. Petridis,<sup>50</sup> A. Petrolini,<sup>21,h</sup> A. Petrov,<sup>72</sup> S. Petrucci,<sup>54</sup>

M. Petruzzo,<sup>23,q</sup> B. Pietrzyk,<sup>6</sup> G. Pietrzyk,<sup>45</sup> M. Pikies,<sup>31</sup> M. Pili,<sup>59</sup> D. Pinci,<sup>28</sup> J. Pinzino,<sup>44</sup> F. Pisani,<sup>44</sup> A. Piucci,<sup>14</sup> V. Placinta,<sup>34</sup> S. Playfer,<sup>54</sup> J. Plews,<sup>49</sup> M. Plo Casasus,<sup>43</sup> F. Polci,<sup>10</sup> M. Poli Lener,<sup>20</sup> A. Poluektov,<sup>8</sup> N. Polukhina,<sup>73,c</sup> I. Polyakov,<sup>63</sup> E. Polycarpo,<sup>2</sup> G. J. Pomery,<sup>50</sup> S. Ponce,<sup>44</sup> A. Popov,<sup>41</sup> D. Popov,<sup>49,13</sup> S. Poslavskii,<sup>41</sup> E. Price,<sup>50</sup> J. Prisciandaro,<sup>43</sup> C. Prouve,<sup>43</sup> V. Pugatch,<sup>48</sup> A. Puig Navarro,<sup>46</sup> H. Pullen,<sup>59</sup> G. Punzi,<sup>26,p</sup> W. Qian,<sup>4</sup> J. Qin,<sup>4</sup> R. Quagliani,<sup>10</sup> B. Quintana,<sup>7</sup> N. V. Raab,<sup>15</sup> B. Rachwal,<sup>32</sup> J. H. Rademacker,<sup>50</sup> M. Rama,<sup>26</sup> M. Ramos Pernas,<sup>43</sup> M. S. Rangel,<sup>2</sup> F. Ratnikov,<sup>39,74</sup> G. Raven,<sup>30</sup> M. Ravonel Salzgeber,<sup>44</sup> M. Reboud,<sup>6</sup> F. Redi,<sup>45</sup> S. Reichert,<sup>12</sup> F. Reiss,<sup>10</sup> C. Remon Alepuz,<sup>76</sup> Z. Ren,<sup>3</sup> V. Renaudin,<sup>59</sup> S. Ricciardi,<sup>53</sup> S. Richards,<sup>50</sup> K. Rinnert,<sup>56</sup> P. Robbe,<sup>9</sup> A. Robert,<sup>10</sup> A. B. Rodrigues,<sup>45</sup> E. Rodrigues,<sup>61</sup> J. A. Rodriguez Lopez,<sup>69</sup> M. Roehrken,<sup>44</sup> S. Roiser,<sup>44</sup> A. Rollings,<sup>59</sup> V. Romanovskiy,<sup>41</sup> A. Romero Vidal,<sup>43</sup> J. D. Roth,<sup>77</sup> M. Rotondo,<sup>20</sup> M. S. Rudolph,<sup>63</sup> T. Ruf,<sup>44</sup> J. Ruiz Vidal,<sup>76</sup> J. J. Saborido Silva,<sup>43</sup> N. Sagidova,<sup>35</sup> B. Saitta,<sup>24,f</sup> V. Salustino Guimaraes,<sup>65</sup> C. Sanchez Gras,<sup>29</sup> C. Sanchez Mayordomo,<sup>76</sup> B. Sanmartin Sedes,<sup>43</sup> R. Santacesaria,<sup>28</sup> C. Santamarina Rios,<sup>43</sup> M. Santimaria,<sup>20,44</sup> E. Santovetti,<sup>27,j</sup> G. Sarpis,<sup>58</sup> A. Sarti,<sup>20,k</sup> C. Satriano,<sup>28,s</sup> A. Satta,<sup>27</sup> M. Saur,<sup>4</sup> D. Savrina,<sup>36,37</sup> S. Schael,<sup>11</sup> M. Schellenberg,<sup>12</sup> M. Schiller,<sup>55</sup> H. Schindler,<sup>44</sup> M. Schmelling,<sup>13</sup> T. Schmelzer,<sup>12</sup> B. Schmidt,<sup>44</sup> O. Schneider,<sup>45</sup> A. Schopper,<sup>44</sup> H. F. Schreiner,<sup>61</sup> M. Schubiger,<sup>45</sup> S. Schulte,<sup>45</sup> M. H. Schune,<sup>9</sup> R. Schwemmer,<sup>44</sup> B. Sciascia,<sup>20</sup> A. Sciubba,<sup>28,k</sup> A. Semennikov,<sup>36</sup> E. S. Sepulveda,<sup>10</sup> A. Sergi,<sup>49</sup> N. Serra,<sup>46</sup> J. Serrano,<sup>8</sup> L. Sestini,<sup>25</sup> A. Seuthe,<sup>12</sup> P. Seyfert,<sup>44</sup> M. Shapkin,<sup>41</sup> T. Shears,<sup>56</sup> L. Shekhtman,<sup>40,x</sup> V. Shevchenko,<sup>72</sup> E. Shmanin,<sup>73</sup> B. G. Siddi,<sup>18</sup> R. Silva Coutinho,<sup>46</sup> L. Silva de Oliveira,<sup>2</sup> G. Simi,<sup>25,o</sup> S. Simone,<sup>16,d</sup> I. Skiba,<sup>18</sup> N. Skidmore,<sup>14</sup> T. Skwarnicki,<sup>63</sup> M. W. Slater,<sup>49</sup> J. G. Smeaton,<sup>51</sup> E. Smith,<sup>11</sup> I. T. Smith,<sup>54</sup> M. Smith,<sup>57</sup> M. Soares,<sup>17</sup> I. Soares Lavra,<sup>1</sup> M. D. Sokoloff,<sup>61</sup> F. J. P. Soler,<sup>55</sup> B. Souza De Paula,<sup>2</sup> B. Spaan,<sup>12</sup> E. Spadaro Norella,<sup>23,q</sup> P. Spradlin,<sup>55</sup> F. Stagni,<sup>44</sup> M. Stahl,<sup>14</sup> S. Stahl,<sup>44</sup> P. Stefko,<sup>45</sup> S. Stefkova,<sup>57</sup> O. Steinkamp,<sup>46</sup> S. Stemmler,<sup>14</sup> O. Stenyakin,<sup>41</sup> M. Stepanova,<sup>35</sup> H. Stevens,<sup>12</sup> A. Stocchi,<sup>9</sup> S. Stone,<sup>63</sup> B. Storaci,<sup>46</sup> S. Stracka,<sup>26</sup> M. E. Stramaglia,<sup>45</sup> M. Straticiu,<sup>34</sup> U. Straumann,<sup>46</sup> S. Strovkov,<sup>75</sup> J. Sun,<sup>3</sup> L. Sun,<sup>67</sup> Y. Sun,<sup>62</sup> K. Swientek,<sup>32</sup> A. Szabelski,<sup>33</sup> T. Szumlak,<sup>32</sup> M. Szymanski,<sup>4</sup> Z. Tang,<sup>3</sup> T. Tekampe,<sup>12</sup> G. Tellarini,<sup>18</sup> F. Teubert,<sup>44</sup> E. Thomas,<sup>44</sup> M. J. Tilley,<sup>57</sup> V. Tisserand,<sup>7</sup> S. T'Jampens,<sup>6</sup> M. Tobin,<sup>32</sup> S. Tolck,<sup>44</sup> L. Tomassetti,<sup>18,g</sup> D. Tonelli,<sup>26</sup> D. Y. Tou,<sup>10</sup> R. Tourinho Jadallah Aoude,<sup>1</sup> E. Tournefier,<sup>6</sup> M. Traill,<sup>55</sup> M. T. Tran,<sup>45</sup> A. Trisovic,<sup>51</sup> A. Tsaregorodtsev,<sup>8</sup> G. Tuci,<sup>26,p</sup> A. Tully,<sup>51</sup> N. Tuning,<sup>29,44</sup> A. Ukleja,<sup>33</sup> A. Usachov,<sup>9</sup> A. Ustyuzhanin,<sup>39,74</sup> U. Uwer,<sup>14</sup> A. Vagner,<sup>75</sup> V. Vagnoni,<sup>17</sup> A. Valassi,<sup>44</sup> S. Valat,<sup>44</sup> G. Valenti,<sup>17</sup> M. van Beuzekom,<sup>29</sup> E. van Herwijnen,<sup>44</sup> J. van Tilburg,<sup>29</sup> M. van Veghel,<sup>29</sup> R. Vazquez Gomez,<sup>44</sup> P. Vazquez Regueiro,<sup>43</sup> C. Vázquez Sierra,<sup>29</sup> S. Vecchi,<sup>18</sup> J. J. Velthuis,<sup>50</sup> M. Veltri,<sup>19,r</sup> A. Venkateswaran,<sup>63</sup> M. Vernet,<sup>7</sup> M. Veronesi,<sup>29</sup> M. Vesterinen,<sup>52</sup> J. V. Viana Barbosa,<sup>44</sup> D. Vieira,<sup>4</sup> M. Vieites Diaz,<sup>43</sup> H. Viemann,<sup>70</sup> X. Vilasis-Cardona,<sup>42,m</sup> A. Vitkovskiy,<sup>29</sup> M. Vitti,<sup>51</sup> V. Volkov,<sup>37</sup> A. Vollhardt,<sup>46</sup> D. Vom Bruch,<sup>10</sup> B. Voneki,<sup>44</sup> A. Vorobyev,<sup>35</sup> V. Vorobyev,<sup>40,x</sup> N. Voropaev,<sup>35</sup> R. Waldi,<sup>70</sup> J. Walsh,<sup>26</sup> J. Wang,<sup>5</sup> M. Wang,<sup>3</sup> Y. Wang,<sup>68</sup> Z. Wang,<sup>46</sup> D. R. Ward,<sup>51</sup> H. M. Wark,<sup>56</sup> N. K. Watson,<sup>49</sup> D. Websdale,<sup>57</sup> A. Weiden,<sup>46</sup> C. Weisser,<sup>60</sup> M. Whitehead,<sup>11</sup> G. Wilkinson,<sup>59</sup> M. Wilkinson,<sup>63</sup> I. Williams,<sup>51</sup> M. Williams,<sup>60</sup> M. R. J. Williams,<sup>58</sup> T. Williams,<sup>49</sup> F. F. Wilson,<sup>53</sup> M. Winn,<sup>9</sup> W. Wislicki,<sup>33</sup> M. Witek,<sup>31</sup> G. Wormser,<sup>9</sup> S. A. Wotton,<sup>51</sup> K. Wyllie,<sup>44</sup> D. Xiao,<sup>68</sup> Y. Xie,<sup>68</sup> A. Xu,<sup>3</sup> M. Xu,<sup>68</sup> Q. Xu,<sup>4</sup> Z. Xu,<sup>6</sup> Z. Xu,<sup>3</sup> Z. Yang,<sup>3</sup> Z. Yang,<sup>62</sup> Y. Yao,<sup>63</sup> L. E. Yeomans,<sup>56</sup> H. Yin,<sup>68</sup> J. Yu,<sup>68,aa</sup> X. Yuan,<sup>63</sup> O. Yushchenko,<sup>41</sup> K. A. Zarebski,<sup>49</sup> M. Zavertyaev,<sup>13,c</sup> D. Zhang,<sup>68</sup> L. Zhang,<sup>3</sup> W. C. Zhang,<sup>3,z</sup> Y. Zhang,<sup>44</sup> A. Zhelezov,<sup>14</sup> Y. Zheng,<sup>4</sup> X. Zhu,<sup>3</sup> V. Zhukov,<sup>11,37</sup> J. B. Zonneveld,<sup>54</sup> and S. Zucchelli<sup>17,e</sup>

(LHCb Collaboration)

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China<sup>4</sup>University of Chinese Academy of Sciences, Beijing, China<sup>5</sup>Institute of High Energy Physics (ihep), Beijing, China<sup>6</sup>University Grenoble Alpes, University Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France<sup>7</sup>Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France<sup>8</sup>Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France<sup>9</sup>LAL, University Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France<sup>10</sup>LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France<sup>11</sup>I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany<sup>12</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany<sup>13</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany<sup>14</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

- <sup>15</sup>*School of Physics, University College Dublin, Dublin, Ireland*
- <sup>16</sup>*INFN Sezione di Bari, Bari, Italy*
- <sup>17</sup>*INFN Sezione di Bologna, Bologna, Italy*
- <sup>18</sup>*INFN Sezione di Ferrara, Ferrara, Italy*
- <sup>19</sup>*INFN Sezione di Firenze, Firenze, Italy*
- <sup>20</sup>*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- <sup>21</sup>*INFN Sezione di Genova, Genova, Italy*
- <sup>22</sup>*INFN Sezione di Milano-Bicocca, Milano, Italy*
- <sup>23</sup>*INFN Sezione di Milano, Milano, Italy*
- <sup>24</sup>*INFN Sezione di Cagliari, Monserrato, Italy*
- <sup>25</sup>*INFN Sezione di Padova, Padova, Italy*
- <sup>26</sup>*INFN Sezione di Pisa, Pisa, Italy*
- <sup>27</sup>*INFN Sezione di Roma Tor Vergata, Roma, Italy*
- <sup>28</sup>*INFN Sezione di Roma La Sapienza, Roma, Italy*
- <sup>29</sup>*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- <sup>30</sup>*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- <sup>31</sup>*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- <sup>32</sup>*AGH—University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- <sup>33</sup>*National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- <sup>34</sup>*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- <sup>35</sup>*Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- <sup>36</sup>*Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- <sup>37</sup>*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>38</sup>*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*
- <sup>39</sup>*Yandex School of Data Analysis, Moscow, Russia*
- <sup>40</sup>*Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- <sup>41</sup>*Institute for High Energy Physics (IHEP), Protvino, Russia*
- <sup>42</sup>*ICCUB, Universitat de Barcelona, Barcelona, Spain*
- <sup>43</sup>*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>44</sup>*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>45</sup>*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>46</sup>*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>47</sup>*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>48</sup>*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>49</sup>*University of Birmingham, Birmingham, United Kingdom*
- <sup>50</sup>*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>51</sup>*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>52</sup>*Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>53</sup>*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>54</sup>*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>55</sup>*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>56</sup>*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>57</sup>*Imperial College London, London, United Kingdom*
- <sup>58</sup>*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>59</sup>*Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>60</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
- <sup>61</sup>*University of Cincinnati, Cincinnati, Ohio, USA*
- <sup>62</sup>*University of Maryland, College Park, Maryland, USA*
- <sup>63</sup>*Syracuse University, Syracuse, New York, USA*
- <sup>64</sup>*Laboratory of Mathematical and Subatomic Physics, Constantine, Algeria*  
[associated to Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil]
- <sup>65</sup>*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil*  
[associated to Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil]
- <sup>66</sup>*South China Normal University, Guangzhou, China*  
(associated to Center for High Energy Physics, Tsinghua University, Beijing, China)
- <sup>67</sup>*School of Physics and Technology, Wuhan University, Wuhan, China*  
(associated to Center for High Energy Physics, Tsinghua University, Beijing, China)

<sup>68</sup>*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China  
(associated to Center for High Energy Physics, Tsinghua University, Beijing, China)*

<sup>69</sup>*Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia  
(associated to LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France)*

<sup>70</sup>*Institut für Physik, Universität Rostock, Rostock, Germany  
(associated to Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)*

<sup>71</sup>*Van Swinderen Institute, University of Groningen, Groningen, Netherlands  
(associated to Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)*

<sup>72</sup>*National Research Centre Kurchatov Institute, Moscow, Russia  
[associated to Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia]*

<sup>73</sup>*National University of Science and Technology “MISIS,” Moscow, Russia  
[associated to Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia]*

<sup>74</sup>*National Research University Higher School of Economics, Moscow, Russia  
(associated to Yandex School of Data Analysis, Moscow, Russia)*

<sup>75</sup>*National Research Tomsk Polytechnic University, Tomsk, Russia  
[associated to Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia]*

<sup>76</sup>*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia—CSIC, Valencia, Spain  
(associated to ICCUB, Universitat de Barcelona, Barcelona, Spain)*

<sup>77</sup>*University of Michigan, Ann Arbor, Michigan, USA  
(associated to Syracuse University,  
Syracuse, New York, USA)*

<sup>78</sup>*Los Alamos National Laboratory (LANL), Los Alamos, New Mexico, USA  
(associated to Syracuse University, Syracuse, New York, USA)*

<sup>†</sup>Deceased.

<sup>a</sup>Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil.

<sup>b</sup>Laboratoire Leprince-Ringuet, Palaiseau, France.

<sup>c</sup>P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.

<sup>d</sup>Università di Bari, Bari, Italy.

<sup>e</sup>Università di Bologna, Bologna, Italy.

<sup>f</sup>Università di Cagliari, Cagliari, Italy.

<sup>g</sup>Università di Ferrara, Ferrara, Italy.

<sup>h</sup>Università di Genova, Genova, Italy.

<sup>i</sup>Università di Milano Bicocca, Milano, Italy.

<sup>j</sup>Università di Roma Tor Vergata, Roma, Italy.

<sup>k</sup>Università di Roma La Sapienza, Roma, Italy.

<sup>l</sup>AGH—University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

<sup>m</sup>LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

<sup>n</sup>Hanoi University of Science, Hanoi, Vietnam.

<sup>o</sup>Università di Padova, Padova, Italy.

<sup>p</sup>Università di Pisa, Pisa, Italy.

<sup>q</sup>Università degli Studi di Milano, Milano, Italy.

<sup>r</sup>Università di Urbino, Urbino, Italy.

<sup>s</sup>Università della Basilicata, Potenza, Italy.

<sup>t</sup>Scuola Normale Superiore, Pisa, Italy.

<sup>u</sup>Università di Modena e Reggio Emilia, Modena, Italy.

<sup>v</sup>H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom.

<sup>w</sup>MSU—Iligan Institute of Technology (MSU-IIT), Iligan, Philippines.

<sup>x</sup>Novosibirsk State University, Novosibirsk, Russia.

<sup>y</sup>Sezione INFN di Trieste, Trieste, Italy.

<sup>z</sup>School of Physics and Information Technology, Shaanxi Normal University (SNNU), Xi'an, China.

<sup>aa</sup>Physics and Micro Electronic College, Hunan University, Changsha City, China.

<sup>ab</sup>Lanzhou University, Lanzhou, China.