UNIVERSITY BIRMINGHAM University of Birmingham Research at Birmingham

Palaeobotanical experiences of plant diversity in deep time. 1:

Cleal, Christopher; Pardoe, Heather; Berry, Christopher; Cascales-Miñana, Borja; Davis, Basil; Diez, Jose; Martina, Filipova-Marinova; Hiesecke, Thomas; Hilton, Jason; Ivanov, Dimiter; Susanne, Leroy; MacElwain, Jennifer; Opluštil, Stanislav; Popa, Mihai; Seyfullah, Leyla; Stolle, Ellen; Thomas, Barry; Uhl, Dieter

DOI:

10.1016/j.palaeo.2021.110481

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

Citation for published version (Harvard):

Cleal, C, Pardoe, H, Berry, C, Cascales-Miñana, B, Davis, B, Diez, J, Martina, F-M, Hiesecke, T, Hilton, J, Ivanov, D, Susanne, L, MacElwain, J, Opluštil, S, Popa, M, Seyfullah, L, Stolle, E, Thomas, B & Uhl, D 2021, 'Palaeobotanical experiences of plant diversity in deep time. 1: How well can we identify past plant diversity in the fossil record?', *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 576, 110481. https://doi.org/10.1016/j.palaeo.2021.110481

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1	Palaeobotanical experiences of plant diversity in deep time. 1:
2	How well can we identify past plant diversity in the fossil record?
3	
4	Christopher J. Cleal ^{a§} , Heather S. Pardoe ^a , Christopher M. Berry ^b , Borja Cascales-Miñana ^c ,
5	Basil A.S. Davis ^d , Jose B. Diez ^e , Mariana V. Filipova-Marinova ^f , Thomas Giesecke ^g , Jason
6	Hilton ^h , Dimiter Ivanov ⁱ , Evelyn Kustatscher ^j , Suzanne A. G. Leroy ^k , Jennifer C. McElwain ^l ,
7	Stanislav Opluštil ^m , Mihai Emilian Popa ⁿ , Leyla J. Seyfullah ^o , Ellen Stolle ^p , Barry A.
8	Thomas ^q and Dieter Uhl ^r
9	^a Department of Natural Sciences, National Museum Wales, Cardiff CF10 3NP, UK,
10	chris.cleal@museumwales.ac.uk, heather.pardoe@museumwales.ac.uk
11	^b School of Earth and Ocean Sciences, Cardiff University, Cardiff CF10 3AT, UK,
12	berrycm@cardiff.ac.uk
13	^c CNRS, Univ. Lille, UMR 8198, Evo-Eco-Paleo, F-59000, Lille, France, borja.cascales-
14	minana@univ-lille.fr
15	^d Institute of Earth Surface Dynamics IDYST, University of Lausanne, Bâtiment Géopolis
16	CH-1015, Lausanne, Switzerland, basil.davis@unil.ch
17	^e Department of Marine Geosciences and Land Management, Universidade de Vigo, 36310
18	Vigo, Spain, jbdiez@uvigo.es
19	^f Varna Regional Museum of History, 41 Maria Louisa Boulevard, 9000 Varna, Bulgaria
20	^g Palaeoecology, Department of Physical Geography, University of Utrecht, P.O. Box 80115,
21	3508 TC, Utrecht, The Netherlands, t.giesecke@uu.nl
22	^h School of Geography, Earth and Environmental Sciences, University of Birmingham,
23	Birmingham, B15 2TT, UK, j.m.hilton@bham.ac.uk

- ⁱ Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Acad.
- 25 G. Bonchev Street, 23, 1113 Sofia, Bulgaria, dimiter@gbg.bg
- ^j Museum of Nature South Tyrol, Bindergasse/Via Bottai 1, Bozen / Bolzano, 39100, Italy,
- 27 evelyn.kustatscher@naturmuseum.it
- 28 ^k Aix Marseille Univ, CNRS, Minist Culture, LAMPEA, UMR 7269, 5 rue du Château de
- 29 l'Horloge, 13094, Aix-en-Provence, France, leroy@mmsh.univ-aix.fr; School of
- 30 Environmental Sciences, University of Liverpool, L69 3GP Liverpool, UK
- 31 ¹ School of Biology and Environmental Science, University College Dublin, Ireland,
- 32 jennifer.mcelwain@gmail.com
- ^m Institute of Geology and Palaeontology, Charles University, Albertov 6, 128 43 Prague 2,
- 34 Czech Republic, stanislav.oplustil@natur.cuni.cz
- ¹Southwest Petroleum University, School of Geosciences and Technology, 8, Xindu Ave.,
- 36 610500 Xindu, Chengdu, China; University of Bucharest, Faculty of Geology and
- 37 Geophysics, Department of Geology and Doctoral School of Geology, Laboratory of
- 38 Palaeontology, 1, N. Bălcescu Ave., 010041, Bucharest, Romania, mihai@mepopa.com
- ^o Department of Palaeontology, University of Vienna, Althanstraße 14, 1090 Vienna, Austria,
- 40 leyla.seyfullah@univie.ac.at
- 41 ^p *EP Research, 59320 Ennigerloh-Westkirchen, Germany, e.stolle.research@mail.de
- 42 ^q Institute of Biological, Environmental and Rural Sciences, University of Aberystwyth,
- 43 Aberystwyth SY23 1NL, UK, bat@aber.ac.uk
- ⁴⁴ ^r Senckenberg Forschungsinstitut und Naturmuseum, 60325 Frankfurt am Main, Germany,
- 45 dieter.uhl@senckenberg.de
- 46 [§] Corresponding author
- 47
- 48

49 Abstract

50 Palaeobotany and palynology are the main direct sources of evidence for studying 51 vegetation diversity dynamics through geological time. However, plant fossil diversity is 52 affected by various factors other than vegetation diversity, which need to be taken into 53 account in such studies. The use of fossil-taxa will potentially inflate perceived plant 54 diversities, requiring taxonomic lists to be normalised. Autochthonous floras provide the 55 most direct evidence of vegetation diversity but these are rare; most plant beds are 56 allochthonous with plant remains that have been subjected to varying levels of fragmentation, 57 transportation and time averaging. Local-scale vegetation diversity is especially difficult to 58 determine from the fossil record, even with rigorous sampling protocols and detailed 59 sedimentological analysis. Landscape-scale and regional-scale vegetation diversities are more 60 reliably determined but usually at the rank of family. Macrofossil and palynological data tend 61 to reveal evidence of different aspects of plant diversity, and the best results are obtained if 62 the two diversity signals are integrated. Despite the inherent difficulties, the plant fossil 63 record provides clear evidence of the dynamic history of vegetation through geological times, 64 including the effects of major processes such as climate changes and mass extinctions. 65

Keywords: Palaeobotany, Palynology, Biodiversity, Taxonomy, Taphonomy, Vegetation

68 1 Introduction

69 Vegetation has played a central role in the evolution of the Earth's biosphere, atmosphere 70 and landscape (Beerling 2007; Davies and Gibling 2010; Wellman 2010; Willis and 71 McElwain 2013); whilst it is possible to envisage a world having evolved with plants but no 72 animals, a world of animals without plants could not function. The raised public awareness of 73 the important ecosystem services provided by vegetation, including carbon capture to help 74 mitigate climate change, providing the foundations of all terrestrial trophic systems, and the 75 psychological benefits it brings to mankind, has resulted in a global research programme on 76 today's plant diversity and ecology (Antonelli et al. 2020). However, this only provides a 77 snapshot of a continuous ecological and evolutionary play that has taken place through some 78 500 million years of "Deep Time". To appreciate properly the significance of events such as the current biodiversity crisis ("the 6th mass extinction") and to anticipate potential outcomes, 79 80 it is vital that we understand this history of vegetation evolution.

Research into vegetation history started over two centuries ago (for summary see
Andrews 1980) but with the primary focus on plant phylogeny (Taylor et al. 2009; Cleal and
Thomas 2019). In recent years, interest in the study of plant fossil diversity has grown (as
summarised by Wing and DiMichele 1992; Willis and McElwain 2013) but investigating it
remains challenging (Wing and DiMichele 1995).

This is one of two papers arising from a workshop on past plant diversity entitled *Tracking changes in plant diversity over the last 400 million years*, which brought together specialists on diversity studies in fossil floras ranging in age from Devonian to Quaternary . The aim was to explore the different analytical methodologies and interpretative approaches used to investigate Phanerozoic plant diversity dynamics. The present contribution addresses what exactly we mean by biodiversity and to what extent can we extract biodiversity patterns from the plant fossil record. We will attempt to look at the relevant issues surrounding both plant macrofossils (i.e. fossils that can normally be seen with the naked eye, including
compressions / impressions / adpressions, casts / moulds and anatomically preserved fossils –
see Cleal and Thomas 2019) and microfossils (pollen, spores and phytoliths). The issues
surrounding sampling and analytical methods used will be discussed in our second paper (in
preparation).

98 2 What is biodiversity?

In biology, biodiversity is sometimes used to refer to functional diversity (the range of
traits in an assemblage) or phylogenetic diversity (the evolutionary breadth of an assemblage)
(Dornelas et al. 2012; Vellend et al. 2011, 2017). Palaeobotanists also sometimes investigate
trait diversity, such as the use of leaf physiognomy for estimating past climatic temperatures
(Wolfe 1993; Glasspool et al 2004). But diversity analyses of the plant fossil record tend to
be overwhelmingly of taxonomic diversity, and it is on this that we will focus here.

105 Taxonomic diversity in ecological studies consists of two factors: taxonomic richness 106 and taxonomic evenness (Tuomisto 2012). Taxonomic richness (the number of taxa present) 107 might be expected to be relatively easy to measure in both modern-day habitats and the fossil 108 record; Magurran (2004) has suggested that this alone can be a sensitive indicator of 109 ecological change. However, total richness can be difficult to determine if there are rare 110 species present, as these may be missed in surveys. A far more nuanced understanding of the 111 functioning of a flora will be obtained by determining its taxonomic evenness using 112 measurements such as Simpson's Index (e.g. Lande 1996; Veech et al. 2002) but this is only 113 really meaningful if it is reflecting the relative numbers of the organisms present. Variations 114 in the productivity of pollen, foliage and seeds between different plant parts (Fig. 1) mean that taxonomic evenness of fossil-taxa in a fossil flora will bear little or no relationship to the 115 116 taxonomic evenness of the original vegetation. The situation is particularly complex with

foliage, especially in pre-Cenozoic floras where the leaves are often compound structures that fragmented in different ways during abscission, transportation and preservation. Taxonomic evenness of a fossil flora may therefore be strongly influenced by taphonomy and how the plants fragmented post-mortem; although such data may provide some evidence as to relative biomass allocation within the vegetation (e.g. Baker and DiMichele 1997), its value for determining taxonomic evenness is limited.

123 Scale will clearly be critical in any diversity study, whether palaeontological or 124 biological (Bennington et al. 2009). R.H. Whittaker (1960) developed the most frequently 125 used concepts of taxonomic diversity for extant biotas, broadly recognised as α -diversity 126 (diversity in particular habitats) and β -diversity (diversity between habitats within a 127 landscape); these were then integrated to provide a γ -diversity (overall diversity within the 128 landscape). R.H. Whittaker (1977) later extended this scheme to include δ -diversity (diversity 129 between landscapes in a biogeographical province) and ε -diversity (overall diversity within 130 that province). However, R.H. Whittaker's (1960) terms were intentionally rather vague and 131 as a consequence have been used by different scientists in different ways (see Swingland 132 2001; Magurran 2004; Hamilton 2005 for reviews).

In an attempt to introduce taxonomic diversity concepts that more realistically reflect the
plant fossil record, Cleal et al. (2012) adopted a more flexible approach similar to those used
by R.J. Whittaker et al. (2001); see also Birks et al. (2016a,b) (Fig. 2):

136 (1) Local-scale diversity: the diversity of plant fossils observed in a single locality and 137 which probably reflects plant diversity within c. 1000 m^2 (c. 30 m x 30 m). In a palynological 138 context, it might more realistically refer to vegetation within up to 1 km². This will broadly 139 equate to the α -diversity of the parent vegetation. 140 (2) Landscape-scale diversity: the diversity of plant fossils observed within a typical 141 depositional basin and which probably reflects plant diversity within up to c. 10^5 km² (c. 300 142 km x 300 km) This will broadly equate to the γ -diversity of the parent vegetation.

143 (3) Regional-scale diversity: the diversity of plant fossils observed within a

144 palaeofloristic province and probably reflects plant diversity within more than 10⁵ km². This

145 will broadly equate to the ε -diversity of the parent vegetation.

146 It is important to remember that the diversities observed in the fossil record (both

147 macrofloral and palynological) represent the diversities of the fossils, and only partially

148 reflect the diversities of the parent vegetation (Gastaldo 1992; Birks et al. 2016). Some of the

149 resulting issues will be discussed later in this paper (Section 4).

150 **3** Taxonomic problems

151 3.1 Macrofossil taxonomy

152 The concept of biodiversity is inevitably tied to taxonomy (Khuroo et al. 2007). In 153 neobotany this is relatively straightforward as the taxonomy is based on whole-organism taxa 154 in which their lifecycles and development can be observed. There will always be 155 disagreements among botanists as to whether a particular genus of plants contains one or 156 more species, or a group of species belong to one or more genera, but at least botanists have 157 whole organisms against which to test their taxonomies.

158 With Cenozoic macrofloras (including Quaternary) it is often still possible to work with

159 whole-plant taxa (e.g. Huang et al. 2016) but the situation is more difficult with older floras

160 where palaeobotanists are dealing with extinct groups. Only rarely are completely

161 reconstructed organisms available to work with; even if a whole, articulated plant is

162 preserved (e.g. the early seed plant *Elkinsia* – Fig. 3) anatomical details are never completely

163 preserved. Palaeobotanists working on these stratigraphically older floras therefore use a

different taxonomic approach. Although this has changed in detail over the years (Cleal and
Thomas 2010), the underlying principle has in effect remained the same since the time of
Sternberg (1820) and Brongniart (1822): different parts of the plant are classified and named
separately as fossil-taxa (Turland et al. 2018, Art. 1.2). Mostly these are fossil-species and
fossil-genera, although in principle they can be of any rank (see Cleal and Shute 2012 for an
example of using fossil-families).

170 Exactly how a fossil-taxon is defined is a subjective matter and is not covered by the 171 regulations in the International Code of Nomenclature (Turland et al. 2018), but this is no 172 different from neobotany. Because of the constraints of the fossil record (e.g. the inability to 173 test hypotheses relating to reproductive isolation or molecular phylogenetics) fossil-taxa have 174 to be defined purely on morphological and/or anatomical criteria. Bateman and Hilton (2009, 175 p. 1256) recommended that each fossil-taxon "...must consistently possess at least one 176 morphological feature that it shares with no other [taxon]" (their autapospecies concept) and 177 should not take into account geographical or stratigraphical criteria. Whether this latter point 178 is always supportable is perhaps a moot point: for instance, is it helpful to place leaves of 179 Permian gigantopterids and of Cenozoic angiosperms in the same fossil-genus simply 180 because they have the same, diagnostic characters (Fig. 3)? However, since most diversity 181 studies at the rank of species or genus tend not to be making comparisons over such long 182 time-scales, this is probably not a significant problem here.

The problem with using fossil-taxa for diversity studies is that a simple summation of the names listed in published taxonomic lists will both significantly overestimate the number of biological species represented, and distort the relative representation of the different plant groups present (Cleal et al. 2012). For instance, in Carboniferous arborescent lycopsids, a single biological species may be represented by up to six separate compression fossil-species, whereas sphenopsids in the same flora may only have four fossil-species (Fig. 4; Table 1). An 189 added complication is that the fossil-taxa of the different plant parts are probably indicative 190 of different taxonomic ranks of the original organism: for instance, Stigmaria ficoides 191 (Sternberg) Brongniart is a fossil-species of phylogenetically conservative lycopsid rootstock 192 that effectively cannot be distinguished across many members of the order, whereas the stems 193 and cones have more sophisticated combinations of derived evolutionary characters and so 194 their fossil-species probably correlate better with the biological species of the organisms. 195 One solution would be only to study whole reconstructed plants (DiMichele and 196 Gastaldo 2008). This is feasible when dealing with higher-ranked taxa such as families (e.g. 197 Anderson et al. 2007) but at the present time there are too few reconstructions to provide 198 meaningful diversity data at the rank of species or genus. A solution is to normalise the 199 dataset by identifying, for each plant group, the plant part whose fossil-taxonomy is most 200 likely to reflect the original, whole-organism taxonomy (e.g. Hilton and Cleal 2007; Cleal et 201 al. 2012). For instance, the study of the late Carboniferous tropical swamps focussed mainly 202 on foliage taxa, except with the arborescent lycophytes for which the outer periderm layer 203 ("bark") of their trunks was used (Table 1; Cleal 2005, 2007, 2008a). Leaf morphotypes have 204 also been successfully used in this way with Cretaceous and Palaeogene angiosperms (e.g. 205 Johnson 2002). Although these vegetative fossil-taxa may not provide the best evidence of 206 phylogenetic relationships (reproductive structures would probably be better for this – e.g. 207 Meyen 1984), they are probably providing a robust reflection of the plant species diversity 208 (Cleal et al. 2012). This will inevitably be imperfect; for instance, cuticle studies of 209 Carboniferous Cordaites and Selaginella foliage have shown that diversities will be 210 significantly underestimated if the identifications are based purely on morphological data 211 (e.g. Thomas 2005; Šimůnek 2007). There is no easy solution to this issue and simply has to 212 be accepted in such diversity studies.

213 The situation is further complicated in that the same plant preserved in different ways 214 (e.g. petrifactions and compressions) will be recorded as different fossil-taxa (Galtier 1986; 215 Bateman and DiMichele 1992; Bateman et al. 1992; Bateman and Hilton 2009; Thomas and 216 Cleal 2020). It is critical, therefore, to ensure that assessments of diversity do not duplicate 217 fossil-taxa in the same assemblage or locality that are preserved in different ways; for 218 instance, if an assemblage should include lycopsid cones as both compressions and 219 permineralisations, the taxonomic list should be normalized so that diversity is not artificially 220 inflated by "double counting".

221 3.2 Palynotaxa

222 Palynological studies on Quaternary floras tend to use whole-plant taxa, based on 223 morphological comparisons with pollen that have been extracted from living plants. It is 224 sometimes possible to distinguish pollen from closely related plant species based purely on 225 morphology but often palynological studies tend to focus mainly on differentiating plants at 226 the generic rank. Attempts have been made to use DNA barcoding to improve the taxonomic 227 resolution in Quaternary studies (e.g. Seppä and Bennett 2003); for instance, Petit et al. 228 (2002) demonstrated that the modern genetic diversity of oak is consistent with the pollen 229 evidence in a study of post-glacial oak migration. However, most Quaternary palynological 230 studies remain essentially morphology-based.

With older floras, the known relationship between the pollen / spores and their parent plants is less certain and so palynologists have developed separate taxonomic schemes (Chaloner 1999). Some proposed taxonomies are completely artificial with the taxa defined purely on morphological criteria with a non-Linnaean nomenclature, such as used in many oil-company palynological databases and in the Biorecords methodology of Hughes (1963) (see Traverse, 2007 for a review). Other taxonomies use a Linnaean-style nomenclature but with taxa that were still essentially morphological (e.g. Potonié 1956, 1958, 1960) and it is this approach which is most usually widely used today in pre-Neogene studies (e.g. Jasper etal. 2010; Stolle 2007, 2012, 2016; Hochuli et al. 2016).

240 Because the botanical affinities of many pre-Neogene palynotaxa are uncertain, it can be difficult to translate observed palynodiversity trends into floristic trends. Thomas (1987) and 241 Mander and Punyasena (2014) suggested that the situation could be improved by revising the 242 243 diagnoses of palynotaxa based on evidence from in situ palynomorphs in fructifications, data which is now being increasingly collated (e.g. Balme 1995; Bek 2017). Experience with 244 245 Palaeogene and Neogene pollen and spores has also shown that a combination of light 246 microscopy and scanning electron microscopy based on individual grains (Ferguson et al. 2007), although very time consuming, can help to improve their assignment to a particular 247 248 plant genus or family, or perhaps even to map it into an established framework represented by 249 one or more phylogenetic trees. (e.g. Grímsson et al. 2011a,b, 2015a,b). Chemical analysis 250 such as using FTIR (Fourier Transform Intra-Red) and fluorescence spectroscopy can also be 251 helpful in determining affinities of particular palynormophs (e.g. Mitsumoto et al. 2009; 252 Steemans et al. 2010; Urban et al. 2010). This approach has shown that more traditional 253 approaches utilizing only light microscopy tend to underestimate the number of taxa present 254 in a palynoflora (Hofmann and Gregor 2018).

An added complication is the variation in morphology of pollen and spores during maturation, as shown for instance in the fern *Weichselia reticulata* (Stokes and Webb) Fontaine (Fig. 5). This is not an issue in most diversity studies on dispersed palynofloras, as plants do not normally release their pollen or spores before they are fully mature. However, if a plant has been subjected to trauma such as a storm, immature pollen and spores may be prematurely released and preserved, and this could inflate the diversity of a palynofloras. Although labour intensive, it is possible to determine whether different morphologies represent different states of maturity or just variability of miospore forms within a species
using sporoderm ultrastructure analysis (e.g. Zavialova et al. 2010).

Because of the problems of classifying stratigraphically older palynotaxa a number of purely morphological suprageneric classifications have been developed (see Traverse 2007 for a review). Especially in Palaeozoic palynofloras, a nested hierarchy of morphological groups (anteturma, turma, subturma, etc.) developed by Potonié (1934) is still widely used, and provides a useful framework for descriptive studies. However, as these groups are strictly morphological, they rarely relate to botanical suprageneric groups and so are of limited use in diversity studies.

271 3.3 Taxonomic rank

Because of the problem of relating pollen and spores to particular plant species, using palynology for species diversity studies can be difficult (Mander and Punyasena 2014); even in the Quaternary where the relationship between pollen and parent plants is better-known, most palynological diversity studies tend to be at the rank of genus or even family (Giesecke et al. 2014). Such studies have nevertheless provided valuable evidence of vegetation dynamics especially at the landscape-scale (Section 5.2).

278 Local- and landscape-scale plant macrofossil diversity studies tend to be based on normalised inventories of fossil-species or possibly fossil-genera (Section 3.1). However, 279 280 species are currently impractical when dealing with diversity changes at regional- or global-281 scales, and over longer time-scales, as the datasets become too large to collate and check 282 objectively by any individual scientist or team. Even where a large amount of species data 283 has been historically accumulated, such as for the Pennsylvanian Subsystem (see comments 284 by Pfefferkorn et al. 2017), there have been few attempts to collate them coherently and 285 critically. Where such collations have been attempted (e.g. Niklas et al. 1980; Lidgard and 286 Crane 1990), methodological and sampling issues occurred (see comments by Niklas and

Tiffney 2010; Cascales-Miñana et al. 2013). Moreover, these early collations were not
published and so cannot be subjected to subsequent critical taxonomic re-assessment, making
the robustness of the resulting analyses difficult to judge.

290 The situation may potentially improve with the development of large-scale computer 291 databases of fossil occurrences, such as those for Cenozoic angiosperms (Xing et al. 2016; 292 Williams et al. 2018). Palaeobotanical data have also been included in the Paleobiology 293 Database (Alroy 2003) although its coverage for plant fossils remains uneven, and is far 294 below that in other groups such as fossil vertebrates. Various numerical approaches have 295 been investigated that aim to overcome the issues of incomplete sampling of such databases 296 (e.g. Silvestro et al. 2015; Beri et al. 2020) but the intractable problem remains of verifying 297 the taxonomic robustness of the data; if the data cannot be trusted, how can the results of any 298 analysis? This is an area where palaeobotany needs to improve in order to catch up with other 299 fossil groups and make sustained impact in analytical methodologies.

300 In the absence of usable databases, the solution adopted in many regional- and global-301 scale macrofloral diversity studies is to analyse changes at the rank of family. Family is the 302 lowest rank of fossil-taxa based almost exclusively on whole organisms and so potentially the 303 dynamics of the fossil-families should be comparable with those of the original parent 304 families. A number of global collations of plant fossil-family distributions through geological 305 time are available (e.g. Harland 1967; Benton 1993; Collinson 1996; Anderson et al. 2007) 306 and they include the evidence on which the records were based and so can be subjected to 307 later critical assessment and potential revision (Cascales-Miñana and Cleal 2014).

308 But how closely do family dynamics mirror diversity dynamics at lower taxonomic 309 ranks? Analyses on modern-day tropical forests suggest that family and species diversity 310 patterns are broadly similar (e.g. Enquist et al. 2002; Jantz et al. 2014) especially if the data 311 are log transformed to reduce the effect of dominant families (La Torre et al. 2007); see also 312 comments by Giesecke et al. (2019) and Reitalu et al. (2019) on Holocene data from Europe. 313 However, this does not take into account the taphonomic filter that fossil floras have been 314 subjected to; many Palaeozoic and Mesozoic plant fossils cannot be assigned to families due 315 to missing, or difficult to deduce, features of reproductive organs or cauline anatomy, 316 suggesting the fossil record of families is incomplete. On the other hand, regional- and 317 global-scale vegetation analyses (e.g. Cascales-Minana et al. 2013) may benefit from using 318 family data because they may help smooth out some of the sampling problems encountered in 319 such large-scale analyses. This is clearly a subject that needs further investigation.

320

4 Representativeness of data

321 There have been many studies looking at the effects of representativeness on diversity 322 studies in the macrofossil record, such as the effects of sampling and taphonomy, but mainly 323 dealing with faunas, notably marine invertebrates (e.g. Kowalewski et al. 2006). However, 324 the issues surrounding such faunal studies are fundamentally different from those facing 325 palaeobotanists and palynologists, as most palaeozoologists have the luxury of dealing with the remains of whole organisms (or at least their hard-parts, such as shells or exoskeletons); 326 327 even vertebrate palaeozoologists tend to deal with whole-organism taxa. Palaeobotanists and 328 palynologists, in contrast, deal almost exclusively with allochthonous and fragmentary 329 remains; there are exceptions, as we will discuss, but these tend to be rare and scattered, and 330 difficult to use in diversity studies. This means that diversity studies on the plant fossil record are addressing quite different questions to those being usually asked by palaeozoologists: 331 332 palaeobotanists and palynologists tend to be looking at broad composition of vegetation 333 either in terms of taxa or biomass rather than looking at changing community structure in terms of individual organisms (e.g. Bambach 1977). 334

335 4.1 Macrofloral data

336 Autochthonous floras (sometimes misleadingly referred to as "Lagerstätten") provide the 337 most reliable data on original plant diversity, especially at a local-scale, but these are rare. 338 One of the best documented is the Devonian Rhynie Chert (e.g. Edwards et al. 2018; 339 Garwood et al. 2019; Strullu-Derrien et al. 2019) where an in situ and almost complete 340 terrestrial biota is preserved including relatively small, herbaceous plants. Autochthonous 341 fossil floras with larger, woody plants are much rarer. There are exceptions, such as the 342 Palaeozoic swamp forests that were rapidly covered by volcanic ash (Sections 5.1, 5.2); but more usually, the so-called T⁰ fossil or submerged forests (DiMichele and Falcon-Lang 2011) 343 344 are only partly autochthonous. They form where an area of forests has been engulfed by a 345 flood of sediment and casts of the stumps have been preserved in situ (e.g. Fig. 6; for other 346 examples see Heyworth and Kidson 1982; Francis 1983; Gastaldo 1985; Pole 2001; Calder et 347 al. 2006; Wagner and Diez 2007; Moir et al. 2010; Stein et al. 2012; Berry and Marshall 348 2015; Thomas and Seyfullah 2015; Falcon-Lang et al. 2016), but most of the herbaceous 349 ground-cover and liana species have been winnowed-out (Thomas 2014). Other types of 350 "fossil forests" consist of petrified logs preserved as log-jams that have been subject to 351 varying degrees of transportation (e.g. Falcon-Lang and Bashforth 2005) and thus also difficult to use for diversity studies. 352

More typically, plant beds occur in fluvio-lacustrine deposits, where disarticulated plant remains have accumulated after varying degrees of transportation either by wind or water (Burnham 1989; Gastaldo et al. 1995, Kędzior and Popa 2013, 2018; Thomas and Cleal 2015). This is in marked contrast to many fossil faunal communities, which tend to be much less prone to transportation (Kidwell and Holland 2002). Many attempts at palaeoecological studies on such plant beds have documented in great detail the sedimentological context where the fossils occur (e.g. Scott 1978, 1989; Gomez et al. 2012; Kędzior and Popa 2013, 360 2018). Detailed, three-dimensional sampling such as in underground coal mines along 361 directional and transversal galleries and in coal extraction chambers in particular can provide 362 valuable data (e.g. Gastaldo 1985; DiMichele and Nelson 1989; Popa 1998, 2011, 2014; 363 DiMichele et al. 2007, 2017; Barbacka et al. 2016). 364 However, the plant remains will have been transported over varying distances, making it difficult to translate the observed distribution of the fossils into original plant diversity 365 366 (Gastaldo 1992). In a few cases, the fossils in such plant beds seem to have been subjected to 367 only limited transport, such as where a river-bank bank has collapsed and the plant remains 368 have become entombed in a crevasse-splay (e.g. Cleal and Thomas 1988; Laveine and Belhis 369 2007). More usually, however, the plant remains are at least partly allochthonous. Actualistic 370 studies suggest they will include only a variable representation of the immediately local 371 vegetation (e.g. Burnham 1989, 1994) mixed with remains derived from riparian vegetation 372 growing some distance upstream (Spicer 1980, 1981; Scheihing and Pfefferkorn 1984; 373 Ferguson 1985; Gastaldo et al. 1987; Gastaldo and Huc 1992). 374 By carefully documenting the co-occurrence of species within individual beds, the 375 composition of individual plant communities can be estimated (e.g. Procter 1994; Bashforth 376 et al. 2010, 2011; Barbacka 2012; Barbacka et al. 2016; Thomas et al. 2020). However, locating those communities in the original vegetation / habitat matrix requires a detailed 377 378 understanding of the sedimentology of the deposits (DiMichele and Gastaldo 2008; Reitalu et 379 al. 2014) and is at best difficult. Moreover, taphonomic factors may distort the observed 380 diversities. Variations in edaphic conditions can cause variable post-mortem decay of the 381 plant tissue (Gastaldo 1992; Gastaldo and Demko 2011). It has been suggested that 382 differential decay of plant groups may distort the species composition (Scott 1979; Wing and 383 DiMichele 1995) although the effect of this may have been exaggerated (Locatelli et al. 2016; Tomescu et al. 2018). More significant may be differential sorting of the plant remains during 384

385 transportation: smaller plant fragments will tend to travel further and softer, heavier plant 386 fragments sink more quickly (e.g. Steart et al. 2002). Some element of time-averaging may 387 even occur within a single plant bed depending on depositional rates of the sediment. 388 Plant remains preserved in shallow marine deposits are usually fragmentary and not 389 concentrated into distinct plant beds, although there can be exceptions caused by storm surges 390 (e.g. Kustatscher et al. 2010). Fossil floras preserved in marine deposits can include the 391 remains of coastal vegetation such as mangroves (e.g. Collinson 1993). During late Permian 392 times, climatic conditions were unfavourable for plant growth in continental Europe and so 393 vegetation tended to be concentrated in coastal areas; remains of this vegetation has been 394 found in shallow marine deposits, preserved particularly during transgression phases 395 (Kustatscher et al. 2017).

396 Some plant macrofossils preserved in marine strata have been interpreted as plant 397 remains washed down from hinterland vegetation (e.g. Rothwell et al. 1996; Rice et al. 1996; 398 Cleal and Rees 2003) and are notably different from what is seen in fluvio-lacustrine plant 399 beds. "Exotic", extra-basinal plant remains have also sometimes been reported from fluvio-400 lacustrine plant beds (e.g. Cleal and Thomas 2004; Uhl 2006; Opluštil et al. 2007). Generally, 401 however, plants growing in places away from rivers or lakes are poorly represented as 402 macrofossils. For instance, grasses, which are obviously major components of terrestrial 403 vegetation today, have a very poor macrofossil record and much of what we know of their 404 evolution is based on palynology (Section 4.2) or dispersed phytoliths derived from their 405 leaves (e.g. Piperno and Pearsall 1998; Strömberg 2004, 2011). 406 Fossil floras can also occur in maar lake deposits (e.g. the Messel World Heritage Site –

407 Collinson et al. 2012). Such lakes are caused by phreatomagmatic explosions resulting from408 the interaction of erupting magma and water, and can occur almost anywhere within a

409 landscape and thus may be surrounded by a different type of vegetation to that growing

adjacent to lakes formed in fluvio-lacustrine settings. For instance, the late Oligocene Norken
fluvio-lacustrine deposits contain predominantly remains of riparian and swamp vegetation
(Uhl et al. 2018) but these are almost totally absent from the nearby, almost contemporary
Enspel maar lake deposits (Köhler and Uhl 2014). Plant remains in such lakes also
experience less hydro-mechanical stress due to water transport and so can preserve delicate
plant structure such as flowers (e.g. Uhl 2015).

416 4.2 Palynological data

417 Palynology has been widely used for Quaternary landscape-scale diversity studies 418 (Giesecke et al. 2014); but translating the data into vegetation patterns can be problematic 419 because of significant variation in pollen productivity from year to year (Andersen 1970; 420 Sugita 1993; Hicks 1985; Barnekow et al. 2007; Pidek et al. 2010; Giesecke et al. 2014). 421 However, this is partly mitigated by most sediment samples representing several years; for 422 example, in the Lake Sapanca sequence (N-W Turkey) sub-annual samples taken at a 5 mm 423 resolution revealed no seasonality in the palynology signal, probably due to bioturbation of 424 the lake sediment (Leroy et al. 2009). On the other hand, in the alternating black and white 425 sediment layers of the Dead Sea (López-Merino et al. 2016), the seasonality of the pollen 426 production was used to determine if the lamina couplets were varves or a laminated sediment. 427 Another problem is the great variation in pollen productivity between different plant 428 species. Current evidence for northern and temperate latitudes suggests that Quaternary 429 palynological data are particularly robust for most trees (with a few notable exceptions such 430 as Larix) and wind-pollinated taxa, and provide a good measure of broad-scale plant richness 431 over several thousands of kilometres (Reitalu et al. 2019); this is less so for tropical 432 environments due to the higher number of insect-pollinated plants. Among herbs, the Poaceae 433 are the most abundant wind-pollinated plants, and their pollen can be widely dispersed. 434 However, the source of Poaceae and Cyperaceae pollen can be difficult to elucidate because

435 these species occur in a wide range of plant communities. It can also be impossible to identify 436 their pollen to species level other than in cultivated cereals (Pardoe 2001; Sjögren et al. 437 2015), although phytoliths can be of help here (see below). Most other herbs tend to be under 438 represented in pollen spectra (Leroy and Roiron 1996) as the pollen are dispersed by other 439 vectors and so are not so abundantly produced; they also often have a lower preservation 440 potential. There have been relatively few studies of the representation of herbs in pollen assemblages (Pardoe 2001; Bunting et al. 2016) but the presence of so called "indicator taxa" 441 442 in pollen samples can give strong evidence that such plants were growing locally (Pardoe 443 1996, 2001, 2006). Data can also be supplemented by evidence from in situ pollen from 444 flowers (e.g. Herendeen et al. 1994) and in exceptional cases from fossils of pollinating 445 animals (e.g. Grímsson et al. 2017).

446 Recent initiatives such as the Pollen Monitoring Programme (PMP) are now helping us gain a greater understanding of the relationship between pollen, vegetation and environmental 447 448 variables. The PMP has been instrumental in the publication of several decades-long records 449 from across Europe (Hicks et al. 1996; Giesecke et al. 2010). The PMP has addressed a 450 variety of problems including the representation of individual taxa (Hicks et al. 1994; Hicks, 451 2001, Ertl et al 2012; Pidek et al. 2010), the representation of plant communities (van der Knaap et al. 2001; Pidek 2004; Gerasimidis et al. 2006), and the influence of sampling 452 453 medium on palynological diversity (Pardoe et al. 2010; Litsitsyna et al. 2012). 454 Although not strictly palynological, phytoliths are another type of plant microfossils that 455 provide valuable evidence of terrestrial vegetation (Strömberg et al. 2018). There can be taphonomic issues due to silica dissolution (Cabanes and Shahack-Gross 2015) but they are 456 457 nevertheless essential indicators of grass diversity in Cenozoic vegetation, which are usually 458 poorly represented as pollen and macrofossils (Piperno and Pearsall 1998; Piperno 2002;

459 Strömberg 2004, 2011; Rashid et al. 2019). There are also records of pre-Cenozoic phytoliths
460 (e.g. Carter 1999) but their affinities remain uncertain.

461 In principle, palynodata can be corrected using R-coefficients (sensu Davis 1963) 462 representing the ratio between the observed pollen abundances and the abundance of plants in 463 the parent vegetation. R-coefficients can be estimated for Quaternary data based on 464 actualistic comparisons between surface pollen data and field vegetation surveys (although 465 even here problems may occur because of some of the mathematical assumption involved -466 Parsons and Prentice 1981). Such an approach is more difficult with tests on the robustness of 467 pre-Quaternary palynodata as often no independent measure of vegetation composition can 468 be used to calculate the R-coefficients. Nevertheless, it has been attempted with the late 469 Carboniferous swamps where available autochthonous macrofloras allow the coefficients to 470 be estimated (Willard 1993; Opluštil et al. 2009). Palynofacies signals can also help in 471 determining the robustness of palynological data by indicating the depositional and 472 palaeoenvironmental situation of the studied strata (e.g. Stolle et al. 2012, pl. 1, fig. 2).

473 *4.3 Pollen and macrofossil data compared*

474 When diversity data from the macrofloral and palynological records are compared (e.g. 475 Leroy and Roiron 1996; Dimitrova et al. 2005; Birks and Bjune 2010; Xiong et al. 2013; Bjune 2014; Looy et al. 2014; DiMichele et al. 2018) rather different signals are often 476 477 revealed both in the plant groups represented and the relative proportions of those plant 478 groups (Fig. 7). The macrofloral record is regarded as giving a more detailed picture of plant 479 species richness, especially at a local-scale (Birks and Birks 1980). However, this tends to 480 represent only a fairly narrow band of habitats, and the much smaller sample sizes usually 481 available compared with palynology will often be insufficient to capture diversity patterns. 482 Palynology, in contrast, will give a better understanding of vegetation patterns across a wider 483 range of habitats and at a landscape-scale (Dimitrova et al. 2010; Costamagna et al. 2018).

484 Because palynological samples may contain palynomorphs from different habitats, it can be 485 difficult to determine the local-scale vegetation patterns within individual habitats; it may 486 also help explain why palynospectra tend to be more diverse than the macrofloras found in 487 the same bed (e.g. Dimitrova et al. 2005). However, as our understanding of the natural 488 affinities of many palynotaxa is improving, palynology is proving increasingly refined 489 evidence of landscape-scale vegetation diversities (Section 5.2). It is evident that one data 490 source is not better than the other for diversity studies: rather, that palaeobotany and 491 palynology are complementary, and are best investigated in tandem (Birks 2000; Kustatscher 492 et al. 2010; Reitalu et al. 2014; Costamagna et al. 2018).

493 **5 Diversity studies**

494 It is beyond the scope of this paper to review all examples of palaeobotanical and 495 palynological diversity studies; the following discussion aims merely to illustrate some of the 496 types of analyses that have been attempted.

497 5.1 Local-scale diversity

498 Most allochthonous fossil macrofloras tend to reflect local-scale plant diversity (Cleal et 499 al. 2012). However, the complexity of the sedimentary systems in which they usually occur 500 (Section 3.1) means that the diversity of each individual bed needs to be analysed separately 501 as each flush of sediment is likely to have remains from a different set of plant communities. 502 Even if the plant beds are rigorously sampled (e.g. Scott 1978, 1979), a detailed 503 understanding of the sedimentology is required before it is possible to unscramble the local-504 scale plant diversity patterns from the mosaic of habitats represented in most allochthonous 505 plant bed (Kedzior and Popa 2013, 2018). 506 Palynological data tend to be even more problematic for local-scale diversity studies due

507 to variations in how far the palynomorphs have been transported. For instance, pollen of

508 modern-day Picea has been found in the Canadian Arctic, 3000 km from its source 509 (Campbell et al. 1999); in the Palaeozoic, conifer pollen appears to have been blown in from 510 a considerable distance (e.g. Bless et al. 1977); some pollen in Carboniferous tropical 511 palynofloras even appear to have originated from high-latitude, Gondwana vegetation 512 (Dimitrova et al. 2011). Even long-distance water transportation of pollen has been reported; 513 for example, Holocene *Podocarpus* pollen that have been found in Nile delta deposits may 514 have been transported > 2,000 km along the river from their source in the Ethiopian 515 Highlands (Leroy 1992). Although such exotics will normally be rare in palynofloras, they 516 represent the end-members of a gradational spectrum of palynomorph abundances reflecting 517 differences in transportation distances, making it difficult to extract local-scale diversity 518 patterns, especially in fluvial and delta settings (Weng et al. 2006).

519 Local-scale past plant diversity is best determined in the rare autochthonous fossil floras 520 although even here the data are often incomplete (Section 4.1). Some of the best examples of 521 autochthonous floras preserving forest vegetation including both the trees and herbaceous 522 plants are in Palaeozoic volcanic ash-fall deposits (e.g. Wagner 1989; Rössler and Barthel 523 1998; Wang et al. 2012; Luthardt et al. 2016). Examples studied in great detail are in an ash 524 band in the early Moscovian Radnice Coal in the Czech Republic, where about 0.5 m of volcanic ash engulfed an area of swamp vegetation. The lower part of the deposit contains in 525 526 situ stumps and the groundcover vegetation, which, because the ash fell almost vertically, 527 was mostly not winnowed out. The upper parts of the ash band, in contrast, includes remains 528 of the upper parts of the trees, together with epiphytes and lianas, brought down by the weight of the ash sometime after the deposit had fallen (Pšenička and Opluštil 2013). A 529 530 careful survey of the plant remains in different levels of the ash deposit (Fig. 8) not only 531 allowed the reconstruction of the taxonomic composition, spatial distribution and density of

532 vegetation cover, but also revealed evidence of plant to plant interactions and living strategies 533 in extraordinary detail (Opluštil et al. 2007, 2009a,b, 2014; Libertín et al. 2009a). 534 Many coals (but not all – Glasspool 2003) are the remains of parautochthonous peat and 535 so, as with modern-day peat deposits (e.g. Mauquoy et al. 2010), have the potential to reveal 536 local-scale plant diversity. When the peat has changed into coal through compaction and 537 diagenesis, however, the plant remains become homogenised and so difficult to identify. 538 Notable exceptions are when the peat has been subjected to early mineralisation that 539 preserves the anatomy of the plant remains in often exquisite detail. Sometimes most or all 540 the peat deposit has been mineralised (e.g. Galtier 2008; Slater et al. 2015) but more 541 commonly the mineralisation is localised, such as in the coal balls (mainly calcitic nodules) 542 found in some Palaeozoic coal seams. There have been several local-scale diversity studies 543 on coal balls (e.g. DiMichele and Phillips 1988; DiMichele et al. 1991; Willard 1993; Baker 544 and DiMichele 1997; DiMichele et al. 2002; Willard et al. 2007), which produced evidence 545 of biomass allocation within the peat, which in turn gave some localised evidence of species 546 diversity.

547 Coal deposits often yield well-preserved palynomorphs, which have been extensively 548 used for biostratigraphical studies revealing evidence of the temporal changes in vegetation 549 (e.g. Smith and Butterworth 1967). If intercalated fine-siliciclastic (shaley) coal-bearing 550 samples are also included, palynological assemblages can be particularly species rich. As 551 with the macrofloras, regional to exotic extra-basinal palynomorphs may also be present (Fig. 552 9), which can be ideal for palynological correlation purposes (e.g. Stolle 2007, 2010), but can confuse local-scale species richness analyses. Palynology has also been used to investigate 553 554 the ecological development of the swamps (e.g. Smith 1962, 1968; Habib and Groth 1967; 555 Jasper et al. 2010; Johnston et al. 2017; Eble et al. 2019) and to look at plant diversity at the 556 rank of genus and higher (e.g. Dimitrova and Cleal 2007; Libertín et al. 2009b; Thomas and

557 Dimitrova 2017), but direct translation of the resulting palynological spectra into plant
558 species diversity is difficult.

559 Another distinctive parautochthonous source of plant remains is amber, mainly of 560 Cretaceous to Neogene age. Amber can be produced by both conifer (Sadowski et al. 2017) 561 and angiosperm trees (Rust et al. 2010), and can result in exquisite preservation, especially of 562 delicate structures such as flowers (Poinar 2002; Gandolfo et al. 2018), fern sori (Sadowski et 563 al. 2019) and even microscopic algae (Schmidt et al. 2006). Some of these deposits have been studied since the middle 19th century, but amber can be a very selective fossil trap (e.g. 564 565 Solórzano Kraemer et al. 2018) and so our understanding of the plant diversity of these 566 forests is still incomplete.

567 5.2 Landscape-scale diversity

568 Studies on compression fossil diversities across depositional basins (e.g. Cleal 2005, 569 2007, 2008a; Goswami and Singh 2013; Huang et al. 2016; Opluštil et al. 2017; Goswami et 570 al. 2018; Roopnarine et al. 2018; Saxena et al. 2020) tend to be based on plant remains from a 571 narrow band of habitats. For instance, compressions from the Pennsylvanian swamps of 572 Euramerica appear to have been dominated by remains of the vegetation growing on clastic 573 substrates such as flood-plains, levees and sand banks, whereas the peat-substrate vegetation, 574 which in fact dominated these swamps, is often poorly represented (Cleal et al. 2012); the 575 peat-substrate vegetation is, in contrast, better represented in the coal ball floras and 576 palynospectra. This is not a problem if the main aim is to document extrinsic effects such as 577 climate or landscape changes, particularly if the sampled habitats are tightly constrained 578 ecologically, but care must be taken not to over-generalise the results in terms of overall 579 vegetation patterns.

580 One of the best sources of detailed data on Palaeozoic landscape-scale diversity are the 581 ash deposits in the Czech Radnice Coal (mentioned in Section 5.1), which have been 582 recorded from numerous localities in both historical collections and several recent 583 excavations. These have allowed lateral variation in the swamp vegetation at one 584 stratigraphical level to be investigated; for instance, studies at the Štilec and Ovčín localities, 585 about 20 km apart, yielded two contrasting assemblages, representing different stages of 586 vegetation succession (Opluštil et al. 2007, 2009a,b, 2014; Libertín et al. 2009a). A similar 587 situation is present in the earliest Permian Wuda ash bed that occurs over an area of more than 60 km², enabling distinct assemblages to be recognised both vertically and laterally 588 589 (Wang et al. 2012).

590 Floras with anatomically preserved petrifactions and permineralisations are more 591 difficult to use for landscape-scale diversity studies. Most such floras tend to be isolated 592 localities reflecting the exceptional conditions that caused the preservation, and so usually 593 only reflect local-scale diversity. The most notable exceptions are the Pennsylvanian-age coal 594 balls floras that occur extensively across the Late Palaeozoic tropical belt but, although they 595 have been the subject of a number of taxonomic collations (e.g. Phillips 1980; Galtier 1997), 596 no detailed landscape-scale diversity studies have been attempted. In palaeozoological 597 studies, such preservational "hot-spots" have proved a problem by suggesting abnormally 598 high diversities at particular stratigraphical levels, often referred to as the "Lagerstätte effect" 599 (e.g. Benton 1995; Butler et al. 2013), but evidence of this distorting effect on plant fossil 600 diversities is less clear (see comments by Cascales-Miñana and Gerrienne 2017). 601 Palynology can provide a more representative picture of landscape-scale diversity as the 602 sediment will contain the pollen from plants growing across the area (Weng et al. 2006). This 603 has proved particularly useful in Quaternary studies where the botanical affinities of the 604 various pollen types are well known (Section 4.2). For instance, palynology has been used to

605 map distribution changes across Europe during the Holocene by Huntley and Birks (1983),

and there have been numerous species-specific studies (Hicks 2001; Brewer et al. 2002;

25.

Giesecke and Bennett 2004; van der Knaap 2004; Latałowa and van der Knaap 2006; Tinner
and Lotter 2006; Giesecke et al. 2007; Pidek et al. 2010; Poska and Pidek 2010). Reitalu et
al. (2019) have demonstrated in their study of modern pollen and plant richness across
northern Europe that the highest correlations were for trees and shrubs and of wind-pollinated
taxa, suggesting that these are the best measures of broad-scale plant richness over several
thousands of kilometres.

613 Improvements in our knowledge of the general affinities of many pre-Neogene

614 palynotaxa (Section 3.2) now allow palynology to identify broad patterns of landscape-scale

615 plant diversity (Abbink et al. 2004; Dimitrova et al. 2005, 2010; Dimitrova and Cleal 2007;

616 Kustatscher et al. 2010; Beri et al. 2018; Franz et al. 2019). However, remaining uncertainties

617 about variations in palynomorph productivity and dispersal between species, and the

618 morphological variation of palynomorphs within plant species, make it difficult to use some

619 taxa for detailed landscape-scale plant diversity studies (Section 4.2).

620 5.3 Regional-scale and global-scale diversities (Evolutionary floras)

621 Studies on global-scale faunal diversity (e.g. Sepkoski 1978, 1979, 1984, 1988; Bambach 622 1977; Powell and Kowalewski 2002) have shown a progressive increase in species doveraity 623 through the Phanerozoic due to an increase in the spatial density of organisms, especially in 624 shallow marine environments (Holland and Sclafani 2015). Similar global and regional 625 studies at the species rank have been attempted in palaeobotany (e.g. Knoll et al. 1979; Niklas 626 et al. 1980) but were hindered by the lack of suitable, taxonomically robust data sets (Section 627 3.3); also by the failure to take into account geographical (especially latitudinal) variations in 628 taxonomic diversity, as has been shown to be an issue with marine invertebrate diversity 629 dynamics (Close et al. 2020). Analogous palaeobotanical studies would, moreover, be unlikely to answer the same sorts of questions of changes in community structure that were 630 631 being investigated in the faunal record (Section 4).

632 Analyses within narrow taxonomic (e.g. Cleal 2008b,c) or stratigraphical (e.g. Cleal et al. 633 2010; Barbacka et al. 2014) limits have been attempted at the regional-scale, which make it 634 practical for the taxonomic robustness of the data to be critically assessed. However, most 635 larger-scale studies have tended to be based at supra-generic ranks, usually family. For instance, global Phanerozoic plant diversity dynamics were interpreted using Evolutionary 636 637 Floras (Fig. 10), identified from a factor analysis of a plant family dataset (Cleal and 638 Cascales-Miñana 2014), and these have been used to describe the broad trajectory of 639 vegetation history (Cleal and Thomas 2019; Cleal 2019). More recently, a similar study on 640 pre-Carboniferous floras at the rank of genus is revealing further details of the early phases of 641 plant terrestrialisation (Capel et al., this volume).

642 There are a number of problems with such large-scale plant diversity studies. The 643 taxonomic robustness of the data used is often uncertain, although for plants this is partly 644 avoided by using family-rank data sets (Section 3.3). More difficult is the robustness of the 645 stratigraphical correlations between widely separated floras. Most fossil floras occur in 646 predominantly terrestrial sequences that lack absolute dating or independent 647 biostratigraphical control (e.g. by marine faunas). In local-scale and landscape-scale studies 648 this is less of a problem as lithostratigraphical correlations are often sufficient to provide a 649 temporal framework for comparisons, but these are inadequate for regional-scale and global-650 scale studies. The palaeobotanical and palynological evidence is itself sometimes used to 651 provide the correlations, but when this is used as the temporal context for the vegetation 652 changes, the arguments become circular. A classic example is the Panchet Formation in India, 653 which is often quoted as justifying the persistence of glossopterids into the Triassic Period, 654 but for which there is in fact no evidence that it is Triassic other than some debatable facies 655 changes and the floras themselves (Saxena et al. 2018).

656 6 Why study plant diversity in deep time?

657 Studies of past plant diversity dynamics are particularly important for providing a 658 comparison with models used to describe the response of vegetation to recent climate change 659 (Willis et al. 2010; Reitalu et al. 2014). For instance, the Quaternary record has indicated that 660 glacial-interglacial changes have induced large-scale shifts in plant distributions (Willis and Bhagwat 2009; Giesecke et al. 2017), although there was sometimes a lag between climatic 661 662 change and vegetation change (Leroy et al. 2011). Some warm-loving and cold-loving 663 deciduous tree species became extinct in Europe during glacial phases (Willis and Niklas 664 2004; Bertini 2010), due not only to climate change, but also to disease, competition and 665 extreme conditions in refugia (Leroy 2007). For those species that survived the glacial 666 phases, refugia such as in southern Europe were essential (Bennett et al. 1991; Leroy and 667 Arpe 2007). In contrast, conifers and some climatically less sensitive angiosperm trees found 668 refugia further north in Europe during glacial phases (crypto-refugia; Willis et al. 2000; 669 Bhagwat and Willis 2008) whereas herbaceous species typical of tundra and steppe 670 vegetation have been forced into upland refugia during the forest dominated phase of the 671 Holocene (Bennett and Provan 2008). It is evident that these refugia have been vital for the 672 shaping of present-day biogeographical patterns and the assemblage of extant communities 673 (Willis and Bhagwat 2009). Refugia have also been used to explain the responses of 674 vegetation to climate changes in Carboniferous tropical swamps (e.g. Falcon-Lang and 675 DiMichele 2010; Looy et al. 2014).

Palynological research has revealed anthropogenic effects on Holocene plant diversity
(Giesecke et al. 2012, 2019). For example, Filipova-Marinova et al. (2014) described an
8,000 year long record of vegetation change at Varna Lake (Bulgaria) and showed how the
vegetation was strongly influenced by human activity, both through woodland clearance and
the establishment of agriculture.

681 Rull (2011, 2013) has explored the drivers of neotropical diversity since early Neogene 682 times, and concluded that it is the result of complex ecological and evolutionary trends 683 initiated by tectonic events and palaeogeographical reorganisations, and was maintained by 684 Pleistocene climatic changes. The palynological record during the Palaeocene – Eocene 685 Thermal Maximum indicated an increase in diversity in tropical (Jaramillo et al. 2010) and 686 polar vegetation (Willard et al. 2019), whereas in temperate latitudes the effects were less 687 marked (e.g. Wing et al. 2003) except sometimes for a change to more fire-prone vegetation 688 (Collinson et al. 2009).

689 Further back in geological time, the analyses are more difficult because we are looking at 690 plants that are only distantly related to modern-day vegetation, but the comparisons can 691 nevertheless be insightful. Many Mesozoic studies have focussed on how vegetation 692 recovered from the Permian - Triassic and Triassic - Jurassic biotic crises (e.g. Grauvogel-693 Stamm and Ash 2005; Yu et al. 2015). For instance, Hochuli et al. (2016) showed the 694 complex pattern of recovery of the post-extinction, Early Triassic vegetation. Various other 695 floral changes were recorded at the Triassic – Jurassic boundary in Greenland (McElwain et 696 al. 2007) and at the Hettangian-Sinemurian boundary in the South Carpathians (Popa 2000). 697 In the Palaeozoic, most attention has been directed to the effect of the Late Palaeozoic 698 Ice Age on plant diversity: did the observed Pennsylvanian - Cisuralian diversity changes in 699 the tropical vegetation cause climate change (e.g. Cleal and Thomas 1999, 2005), or did the 700 climate change cause the vegetation changes (e.g. Pfefferkorn et al. 2008, 2017), or were the 701 two interlocked through feedback loops (Cleal et al. 2010)? Other links that have also been 702 explored are between the diversification of the first woody forests in Late Devonian times 703 and a significant change in ocean water chemistry that caused major reduction in marine 704 faunal diversity (Algeo and Scheckler 1998); and between the very early development of

plant diversity during Ordovician times and global cooling and glaciation (Servais et al.2019).

707 Another major theme of research has been the effect of mass extinctions on vegetation 708 (McElwain and Punyasena 2007; Cascales-Miñana et al. 2013). Clearly extensive destruction 709 of vegetation occurred during three of the five classic "mass extinctions" of Sepkoski (1978, 710 1979, 1984): at the boundaries between the Permian – Triassic (e.g. Looy et al. 1999; Hochuli et al. 2016, 2017), Triassic – Jurassic (e.g. McElwain et al 2007; Mander et al. 2013; 711 712 McElwain 2018) and Cretaceous - Palaeogene (e.g. Vajda and Bercovici 2014). However, a 713 key criterion for recognising a true mass extinction (Raup and Sepkoski 1982) is that it 714 should significantly disrupt the overall trajectory of evolution and this did not occur with 715 plants at most of these biotic crises (McElwain and Punyasena 2007; Cascales-Miñana 2011, 716 2012; Cascales-Miñana and Cleal 2011; Cascales-Miñana and Diez 2012; Cascales-Miñana et 717 al. 2013). Only at the Permian – Triassic boundary does there seem to have been any 718 significant clade disruption (Cascales-Miñana and Cleal 2014; Cascales-Miñana et al. 2016), 719 and even here the pattern of extinction was more complex than with the marine faunas 720 (Hochuli et al. 2016; Nowak et al. 2019). The fossil record seems to suggest that plants were 721 much less vulnerable to biotic crises compared with faunas (Traverse 1988; McElwain and 722 Punyasena 2007; McElwain et al. 2007; Cascales-Miñana et al. 2018). 723 Identifying vegetation diversity patterns at all scales from the plant fossil record is 724 clearly not easy; whatever the sampling protocol and analytical methods used, the fact will 725 remain that the observed diversity patterns are of the fossils rather than purely of the original vegetation. Nevertheless, the fossil record is the only direct evidence we have of how 726 727 vegetation has changed through time. By bringing together data from the palaeobotanical and 728 palynological records and interpreting it within the context of the taphonomic filter through 729 which the fossils have passed (Fig. 11) will allow us to understand better how plant-life has

730	responded to changes in climate, landscape and continental configurations, and to the
731	dramatic ecological crises often referred to as mass extinctions.

732 7 Conclusions

A deeper appreciation of the history of vegetation dynamics can inform present-day 733 734 landscape management and predictions of future biodiversity and climate. For example, the 735 plant fossil record can provide evidence of the speed at which plants can track climate change 736 and this may prove valuable to predict the response of today's plant vegetation to global 737 warming. It can also provide empirical data to help support and improve models of the 738 dynamic interactions of modern-day vegetation, atmosphere and climate. Evidence from the 739 fossil record clearly has the potential for making a significant contribution to understanding 740 the world today, and emphasises the importance of close co-operation between 741 palaeoecologists and ecologists. Provided that the context of the data is properly understood, 742 including the taxonomy of the fossils (not just the taxonomic names used but what the fossil-743 taxa mean), the taphonomic processes that caused the fossil to be preserved, and the detailed 744 temporal (stratigraphical) correlations, an underlying signal of vegetation diversity remains 745 waiting to be discovered.

746 Acknowledgements

The authors are grateful to the European Science Foundation for supporting the
Exploratory Workshop entitled *Tracking changes in plant diversity over the last 400 million years*, held in 2014 at National Museum Wales, Cardiff, UK. Ian Glasspool is thanked for
providing the images used in Fig. 3. This is a contribution to National Museum Wales
Research Project NS25 *Plant diversity through 450 million years of Earth history*.

752 8 References

- 753 Abbink, O.A., van Konijnenburg-van Cittert, J.H.A., Visscher, H., 2004. A sporomorph
- roup model for the Northwest European Jurassic-Lower Cretaceous: concepts and
- framework. Netherlands Journal of Geosciences 83, 17–31.
- 756 Algeo, T.J., Scheckler, S.E., 1998. Terrestrial-marine teleconnections in the Devonian: links
- between the evolution of land plants, weathering processes, and marine anoxic events.
- Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences
 353, 113–130.
- Alroy, J., 2003. Global databases will yield reliable measures of global biodiversity.
- 761 Paleobiology 29, 26–29.
- Andersen, S.T., 1970. The relative pollen productivity and pollen representation of north
- European trees, and correction factors for tree pollen spectra. Danmarks Geologiske
- 764 Undersögelse, Raekke 2, 1–99.
- Anderson, J.M., Anderson, H.M., Cleal, C.J., 2007. Brief history of the gymnosperms:
- real classification, biodiversity, phytogeography and ecology. Strelitzia 20, 1–279.
- Andrews, H.N., 1980. The fossil hunters. In search of ancient plants. Cornell University
 Press, Ithaca and London.
- Antonelli, A., Fry, C., Smith, R.J., Simmonds, M.S.J., Kersey, P.J., Pritchard, H.W., Abbo,
- 770 M.S., Acedo, C., Adams, J., Ainsworth, A.M., Allkin, B., Annecke, W., Bachman, S.P.,
- 771 Bacon, K., Bárrios, S., Barstow, C., Battison, A., Bell, E., Bensusan, K., Bidartondo, M.I.,
- 772 Blackhall-Miles, R.J., Borrell, J.S., Brearley, F.Q., Breman, E., Brewer, R.F.A., Brodie, J.,
- 773 Cámara-Leret, R., Campostrini Forzza, R., Cannon, P., Carine, M., Carretero, J., Cavagnaro,
- 774 T.R., Cazar, M.-E., Chapman, T., Cheek, M., Clubbe, C., Cockel, C., Collemare, J., Cooper,
- A., Copeland, A.I., Corcoran, M., Couch, C., Cowell, C., Crous, P., da Silva, M., Dalle, G.,

- 776 Das, D., David, J.C., Davies, L., Davies, N., De Canha, M.N., de Lirio, E.J., Demissew, S.,
- 777 Diazgranados, M., Dickie, J., Dines, T., Douglas, B., Dröge, G., Dulloo, M.E., Fang, R.,
- Farlow, A., Farrar, K., Fay, M.F., Felix, J., Forest, F., Forrest, L.L., Fulcher, T., Gafforov, Y.,
- 779 Gardiner, L.M., Gâteblé, G., Gaya, E., Geslin, B., Gonçalves, S.C., Gore, C.J.N., Govaerts,
- 780 R., Gowda, B., Grace, O.M., Grall, A., Haelewaters, D., Halley, J.M., Hamilton, M.A.,
- 781 Hazra, A., Heller, T., Hollingsworth, P.M., Holstein, N., Howes, M.-J.R., Hughes, M.,
- Hunter, D., Hutchinson, N., Hyde, K., Iganci, J., Jones, M., Kelly, L.J., Kirk, P., Koch, H.,
- 783 Krisai-Greilhuber, I., Lall, N., Langat, M.K., Leaman, D.J., Leão, T.C., Lee, M.A., Leitch,
- I.J., Leon, C., Lettice, E., Lewis, G.P., Li, L., Lindon, H., Liu, J.S., Liu, U., Llewellyn, T.,
- 785 Looney, B., Lovett, J.C., Łuczaj, Ł., Lulekal, E., Maggassouba, S., Malécot, V., Martin, C.,
- 786 Masera, O.R., Mattana, E., Maxted, N., Mba, C., McGinn, K.J., Metheringham, C., Miles, S.,
- 787 Miller, J., Milliken, W., Moat, J., Moore, P.G.P., Morim, M.P., Mueller, G.M., Muminjanov,
- H., Negrão, R., Nic Lughadha, E., Nicolson, N., Niskanen, T., Nono Womdim, R., Noorani,
- A., Obreza, M., O'Donnell, K., O'Hanlon, R., Onana, J.-M., Ondo, I., Padulosi, S., Paton, A.,
- 790 Pearce, T., Pérez Escobar, O.A., Pieroni, A., Pironon, S., Prescott, T.A.K., Qi, Y.D., Qin, H.,
- 791 Quave, C.L., Rajaovelona, L., Razanajatovo, H., Reich, P.B., Rianawati, E., Rich, T.C.G.,
- 792 Richards, S.L., Rivers, M.C., Ross, A., Rumsey, F., Ryan, M., Ryan, P., Sagala, S., Sanchez,
- 793 M.D., Sharrock, S., Shrestha, K.K., Sim, J., Sirakaya, A., Sjöman, H., Smidt, E.C., Smith, D.,
- 794 Smith, P., Smith, S.R., Sofo, A., Spence, N., Stanworth, A., Stara, K., Stevenson, P.C., Stroh,
- P., Suz, L.M., Tambam, B.B., Tatsis, E.C., Taylor, I., Thiers, B., Thormann, I., Trivedi, C.,
- 796 Twilley, D., Twyford, A.D., Ulian, T., Utteridge, T., Vaglica, V., Vásquez-Londoño, C.,
- 797 Victor, J., Viruel, J., Walker, B.E., Walker, K., Walsh, A., Way, M., Wilbraham, J., Wilkin,
- P., Wilkinson, T., Williams, C., Winterton, D., Wong, K.M., Woodfield-Pascoe, N.,
- 799 Woodman, J., Wyatt, L., Wynberg, R., Zhang, B.G., 2020. State of the world's plants and
- fungi 2020. Royal Botanic Gardens, Kew. DOI: https://doi.org/10.34885/172

- Appleton, P., Malpas, J., Thomas, B.A., Cleal, C.J., 2011. The Brymbo fossil forest. Geology
 Today 27, 107–113.
- 803 Baker, R.A., DiMichele, W.A., 1997. Biomass allocation in Late Pennsylvanian coal-swamp
- 804 plants. Palaios 12, 127–132.
- 805 Balme, B.E., 1995. Fossil in situ spores and pollen grains: an annotated catalogue. Review of
- 806 Palaeobotany and Palynology 87, 81–323.
- 807 Bambach, R.K., 1977. Species richness in marine benthic habitats through the Phanerozoic.
- 808 Paleobiology 3, 152–167.
- 809 Barbacka, M., 2012. Biodiversity and reconstruction of Early Jurassic flora from the Mecsek
- 810 Mountains (southern Hungary). Acta Palaeobotanica 51, 127–179.
- 811 Barbacka, M., Bodor, E., Jarzynka, A., Kustatscher, E., Pacyna, G., Popa, M.E., Scanu, G.G.,
- 812 Thévenard, F., Ziaja, J., 2014. European Jurassic floras: statistics and palaeoenvironmental
- 813 proxies. Acta Palaeobotanica 54, 173–195.
- 814 Barbacka, M., Popa, M.E., Mitka, J., Bodor, E., Püspöki, Z., McIntosh, R.W., 2016. A
- 815 quantitative approach for identifying plant ecogroups in the Romanian Early Jurassic
- 816 terrestrial vegetation. Palaeogeography, Palaeoclimatology, Palaeoecology 446, 44–54.
- 817 Barnekow, L., Loader, N.J., Hicks, S., Froyd, C.A., Goslar, T., 2007. Strong correlation
- 818 between summer temperature and pollen accumulation rates for *Pinus sylvestris*, *Picea abies*
- 819 and Betula spp. in a high-resolution record from northern Sweden Journal of Quaternary
- 820 Science 22, 653–658.
- 821 Bashforth, A.R., Falcon-Lang, H.J., Gibling, M.R., 2010. Vegetation heterogeneity on a Late
- 822 Pennsylvanian braided-river plain draining the Variscan Mountains, La Magdalena Coalfield,
- 823 northwestern Spain. Palaeogeography, Palaeoclimatology, Palaeoecology 292, 367–390.

- 824 Bashforth, A.R., Drábková, J., Opluštil, S., Gibling, M.R., Falcon-Lang, H.J., 2011.
- 825 Landscape gradients and patchiness in riparian vegetation on a Middle Pennsylvanian
- 826 braided-river plain prone to flood disturbance (Nýřany Member, Central and Western
- 827 Bohemian Basin, Czech Republic). Review of Palaeobotany and Palynology 163, 153–189.
- 828 Bateman, R.M., DiMichele, W.A., 1992. The rhizomorphic lycopsids: A case-study in
- paleobotanical classification. Sytematic Botany 21, 535–552.
- 830 Bateman, R.M., Hilton, J., 2009. Palaeobotanical systematics for the phylogenetic age:
- 831 applying organ-species, form-species and phylogenetic species concepts in a framework of
- reconstructed fossil and extant whole–plants. Taxon 58, 1254–1280.
- 833 Bateman, R.M., DiMichele, W.A., Willard, D.A., 1992. Experimental cladistics-analysis of
- anatomically preserved arborescent lycopsids from the Carboniferous of Euramerica An
- essay on palaeobotanical phylogenies. Annals of the Missouri Botanical Garden 79, 500–559.
- 836 Beerling, D., 2007. The emerald planet: how plants changed Earth's history. Oxford
- 837 University Press, Oxford.
- Bek, J., 2017. Paleozoic in situ spores and pollen. Lycopsida. Palaeontographica Abteilung B
 296, 1–111.
- Bennett, K.D., Provan, J., 2008. What do we mean by refugia? Quaternary Science Reviews
 27, 2449–2455.
- 842 Bennett, K.D., Tzedakis, P.C., Willis, K.J., Quaternary refugia of north European trees.
- S43 Journal of Biogeography 18, 103–115.
- 844 Bennington, J.B., Dimichele, W.A., Badgley, C., Bambach, R.K., Barrett, P.M.,
- 845 Behrensmeyer, A.K., Bobe, R., Burnham, R.J., Daeschler, E.B., Dam, J.V., Eronen, J.T., et
- al., 2009. Critical issues of scale in paleoecology. Palaios 24, 1–4.
- 847 Benton, M.J. (Ed.), 1993. The Fossil Record 2. Chapman and Hall, London.
- 848 Benton, M.J., 1995. Diversification and extinction in the history of life. Science 268, 52–58.
- 849 Beri, A., Martínez-Blanco, X., Tejera, L., Piñeyro, A., Souza, P.A., 2018. Palynodiversity
- 850 patterns and paleoclimatic changes in the late Paleozoic in Brazil and Uruguay. Boletín
- 851 Geológico y Minero 129, 599–614.
- 852 Beri, Á., Martínez-Blanco, X., Varela, L., di Pasquo, M., de Souza, P.A., 2020. Sampling
- 853 biases and Paleozoic sporomorphs diversity dynamics in Western Gondwana strata. Journal
- of South American Earth Sciences 98, doi.org/10.1016/j.jsames.2019.102457.
- 855 Berry, C.M., Marshall, J.E., 2015. Lycopsid forests in the early Late Devonian
- paleoequatorial zone of Svalbard. Geology 43, 1043–1046.
- Bertini, A., 2010. Pliocene to Pleistocene palynoflora and vegetation in Italy: state of the art.
 Quaternary International 225, 5–24.
- 859 Bhagwat, S.A., Willis, K.J., 2008. Species persistence in northerly glacial refugia of Europe:
- a matter of chance or biogeographical traits? Journal of Biogeography 35, 465–482.
- 861 Birks, H.H., Birks, H.J.B., 2000. Future uses of pollen analysis must include plant
- 862 macrofossils. Journal of Biogeography 27, 31–35.
- 863 Birks, H.H., Bjune, A.E., 2010. Can we detect a west-Norwegian treeline from modern
- 864 samples of plant remains and pollen? Results from the DOORMAT project. Vegetation
- 865 History and Archaeobotany 19, 325–340.
- 866 Birks, H.J.B., Birks, H.H., 1980. Quaternary Palaeoecology. Edward Arnold, London.
- 867 Birks, H.J.B., Felde, V.A., Seddon, A.W., 2016a. Biodiversity trends within the Holocene.
- 868 The Holocene 26, 994–1001.

- 869 Birks, H.J.B., Felde, V.A., Bjune, A.E., Grytnes, J.A., Seppä, H., Giesecke, T., 2016b. Does
- 870 pollen-assemblage richness reflect floristic richness? A review of recent developments and
- future challenges. Review of Palaeobotany and Palynology 228, 1–25.
- 872 Birks, H.J.B., Line, J.M., 1992. The use of rarefaction analysis for estimating palynological
- 873 richness from Quaternary pollen-analytical data. The Holocene 2, 1–10.
- Bjune, A.E., 2014. After 8 years of annual pollen trapping across the treeline in western
- Norway: are the data still anomalous? Vegetation History and Archaeobotany 23, 299–308.
- 876 Bless, M.J.M., Loboziak, S., Streel, M., 1977. An upper Westphalian C 'hinterland'
- 877 microflora from the Haaksbergen-1 Borehole (Netherlands). Mededelingen Rijks
- 878 Geologische Dienst 28, 135–147.
- 879 Brewer, S. Cheddadi, R., de Beaulieu, J.L., Reille, M., 2002. The spread of deciduous
- *Quercus* throughout Europe since the last glacial period. Forest Ecology and Management
 156, 27–48.
- 882 Brongniart, A., 1822. Sur la classification et la distribution des végétaux fossiles en général,
- 883 et sur ceux des terrains de sédiment supérieur en particulier. Introduction, Chapitre I.
- 884 Mémoires du Muséum d'Histoire Naturelle 8, 203–240.
- Bunting, M.J., Grant, M.J., Waller, M.P., 2016. Pollen signals of ground flora in managed
 woodlands. Review of Palaeobotany and Palynology 224, 121–133.
- 887 Burnham, R.J., 1989. Relationships between standing vegetation and leaf litter in a
- paratropical forest: implications for paleobotany. Review of Palaeobotany and Palynology 58,
 5–32.
- 890 Burnham, R.J., 1994. Paleoecological and floristic heterogeneity in the plant-fossil record: an
- analysis based on the Eocene of Washington. U.S. Geological Survey Bulletin 2085, B1-
- 892 B35.

- 893 Butler, R.J., Benson, R.B., Barrett, P.M., 2013. Pterosaur diversity: untangling the influence
- 894 of sampling biases, Lagerstätten, and genuine biodiversity signals. Palaeogeography,
- 895 Palaeoclimatology, Palaeoecology 372, 78–87.
- 896 Cabanes, D., Shahack-Gross, R., 2015. Understanding fossil phytolith preservation: The role
- of partial dissolution in paleoecology and archaeology. PLoS ONE 10(5), e0125532
- 898 Calder, J.H., Gibling, M.R., Scott, A.C., Davies, S.J., Hebert, B.L., Greb, S.F., DiMichele,
- 899 W.A., 2006. A fossil lycopsid forest succession in the classic Joggins section of Nova Scotia:
- 900 paleoecology of a disturbance-prone Pennsylvanian wetland, in: Greb, S.F., DiMichele, W.A.
- 901 (Eds), Wetlands through time. Geological Society of America Special Paper 399, pp. 169-
- 902 195.
- Campbell, I.D., McDonald, K., Flannigan, M.D., Kringayark, J., 1999. Long-distance
 transport of pollen into the Arctic. Nature 399, 29–30.
- 905 Carter, J.A., 1999. Late Devonian, Permian and Triassic phytoliths from Antarctica.
- 906 Micropaleontology 45, 56–61.
- 907 Cascales-Miñana, B., 2011. New insights into the reading of Paleozoic plant fossil record
- 908 discontinuities. Historical Biology 23, 115–130.
- 909 Cascales-Miñana, B., 2012. Disentangling temporal patterns in our perception of the fossil
- 910 history of gymnosperms. Historical Biology 24, 143–159.
- 911 Cascales-Miñana, B., Cleal, C.J., 2011. Plant fossil record and survival analysis. Lethaia 45,
 912 71–82.
- 913 Cascales-Miñana, B., Cleal, C.J., 2014. The plant fossil record reflects just two great
- 914 extinction events. Terra Nova 26, 195–200.

- 915 Cascales-Miñana, B., Gerrienne, P., 2017. Teruelia diezii gen. et sp. nov.: an early
- 916 polysporangiophyte from the Lower Devonian of the Iberian Peninsula. Palaeontology 60,917 199–212.
- 918 Cascales-Miñana, B., Cleal, C. J., Diez, J.B., 2013. What is the best way to measure
- 919 extinction? A reflection from the palaeobotanical record. Earth-Science Reviews 124, 126–
 920 147.
- 921 Cascales-Miñana, B., Diez, J.B., Gerrienne, P., Cleal, C.J., 2016. A palaeobotanical
- perspective on the great end-Permian biotic crisis. Historical Biology 28, 1066–1074.
- 923 Cascales-Miñana, B., Servais, T., Cleal, C.J., Gerrienne, P., Anderson, J., 2018. Plants-the
- great survivors! Geology Today 34, 224–229.
- 925 Cascales-Miñana, B., Diez, J.B., 2012. The effect of singletons and interval length on
- 926 interpreting diversity trends from the palaeobotanical record. Palaeontologia Electronica 15927 (1), 6A.
- 928 Chaloner, W.G., 1999. Taxonomic and nomenclatural alternatives, in: Jones, T.P., Rowe,
- N.P. (Eds.), Fossil plants and spores. Modern techniques. Geological Society, London, pp
 179–183.
- 931 Cleal, C.J., 2005. The Westphalian macrofloral record from the cratonic central Pennines
- Basin, UK. Zeitschrift der Deutschen Gesellschaft für Geowissenschaften 156, 387–410.
- 933 Cleal, C.J., 2007. The Westphalian-Stephanian macrofloral record from the South Wales
- 934 Coalfield. Geological Magazine 144, 465–486.
- 935 Cleal, C.J., 2008a. Westphalian-Stephanian macrofloras of the southern Pennines Basin, UK.
 936 Studia Geologica Polonica 129, 25–41.

- 937 Cleal, C.J., 2008b. Palaeofloristics of Middle Pennsylvanian lyginopteridaleans in Variscan
- 938 Euramerica. Palaeogeography, Palaeoclimatology, Palaeoecology 261, 1–14.
- 939 Cleal, C.J., 2008c. Palaeofloristics of Middle Pennsylvanian medullosaleans in Variscan
- 940 Euramerica. Palaeogeography, Palaeoclimatology, Palaeoecology 268, 164–180.
- 941 Cleal, C.J., 2019 (in press). Paleozoic plants. In Encyclopaedia of Earth sciences. Elsevier,
- 942 Amsterdam.
- 943 Cleal, C.J., Cascales-Miñana, B., 2014. Composition and dynamics of the great Phanerozoic
- Evolutionary Floras. Lethaia 47, 469–484.
- 945 Cleal, C.J., Rees, P.M., 2003. The Middle Jurassic flora from Stonesfield, Oxfordshire, UK.
- 946 Palaeontology 46, 739–801.
- 947 Cleal, C.J., Shute, C.H., 2012. The systematic and palaeoecological value of foliage anatomy
- 948 in Late Palaeozoic medullosalean seed-plants. Journal of Systematic Palaeontology 10, 765–
 949 800.
- Cleal, C.J., Thomas, B.A., 1988. The Westphalian fossil floras from the Cattybrook Claypit,
 Avon (Great Britain). Geobios 21, 409–433.
- Cleal, C.J., Thomas, B.A., 1999. Tectonics, tropical forest destruction and global warming in
 the Late Palaeozoic. Acta Palaeobotanica, Supplement 2, 17–19.
- 954 Cleal, C.J., Thomas, B.A., 2004. Late Carboniferous palaeobotany of the upper Bideford
- 955 Formation, north Devon: a coastal setting for a Coal Measures flora. Proceedings of the
- 956 Geologists' Association 115, 267–281.
- 957 Cleal, C.J., Thomas, B.A., 2005. Palaeozoic tropical rainforests and their effect on global
- 958 climates: is the past the key to the present? Geobiology 3, 13–31.

- 959 Cleal, C.J., Thomas, B.A., 2010. Botanical nomenclature and plant fossils. Taxon 59, 261–
 960 268.
- 961 Cleal, C.J., Thomas, B.A., 2019. Introduction to plant fossils. Second edition. Cambridge
- 962 University Press, Cambridge.
- 963 Cleal, C.J., Opluštil, S., Thomas, B.A., Tenchov, Y., 2010. Late Moscovian terrestrial biotas
- and palaeoenvironments of Variscan Euramerica. Netherlands Journal of Geosciences 88,
 181–278.
- 966 Cleal, C.J., Uhl, D., Cascales-Miñana, B., Thomas, B.A., Bashforth, A.R., King, S.C.,
- 967 Zodrow, E.L., 2012. Plant biodiversity changes in Carboniferous tropical wetlands. Earth-
- 968 Science Reviews 114, 124–155.
- 969 Close, R.A., Benson, R.B.J., Saupe, E.E., Clapham, M.E., Butler, R.J., 2020. The spatial
- 970 structure of Phanerozoic marine animal diversity. Science 368, 420-424.
- 971 Collinson, M.E., 1993. Taphonomy and fruiting biology of recent and fossil Nypa. Special
- 972 Papers in Palaeontology 49, 165–180.
- 973 Collinson, M.E., 1996. "What use are fossil ferns?" 20 years on: with a review of the fossil
- 974 history of extant pteridophyte families and genera, in: Camus, M.G., Johns, R.J. (Eds.),
- 975 Pteridology in perspective. Royal Botanic Gardens, Kew, pp. 349–394.
- 976 Collinson, M.E., Manchester, S.R., Wilde, V., 2012. Fossil fruits and seeds of the Middle
- 977 Eocene Messel biota, Germany. Abhandlungen der Senckenberg Gesellschaft für
- 978 Naturforschung 570, 251 pp.
- 979 Collinson, M.E., Steart, D.C., Harrington, G.J., Hooker, J.J., Scott, A.C., Allen, L.O.,
- 980 Glasspool, I.J., Gibbons, S.J., 2009. Palynological evidence of vegetation dynamics in
- 981 response to palaeoenvironmental change across the onset of the Paleocene-Eocene Thermal
- 982 Maximum at Cobham, Southern England. Grana 48, 38–66.

- 983 Costamagna, L.G., Kustatscher, E., Scanu, G.G., Del Rio, M., Pittau, P., van Konijnenburg-
- van Cittert, J.H., 2018. A palaeoenvironmental reconstruction of the Middle Jurassic of
- 985 Sardinia (Italy) based on integrated palaeobotanical, palynological and lithofacies data
- 986 assessment. Palaeobiodiversity and Palaeoenvironments 98, 111–138.
- 987 Davies, N.S., Gibling, M.R., 2010. Cambrian to Devonian evolution of alluvial systems: the
- 988 sedimentological impact of the earliest land plants. Earth-Science Reviews 98, 171–200.
- Davis, M.B., 1963. On the theory of pollen analysis. American Journal of Science 261, 897–
 990 912.
- 991 Diez, J.B., Sender, L.M., Villanueva-Amadoz, U., Ferrer, J., Rubio, C., 2005. New data
- 992 regarding Weichselia reticulata: Soral clusters and the spore developmental process. Review
- 993 of Palaeobotany and Palynology 135, 99–107
- DiMichele, W.A., Falcon-Lang, H.J., 2011. Pennsylvanian 'fossil forests' in growth position
- 995 (T⁰ assemblages): origin, taphonomic bias and palaeoecological insights. Journal of the
- 996 Geological Society, London 168, 585–605.
- DiMichele, W.A., Gastaldo, R.A., 2008. Plant paleoecology in deep time. Annals of the
 Missouri Botanical Garden 95, 144–198.
- 999 DiMichele, W.A., Nelson, W.J., 1989. Small-scale spatial heterogeneity in Pennsylvanian-
- age vegetation from the roof shale of the Springfield Coal (Illinois Basin). Palaios 4, 276–
- 1001 280.
- 1002 DiMichele, W.A., Phillips, T.L., 1988. Paleoecology of the Middle Pennsylvanian-age Herrin
- 1003 coal swamp (Illinois) near a contemporaneous river system, the Walshville paleochannel.
- 1004 Review of Palaeobotany and Palynology 56, 151–176.

- DiMichele, W.A., Philips, T.L., McBrinn, G.E., 1991. Quantitative analysis and paleoecology
 of the Secor coal and roof shale floras (Middle Pennsylvanian, Oklahoma). Palaios 6, 390–
 409.
- 1008 DiMichele, W.A., Phillips, T.L., Nelson, W.J., 2002. Place vs. time and vegetational
- 1009 persistence: a comparison of four tropical mires from the Illinois Basin during the height of
- 1010 the Pennsylvanian Ice Age. International Journal of Coal Geology 50, 43–72.
- 1011 DiMichele, W.A., Falcon-Lang, H.J., Nelson, W.J., Elrick, S.D., Ames, P.R., 2007.
- 1012 Ecological gradients within a Pennsylvanian mire forest. Geology 35, 415–418.
- 1013 DiMichele, W.A., Elrick, S.D., Nelson, W.J., 2017. Vegetational zonation in a swamp forest,
- 1014 Middle Pennsylvanian, Illinois Basin, USA, indicates niche differentiation in a wetland plant
- 1015 community. Palaeogeography, Palaeoclimatology, Palaeoecology 487, 71–92.
- 1016 DiMichele, W.A., Hook, R.W., Kerp, H., Hotton, C.L., Looy, C.V., Chaney, D.S., 2018.
- 1017 Lower Permian Flora of the Sanzenbacher Ranch, Clay County, Texas, in: Krins, M., Harper,
- 1018 C.J., Cúneo, R., Rothwell, G.E. (eds.), Transformative paleobotany: papers to commemorate
- 1019 the life and legacy of Thomas N. Taylor. Elsevier, Amsterdam, pp. 95–126.
- 1020 Dimitrova, T.K., Cleal, C.J., 2007. Palynological evidence for late Westphalian-early
- 1021 Stephanian vegetation change in the Dobrudzha Coalfield, NE Bulgaria. Geological
- 1022 Magazine 144, 513–524.
- 1023 Dimitrova, T.K., Cleal, C.J., Thomas, B.A., 2005. Palynology of late Westphalian early
- Stephanian coal-bearing deposits in the eastern South Wales Coalfield. Geological Magazine
 142, 809–821.
- 1026 Dimitrova, T.K., Cleal, C.J., Thomas, B.A., 2011. Palynological evidence for Pennsylvanian
- 1027 extra-basinal vegetation in Atlantic Canada. Journal of the Geological Society 168, 559–569.

- 1028 Dimitrova, T.K., Zodrow, E.L., Cleal, C.J., Thomas, B.A., 2010. Palynological evidence for
- 1029 Pennsylvanian (Late Carboniferous) vegetation change in the Sydney Coalfield, eastern
- 1030 Canada. Geological Journal 45, 388–396.
- 1031 Dornelas, M., Magurran, A.E., Buckland, S.T., Chao, A., Chazdon, R.L., Colwell, R.K.,
- 1032 Curtis, T., Gaston, K.J., Gotelli, N.J., Kosnik, M.A., McGill, B., 2012. Quantifying temporal
- 1033 change in biodiversity: challenges and opportunities. Proceedings of the Royal Society B:
- 1034 Biological Sciences 280, 20121931
- 1035 Eble, C.F., Greb, S.F., Williams, D.A., Hower, J.C., O'Keefe, J.M., 2019. Palynology,
- 1036 organic petrology and geochemistry of the Bell coal bed in Western Kentucky, Eastern
- 1037 Interior (Illinois) Basin, USA. International Journal of Coal Geology 213, 103264.
- 1038 Edwards, D., Kenrick, P., Dolan, L., 2018, History and contemporary significance of the
- 1039 Rhynie cherts-our earliest preserved terrestrial ecosystem. Philosophical Transactions of the
- 1040 Royal Society of London. Series B 373(1739), 20160489.
- Enquist, B.J., Haskell, J.P., Tiffney, B.H., 2002. General patterns of taxonomic and biomass
 partitioning in extant and fossil plant communities. Nature 419, 610–613.
- 1043 Ertl, C., Pessi, A.M., Huusko, A., Hicks, S., Kubin, E., Heino, S., 2012, Assessing the
- 1044 proportion of "extra-local" pollen by means of modern aerobiological and phenological
- 1045 records An example from Scots pine (Pinus sylvestris L.) in northern Finland. Review of
- 1046 Palaeobotany and Palynology 185, 1–12.
- 1047 Falcon-Lang, H.J., Bashforth, A.R., 2005. Morphology, anatomy, and upland ecology of
- 1048 large cordaitalean trees from the Middle Pennsylvanian of Newfoundland. Review of
- 1049 Palaeobotany and Palynology 135, 223–243.
- 1050 Falcon-Lang, H.J. and DiMichele, W.A., 2010. What happened to the coal forests during
- 1051 Pennsylvanian glacial phases? Palaios 25, 611–617.

- 1052 Falcon-Lang, H.J., Kurzawe, F., Lucas, S.G., 2016. A Late Pennsylvanian coniferopsid forest
- 1053 in growth position, near Socorro, New Mexico, USA: tree systematics and palaeoclimatic
- 1054 significance. Review of Palaeobotany and Palynology 225, 67–83.
- 1055 Ferguson, D.K., 1985. The origin of leaf assemblages new light on an old problem. Review
- 1056 of Palaeobotany and Palynology 46, 117–188.
- Ferguson, D.K., 2005. Plant taphonomy: ruminations on the past, the present, and the future.Palaios 20, 418–428.
- 1059 Ferguson, D.K., Zetter, R., Paudayal, K.N., 2007. The need for SEM in palaeopalynology.
- 1060 Comptes Rendus Palévol 6, 423–430.
- 1061 Filipova-Marinova, M.V., Kvavadze, E.V., Connor, S.E., Sjögren, P., 2010. Estimating
- absolute pollen productivity for some European Tertiary-relict taxa. Vegetation History and
 Archaeobotany 19, 351–364.
- 1064 Francis, J.E., 1983. The dominant conifer of the Jurassic Purbeck formation, England.
- 1065 Palaeontology 26, 277–294.
- 1066 Franz, M., Kustatscher, E., Heunisch, C., Niegel, S., Röhling, H.G., 2019. The
- 1067 Schilfsandstein and its flora; arguments for a humid mid-Carnian episode? Journal of the
- 1068 Geological Society 176, 133–148.
- 1069 Galtier, J., 1986. Taxonomic problems due to preservation: comparing compression and
- 1070 permineralised taxa, in: Spicer, R.A., Thomas, B.A (Eds.), Systematic and taxonomic
- 1071 approaches in palaeobotany. Oxford University Press, Oxford (Systematics Association,
- 1072 Special Volume 31), pp. 1–16.
- 1073 Galtier, J., 1997. Coal-ball floras of the Namurian-Westphalian of Europe. Review of
- 1074 Palaeobotany and Palynology 95, 51–72.

- 1075 Galtier, J., 2008. A new look at the permineralized flora of Grand-Croix (Late Pennsylvanian,
- 1076 Saint-Etienne basin, France). Review of Palaeobotany and Palynology 152, 129–140.
- 1077 Gandolfo, M.A., Nixon, K.C., Crepet, W.L., Grimaldi, D.A., 2018. A late Cretaceous
- 1078 fagalean inflorescence preserved in amber from New Jersey. American Journal of Botany
- 1079 105, 1424–1435.
- 1080 Garwood, R.J., Oliver, H., Spencer, A.R.T., 2019. An introduction to the Rhynie chert.
- 1081 Geological Magazine 157, 47–64.
- 1082 Gastaldo, R.A., 1985. Implications on the paleoecology of autochthonous lycopods in clastic
- 1083 sedimentary environments of the Early Pennsylvanian of Alabama. Palaeogeography,
- 1084 Palaeoclimatology, Palaeoecology 53, 191–212.
- Gastaldo, R.A. 1992. Taphonomic considerations for plant evolutionary investigations. The
 Palaeobotanist 41, 211–223.
- 1087 Gastaldo, R.A., Demko, T.M., 2011. The relationship between continental landscape
- 1088 evolution and the plant-fossil record: long term hydrologic controls on preservation, in:
- Allison, P.A., Bottjer, D.J. (Eds.), Taphonomy. Process and bias through time. Springer
 Netherlands, pp. 249-285.
- 1091 Gastaldo, R.A., Douglass, D.P., McCarroll, S.M., 1987. Origin, characteristics, and
- 1092 provenance of plant macrodetritus in a Holocene crevasse splay mobile delta, Alabama.
- 1093 Palaios 2, 229–240.
- 1094 Gastaldo, R.A., Huc, A.-Y., 1992. Sediment facies, depositional environments, and
- 1095 distribution of phytoclasts in the Recent Mahakam River Delta, Kalimantan, Indonesia.
- 1096 Palaios 7, 574–590.

Gastaldo, R.A., Pfefferkorn, H.W., DiMichele, W.A., 1995. Taphonomic and sedimentologic
characterization of roof-shale floras. Memoirs of the Geological Society of America 185,
341–352.

- 1100 Gerasimidis, A., Panajiotidis, S., Hicks, S., Athanasiadis, N., 2006. An eight-year record of
- 1101 pollen deposition in the Pieria mountains (N. Greece) and its significance for interpreting
- 1102 fossil pollen assemblages. Review of Palaeobotany and Palynology 141, 231–243.
- 1103 Giesecke, T., Ammann, B., Brande, A., 2014. Palynological richness and evenness: insights
- 1104 from the taxa accumulation curve. Vegetation History and Archaeobotany 23, 217–228.
- 1105 Giesecke, T., Bennett, K.D., 2004. The Holocene spread of Picea abies (L.) Karst. in
- 1106 Fennoscandia and adjacent areas. Journal of Biogeography 31, 1–26.
- 1107 Giesecke, T., Hickler, T., Kunkel, T., Sykes, M.T., Bradshaw R.H.W., 2007. Towards an
- understanding of the Holocene distribution of *Fagus sylvatica* L. Journal of Biogeography 34,
 1109 118–131.
- 1110 Giesecke, T., Fontana, S.L., van der Knaap, W.O., Pardoe, H.S., Pidek, I.A., 2010. From
- 1111 early pollen trapping experiments to the pollen monitoring programme. Vegetation History
- 1112 and Archaeobotany 19, 247–258.
- 1113 Giesecke, T., Wolters, S., Jahns, S., Brande, A., 2012. Exploring Holocene changes in
- 1114 palynological richness in Northern Europe did postglacial immigration matter? PLos One 7
- 1115 e51624. Doi:10.1371/journal.pone.0051624.
- 1116 Giesecke, T., Davis, B., Brewer, S. Finsinger, W., Wolters, S., Blaauw, M., de Beaulieu, J.-
- 1117 L., Binney, H., Fyfe, R.M., Gaillard, M.-J., Gil-Romera, G., van der Knaap, W.O., Kuneš, P.,
- 1118 Kühl, N., van Leeuwen, J.F.N., Leydet, M., Lotter, A.F., Ortu, E., Semmler, M., Bradshaw,
- 1119 R.H.W., 2014. Towards mapping the late Quaternary vegetation change of Europe.
- 1120 Vegetation History and Archaeobotany 23, 75–86.

- 1121 Giesecke, T., Wolters, S., van Leeuwen, J.F., van der Knaap, P.W., Leydet, M., Brewer, S.,
- 1122 2019. Postglacial change of the floristic diversity gradient in Europe. Nature
- 1123 Communications 10(1), 1-7.
- 1124 Glasspool, I.J., 2003. Hypautochthonous–allochthonous coal deposition in the Permian,
- 1125 South African, Witbank Basin No. 2 seam; a combined approach using sedimentology, coal
- 1126 petrology and palaeontology. International Journal of Coal Geology 53, 81–135.
- 1127 Glasspool, I.J., Hilton, J., Collinson, M.E., Wang, S.J., 2004. Foliar physiognomy in
- 1128 Cathaysian gigantopterids and the potential to track Palaeozoic climates using an extinct plant
- 1129 group. Palaeogeography, Palaeoclimatology, Palaeoecology 205, 69–110.
- 1130 Gomez, B., Gillot, T., Daviero-Gomez, V., Coiffard, C., Spagna, P., Yans, J., 2012.
- 1131 Mesofossil plant remains from the Barremian of Hautrage (Mons Basin, Belgium), with
- 1132 taphonomy, paleoecology, and paleoenvironment insights, in: Godefroit, P. (Ed.), Bernissart
- 1133 dinosaurs and Early Cretaceous terrestrial ecosystems. Indiana University Press,
- 1134 Bloomington, pp. 97–112.
- 1135 Goswami, S., Singh, K.J., 2013. Floral biodiversity and geology of the Talcher Basin, Orissa,
- 1136 India during the Permian–Triassic interval. Geological Journal 48, 39–56.
- 1137 Goswami, S., Saxena, A., Singh, K.J., Chandra, S., Cleal, C.J., 2018. An appraisal of the
- 1138 Permian palaeobiodiversity and geology of the Ib-River Basin, eastern coastal area, India.
- 1139 Journal of Asian Earth Sciences 157, 283–301.
- 1140 Grauvogel-Stamm, L., Ash, S.R., 2005. Recovery of the Triassic land flora from the end-
- 1141 Permian life crisis. Comptes Rendus Palevol 4, 593–608.
- 1142 Grímsson, F., Zetter, R., 2011a. Combined LM and SEM study of the Middle Miocene
- 1143 (Sarmatian) palynoflora from the Lavanttal Basin, Austria: Part II. Pinophyta (Cupressaceae,
- 1144 Pinaceae and Sciadopityaceae). Grana 50, 262–310.

- 1145 Grímsson, F., Zetter, R., Baal C., 2011b. Combined LM and SEM study of the Middle
- 1146 Miocene (Sarmatian) palynoflora from the Lavanttal Basin, Austria: Part I. Bryophyta,
- 1147 Lycopodiophyta, Pteridophyta, Ginkgophyta, and Gnetophyta. Grana 50, 102–28.
- 1148 Grímsson, F., Grimm, G.W., Meller, B., Bouchal, J.M., Zetter, R., 2015a. Combined LM and
- 1149 SEM study of the middle Miocene (Sarmatian) palynoflora from the Lavanttal Basin, Austria:
- 1150 Part IV. Magnoliophyta 2 Fagales to Rosales. Grana 55, 101–63.
- 1151 Grímsson, F., Meller, B., Bouchal, J.M., Zetter, R., 2015b. Combined LM and SEM study of
- 1152 the middle Miocene (Sarmatian) palynoflora from the Lavanttal Basin, Austria: part III.
- 1153 Magnoliophyta 1 Magnoliales to Fabales. Grana 54, 85–128.
- 1154 Grímsson, F., Zetter, R., Labandeira, C.C., Engel, M.S., Wappler, T., 2017. Taxonomic
- description of *in situ* bee pollen from the middle Eocene of Germany. Grana 56: 37–70.
- 1156 Habib, D., Groth, P.K., 1967. Paleoecology of migrating Carboniferous peat environments.
- 1157 Palaeogeography, Palaeoclimatology, Palaeoecology 3, 185–195.
- 1158 Hamilton, A.J., 2005. Species diversity or biodiversity? Journal of Environmental
- 1159 Management 75, 89–92.
- Harland, W.B. (Ed.), 1967. The Fossil Record: a symposium with documentation. GeologicalSociety, London.
- 1162 Herendeen, P.S., Crepet, W.L., Nixon, K.C., 1994. Fossil flowers and pollen of Lauraceae
- 1163 from the Upper Cretaceous of New Jersey. Plant Systematics and Evolution 189, 29–40.
- 1164 Heyworth, A., Kidson, C., 1982. Sea-level changes in southwest England and Wales.
- 1165 Proceedings of the Geologists' Association 93, 91–111.
- 1166 Hicks, S., 1985. Modern pollen deposition records from Kuusamo, Finland I. Seasonal and
- 1167 annual variation. Grana 24,167–184.

- Hicks, S., 2001. The use of annual arboreal pollen deposition values for delimiting tree-lines
 in the landscape and exploring models of pollen dispersal. Review of Palaeobotany and
 Palynology 117, 1–29.
- 1171 Hicks, S., Ammann, B., Latałowa, M., Pardoe, H., Tinsley, H., 1996. European Pollen
- 1172 Monitoring Programme: Project Description and Guidelines. Oulu University Press, Oulu.
- 1173 Hicks, S., Helander, M., Heino, S., 1994. Birch pollen production, transport and deposition
- 1174 for the period 1984-1993 at Kevo, Finland. Aerobiologia 10, 183–191.
- 1175 Hilton, J., Cleal, C.J., 2007. The relationship between Euramerican and Cathaysian tropical
- 1176 floras in the Late Palaeozoic: palaeobiogeographical and palaeogeographical implications.
- 1177 Earth–Science Reviews 85, 85–116.
- 1178 Hochuli, P.A., Sanson-Barrera, A., Schneebeli-Hermann, E., Bucher, H., 2016. Severest crisis
- 1179 overlooked—Worst disruption of terrestrial environments postdates the Permian–Triassic
- 1180 mass extinction. Nature Scientific Reports, 6:28372 | DOI: 10.1038/srep28372.
- 1181 Hochuli, P.A., Schneebeli-Hermann, E., Mangerud, G., Bucher, H. 2017. Evidence for
- atmospheric pollution across the Permian-Triassic transition. Geology 45, 1123–1126.
- 1183 Hofmann, C.-C., Gregor, H.J., 2018. Scanning electron microscope investigations of pollen
- 1184 from an atypical mid-Eocene coal facies in Stolzenbach mine (PreußenElektra) near Borken
- 1185 (Kassel, Lower Hesse, Germany). Review of Palaeobotany and Palynology 252, 41–63.
- Holland, S.M., Sclafani, J.A., 2015. Phanerozoic diversity and neutral theory. Paleobiology41, 369–376.
- 1188 Huang, Y., Jia, L., Wang, Q., Mosbrugger, V., Utescher, T., Su, T., Zhou, Z., 2016. Cenozoic
- 1189 plant diversity of Yunnan: a review. Plant Diversity 38, 271–282.
- Hughes, N.F., 1963. The assignment of species of fossils to genera. Taxon 12, 336–337.

- 1192 Huntley, B., Birks, H.J.B., 1983. An Atlas of Past and Present Pollen Maps for Europe: 0-
- 1193 13000 Years Ago. Cambridge University Press, Cambridge.
- 1194 Jantz, N., Homeier, J., Behling, H., 2014. Representativeness of tree diversity in the modern
- pollen rain of Andean montane forests. Journal of Vegetation Science 25, 481–490.
- 1196 Jaramillo, C., Ochoa, D., Contreras, L., Pagani, M., Carvajal-Ortiz, H., Pratt, L.M., Krishnan,
- 1197 S., Cardona, A., Romero, M., Quiroz, L., Rodriguez, G., et al. 2010. Effects of rapid global
- 1198 warming at the Paleocene-Eocene boundary on neotropical vegetation. Science 330, 957–
- 1199 961.
- 1200 Jasper, K., Hartkopf-Fröder, C., Flajs, G., Littke, R., 2010. Palaeoecological evolution of
- 1201 Duckmantian wetlands in the Ruhr Basin (western Germany): A palynological and coal
- 1202 petrographical analysis. Review of Palaeobotany and Palynology 162, 123–145.
- 1203 Johnson, K.R., 2002. The megaflora of the Hell Creek and lower Fort Union formations in
- 1204 the western Dakotas: Vegetational response to climate change, the Cretaceous-Tertiary
- 1205 boundary event, and rapid marine transgression, in: Hartman, J.H., Johnson, K.R., Nichols,
- 1206 D.J. (Eds.), The Hell Creek Formation and the Cretaceous-Tertiary boundary in the northern
- 1207 Great Plains: An integrated continental record of the end of the Cretaceous. Geological
- 1208 Society of America, Special Papers 361, 329–391.
- 1209 Johnston, M.N., Eble, C.F., O'Keefe, J.M., Freeman, R.L., Hower, J.C., 2017. Petrology and
- 1210 palynology of the Middle Pennsylvanian Leatherwood coal bed, Eastern Kentucky:
- 1211 Indications for depositional environments. International Journal of Coal Geology 181, 23–38.
- 1212 Kędzior, A., Popa, M.E., 2013. Sedimentology of the Early Jurassic terrestrial Steierdorf
- 1213 Formation in Anina, Colonia Cehă Quarry, South Carpathians, Romania. Acta Geologica
- 1214 Polonica 63, 175–199.

- 1215 Kędzior, A., Popa, M.E., 2018. An Early Jurassic braided river system from Mehadia, South
- 1216 Carpathians, Romania. Geological Quarterly 62, 415–432.
- 1217 Khuroo, A.A., Dar, G.H., Khan, Z.S., Malik, A.H., 2007. Exploring an inherent interface
- 1218 between taxonomy and biodiversity: current problems and future challenges. Journal for
- 1219 Nature Conservation 15, 256–261.
- 1220 Kidwell, S.M., Holland, S.M., 2002. The quality of the fossil record: implications for
- 1221 evolutionary analyses. Annual Review of Ecology and Systematics 33, 561–88.
- 1222 Knoll, A.H., Niklas, K.J., Tiffney, B.H., 1979. Phanerozoic land-plant diversity in North
- 1223 America. Science 206, 1400–1402.
- 1224 Köhler, J., Uhl, D., 2014. Die Blatt- und Karpoflora der oberoligozänen Fossillagerstätte
- 1225 Enspel (Westerwald, Rheinland-Pfalz, W-Deutschland). Mainzer naturwissenschaftliches
 1226 Archiv, Beihefte 35, 1–87.
- 1227 Kowalewski, M., Kiessling, W., Aberhan, M., Fürsich, F.T., Scarponi, D., Wood, S.L.B.,
- 1228 Hoffmeister, A.P., 2006. Ecological, taxonomic, and taphonomic components of the post-
- 1229 Paleozoic increase in sample-level species diversity of marine benthos. Paleobiology 32,
- 1230 533–561.
- 1231 Kustatscher, E., van Konijnenburg-van Cittert, J.H.A., Roghi, G., 2010. Macrofloras and
- 1232 palynomorphs as possible proxies for palaeoclimatic and palaeoecological studies: A case
- 1233 study from the Pelsonian (Middle Triassic) of Kühwiesenkopf/Monte Prà della Vacca (Olang
- 1234 Dolomites, N-Italy). Palaeogeography, Palaeoclimatology, Palaeoecology 290, 71–80.
- 1235 Kustatscher, E., Bernardi, M., Petti, F.M., Franz, M., van Konijnenburg-van Cittert, J.H.A.,
- 1236 Kerp, H., 2017. Sea-level changes in the Lopingian (late Permian) of the north-western
- 1237 Tethys and their effects on the terrestrial palaeoenvironments, biota and fossil preservation.
- 1238 Global and Planetary Change 148, 166–180.

- 1239 Lande, R., 1996. Statistics and partitioning of species diversity, and similarity among
- 1240 multiple communities. Oikos 76, 5–13.
- 1241 Latałowa, M., van der Knaap, W.O., 2006. Late Quaternary expansion of Norway spruce
- 1242 Picea abies (L.) Karst. in Europe according to pollen data. Quaternary Science Reviews 25,
- 1243 2780–2805.
- 1244 La Torre, M.D.L.Á., Herrando-Pérez, S., Young, K.R., 2007. Diversity and structural patterns
- 1245 for tropical montane and premontane forests of central Peru, with an assessment of the use of
- 1246 higher-taxon surrogacy. Biodiversity and Conservation 16, 2965–2988.
- 1247 Laveine, J.P., Belhis, A., 2007. Frond architecture of the seed-fern Macroneuropteris
- 1248 scheuchzeri, based on Pennsylvanian specimens from the Northern France coal field.
- 1249 Palaeontographica, Abteilung B 277, 1–41.
- 1250 Leroy, S.A.G., 1992. Palynological evidence of *Azolla nilotica* Dec. in recent Holocene of
- eastern Nile Delta, and its environment. Vegetation History and Archaeobotany 1, 43-52.
- 1252 Leroy, S.A.G., Arpe, K., 2007. Glacial refugia for summer-green trees in Europe and south-
- 1253 west Asia as proposed by ECHAM3 time-slice atmospheric model simulations. Journal of
- 1254 Biogeography 34, 2115–2128.
- 1255 Leroy, S.A.G., Arpe, K., Mikolajewicz, U., 2011. Vegetation context and climatic limits of
- the early Pleistocene hominin dispersal in Europe. Quaternary Science Reviews 30, 1448–
 1463.
- 1258 Leroy, S.A.G., Boyraz, S., Gürbüz, A., 2009. High-resolution palynological analysis in Lake
- 1259 Sapanca as a tool to detect earthquakes on the North Anatolian Fault. Quaternary Science
- 1260 Reviews 28, 2616–2632.
- 1261 Leroy, S.A.G., Roiron, P., 1996. Final Pliocene macro and micro floras of Bernasso
- 1262 (Escandorgue, France). Review of Palaeobotany and Palynology 94, 295–328.

- 1263 Libertín, M., Opluštil, S., Pšenièka, J., Bek, J., Sýkarová, I., Daškova, J., 2009a. Middle
- 1264 Pennsylvanian pioneer plant assemblage buried in situ by volcanic ash-fall, central Bohemia,
- 1265 Czech Republic. Review of Palaeobotany and Palynology 155, 204–233.
- 1266 Libertín, M., Dašková, J., Opluštil, S., Bek, J., Edress, N., 2009b. A palaeoecological model
- 1267 for a vegetated early Westphalian intramontane valley (Intra-Sudetic Basin, Czech
- 1268 Republic). Review of Palaeobotany and Palynology 155, 175–203.
- Lidgard, S., Crane, P.R., 1990. Angiosperm diversification and Cretaceous floristic trends: a
 comparison of palynofloras and leaf macrofloras. Paleobiology 16, 77–93.
- 1271 Litsitsyna, O.V, Hicks, S. Huusko, A., 2012. Do moss samples, pollen traps and modern lake
- sediments all collect pollen in the same way? Vegetation History and Archaeobotany 21,
- 1273 187–199.
- 1274 Locatelli, E.R., Krajewski, L., Chochinov, A.V., Laflamme, M., 2016. Taphonomic variance
- 1275 between marattialean ferns and medullosan seed ferns in the Carboniferous Mazon Creek
- 1276 Lagerstätte, Illinois, USA. Palaios 31, 97–110.
- 1277 Looy, C.V., Brugman, W.A., Dilcher, D.L., Visscher, H., 1999. The delayed resurgence of
- 1278 equatorial forests after the Permian–Triassic ecologic crisis. Proceedings of the National
- 1279 Academy of Sciences 96, 13857–13862.
- 1280 Looy, C.V., Stevenson, R.A., van Hoof, T.B., Mander, L., 2014. Evidence for coal forest
- refugia in the seasonally dry Pennsylvanian tropical lowlands of the Illinois Basin, USA.
- 1282 PeerJ 2, p.e630.
- 1283 López-Merino, L., Leroy, S.A.G., Eshel, A., Epshtein, V., Belmaker, R., Bookman, R., 2016.
- 1284 Using palynology to re-assess the Dead Sea laminated sediments indeed varves? Quaternary
- 1285 Science Reviews 140, 49–66.

- 1286 Luthardt, L., Rößler, R., Schneider, J.W., 2016. Palaeoclimatic and site-specific conditions in
- 1287 the early Permian fossil forest of Chemnitz Sedimentological, geochemical and
- palaeobotanical evidence. Palaeogeography, Palaeoclimatology, Palaeoecology 441, 627–
 652.
- 1290 McElwain, J.C., 2018. Paleobotany and global change: Important lessons for species to
- biomes from vegetation responses to past global change. Annual Review of Plant Biology 69,761–787.
- 1293 McElwain, J.C., Punyasena, S.W., 2007. Mass extinction events and the plant fossil record.
- 1294 Trends in Ecology and Evolution 22, 548–557.
- 1295 McElwain, J.C., Popa, M.E., Hesselbo, S.P., Haworth, D.M., Surlyk, F., 2007.
- 1296 Macroecological responses of terrestrial vegetation to climatic and atmospheric change across
- 1297 the Triassic/Jurassic boundary in East Greenland. Paleobiology 33, 547–573.
- 1298 Magurran, A.E., 2004. Measuring biological diversity. Wiley-Blackwell, Oxford.
- 1299 Mander, L., Punyasena, S.W., 2014. On the taxonomic resolution of pollen and spore records
- 1300 of Earth's vegetation. International Journal of Plant Sciences 175, 931–945.
- 1301 Mander, L., Kürschner, W.M., McElwain, J.C., 2013. Palynostratigraphy and vegetation
- 1302 history of the Triassic–Jurassic transition in East Greenland. Journal of the Geological
- 1303 Society 170, 37–46.
- 1304 Mauquoy, D., Hughes, P.D.M., Van Geel, B., 2010. A protocol for plant macrofossil analysis
- 1305 of peat deposits. Mires and Peat 7, 1–5.
- 1306 Meyen, S.V., 1984. Basic features of gymnosperm systematics and phylogeny as evidenced
- 1307 by the fossil record. Botanical Review 50, 1–111.

- 1308 Mitsumoto, K., Yabusaki, K., Aoyagi, H., 2009. Classification of pollen species using
- 1309 autofluorescence image analysis. Journal of Bioscience and Bioengineering 107, 90–94.
- 1310 Moir, A., Leroy, S.A.G., Brown, D., Collins, P., 2010. Dendrochronological evidence for a
- 1311 lower water table on peatland around 3200-3000 BC from sub-fossil pine in northern
- 1312 Scotland. The Holocene 20, 931–942.
- 1313 Niklas, K.J., Tiffney, B.H., 2010. The quantification of plant biodiversity through time.
- 1314 Philosophical Transactions of the Royal Society of London, Series B 345, 35–44.
- 1315 Niklas, K.J., Tiffney, B.H., Knoll, A.H., 1980. Apparent changes in the diversity of fossil
- 1316 plants. Evolutionary Biology 12, 1–89.
- 1317 Nowak, H., Schneebeli-Hermann, E., Kustatscher, E., 2019. A non-extinction event for plants
- during the end-Permian mass extinction. Nature Communications, doi.org/10.1038/s41467-018-07945-w.
- 1320 Opluštil, S., Pšenička, J., Libertín, M., Šimůnek, Z., 2007. Vegetation patterns of
- 1321 Westphalian and lower Stephanian mire assemblages preserved in tuff beds of the continental
- basins of Czech Republic. Review of Palaeobotany and Palynology 143, 107–154.
- 1323 Opluštil, S., Pšenička, J., Libertín, M., Bashforth, A., Šimůnek, Z., 2009a. A Middle
- 1324 Pennsylvanian (Bolsovian) peat-forming forest preserved in situ in volcanic ash of the
- 1325 Whetstone Horizon in the Radnice Basin, Czech Republic. Review of Palaeobotany and
- 1326 Palynology 155, 234–274.
- 1327 Opluštil, S., Pšenička, J., Libertín, M., Bek, J., Dašková, J., Šimůnek, Z., Drábková, J.,
- 1328 2009b. Composition and structure of an *in situ* Middle Pennsylvanian peat-forming plant
- 1329 assemblage in volcanic ash, Radnice Basin (Czech Republic). Palaios 24, 726–746.
- 1330 Opluštil, S., Pšenička, J., Bek, J., Wang. J., Feng, Z., Libertín, M., Šimůnek, Z., Bureš, J.,
- 1331 Drábková, J., 2014. T⁰ peat-forming plant assemblage preserved in growth position by

- 1332 volcanic ash-fall: A case study from the Middle Pennsylvanian of the Czech Republic.
- 1333 Bulletin of Geosciences 89, 773–818.
- 1334 Opluštil, S., Šimůnek, Z., Pšenička, J., Bek, J., Libertín, M., 2017. A 25 million year
- 1335 macrofloral record (Carboniferous–Permian) in the Czech part of the Intra-Sudetic Basin;
- 1336 biostratigraphy, plant diversity and vegetation patterns. Review of Palaeobotany and
- 1337 Palynology 244, 241–307.
- 1338 Pan, Y.-L., Hill, S.C., Pinnick, R.G., House, J.M., Flagan, R.C., Chang, R.K., 2011. Dual-
- 1339 excitation-wavelength fluorescence spectra and elastic scattering for differentiation of single
- 1340 airborne pollen and fungal. Atmospheric Environment 14, 1555–1563.
- 1341 Pardoe, H.S., 1996. Micro-scale patterns of modern pollen deposition within three alpine
- 1342 plant communities. New Phytologist 132, 327–341.
- Pardoe, H.S., 2001. The representation of taxa in surface pollen spectra on alpine and subalpine glacier forelands in southern Norway. Review of Palaeobotany and Palynology 117,
 63–78.
- 1346 Pardoe, H.S., 2006. Surface pollen deposition on glacier forelands in southern Norway I:
- 1347 local patterns of representation and source area at Storbreen, Jotunheimen. The Holocene 16,1348 1149–1161.
- 1349 Pardoe, H.S., Giesecke, T., van der Knaap, W.O., Svitavská-Svobodová, H., Kvavadze, E.V.,
- 1350 Panajiotidis, S., Gerasimidis, A., Pidek, I.A., Zimny, M., Świeta-Musznicka, J., Latalowa,
- 1351 M., Noryskiewicz, A.M., Bozilova, E., Tonkov, S., Filipova-Marinova, M.V., van Leeuwen,
- 1352 J.F.N., Kalniņa, L., 2010. Comparing pollen spectra from modified Tauber traps and moss
- 1353 samples: examples from a selection of woodlands across Europe. Vegetation history and
- 1354 Archaeobotany 19, 271–283.

- 1355 Parsons, R.W., Prentice, I.C., 1981. Statistical approaches to R-values and the pollen—
- 1356 vegetation relationship. Review of Palaeobotany and Palynology 32, 127–152.
- 1357 Petit, R.J., Brewer, S., Bordacs, S., et al., 2002. Identification of refugia and post-glacial
- 1358 colonisation routes of European white oaks based on chloroplast DNA and fossil pollen
- evidence. Forest Ecology and Management 156, 49–74.
- 1360 Pfefferkorn, H.W., Gastaldo, R.A., DiMichele, W.A., Phillips, T.L., 2008. Pennsylvanian
- tropical floras from the United States as a record of changing climate, in: Fielding, C.R.,
- 1362 Frank, T.D., Isbell, J.L. (Eds.), Resolving the Late Paleozoic Ice Age in Time and Space.
- 1363 Geological Society of America Special Paper 441, 305–316.
- 1364 Pfefferkorn, H.W., Gastaldo, R.A., DiMichele, W.A., 2017. Impact of an icehouse climate
- interval on tropical vegetation and plant evolution. Stratigraphy 14, 365–376.
- 1366 Phillips, T.L., 1980. Stratigraphic and geographic occurrences of permineralized coal-swamp
- 1367 plants–Upper Carboniferous and Europe, in: Dilcher, D.L., Taylor, T.N. (Eds.),
- 1368 Biostratigraphy of fossil plants. Hutchinson & Ross, Stroundsburg, PA, pp. 25–92.
- 1369 Pidek, I.A., 2004. Preliminary results of pollen trapping in the region of the Roztocze
- 1370 National Park (SE Poland). Annales of the University M. Curie-Sklodowska, Sect. B 49,
- 1371 143–159.
- 1372 Pidek, I.A., Svitavská-Svobodová, H., van der Knaap, W.O., Noryśkiewicz, A.M., Filbrandt-
- 1373 Czaja, A., Noryśkiewicz, B., Latałowa, M., Zimny, M., Święta-Musznicka, J., Bozilova, E.,
- 1374 Tonkov, S., Filipova-Marinova, M., Poska, A., Giesecke, T., Gikov, A., 2010. Variation in
- 1375 annual pollen accumulation rates of *Fagus* along a N–S transect in Europe based on pollen
- 1376 traps. Vegetation history and Archaeobotany 19, 259–270.
- 1377 Piperno, D.R., 2002. Phytoliths, in: Mulholl, S.C, Rapp, G. (eds.) Tracking environmental
- 1378 change using lake sediments. Springer, Dordrecht, pp. 235–251.

- 1379 Piperno, D.R., Pearsall, D.M., 1998. The silica bodies of tropical American grasses:
- 1380 morphology, taxonomy, and implications for grass systematics and fossil phytolith
- 1381 identification. Smithsonian Institution, Washington DC (Contributions to Botany 85).
- 1382 Poinar, G., jr, 2002. Fossil palm flowers in Dominican and Baltic amber. Botanical Journal of
- 1383 the Linnean Society 139, 361–367.
- 1384 Pole, M., 2001. Repeated flood events and fossil forests at Curio Bay (Middle Jurassic), New
- 1385 Zealand. Sedimentary Geology 144, 223–242.
- 1386 Popa, M.E., 1998. The Liassic continental flora of Romania: Systematics, stratigraphy and
- 1387 paleoecology. Acta Botanica Horti Bucurestensis 1997–1998, 177–184.
- 1388 Popa, M.E., 2000. Aspects of Romanian Early Jurassic palaeobotany and palynology. Part III.
- 1389 Phytostratigraphy of the Getic Nappe. Acta Palaeontologica Romaniae 2, 377–386.
- 1390 Popa, M.E., 2011. Field and laboratory techniques in plant compressions: an integrated
- approach. Acta Palaeontologica Romaniae 7, 279–283.
- 1392 Popa, M.E., 2014. Early Jurassic bennettitalean reproductive structures of Romania.
- 1393 Palaeobiodiversity and Palaeoenvironments 94, 327–362.
- 1394 Poska, A., Pidek, I.A., 2010. Pollen dispersal and deposition characteristics of Abies alba,
- 1395 Fagus sylvatica and Pinus sylvestris, Roztocze region (SE Poland). Vegetation History and
- 1396 Archaeobotany 19, 91–101.
- 1397 Potonié, R., 1934. Zur Morphologie der fossilen Pollen und Sporen. Arbeiten aus dem Institut
- 1398 für Paläobotanik und Petrographie Brennsteine 4, 5–24
- 1399 Potonié, R., 1956. Synopsis der Gattungen der Sporae dispersae, Teil 1: Sporites.
- 1400 Geologisches Jahrbuch 23, 1–103.

- 1401 Potonié, R., 1958. Synopsis der Gattungen der Sporae dispersae. II. Teil: Sporites
- 1402 (Nachträge), Saccites, Aletes, Praecolpates, Polyplicates, Monocolpates. Geologisches
- 1403 Jahrbuch 31, 1–114.
- 1404 Potonié, R., 1960. Synopsis der Gattungen der Sporae dispersae: III. Teil: Nachträge Sporites,
- 1405 Fortsetzung *Pollenites* mit Generalregister zu Teil I-II. Geologisches Jahrbuch 39, 1–189.
- 1406 Powell, M.G., Kowalewski, M., 2002. Increase in evenness and sampled alpha diversity
- 1407 through the Phanerozoic: comparison of early Paleozoic and Cenozoic marine fossil
- 1408 assemblages. Geology 30, 31–334.
- 1409 Procter, C.J., 1994. Carboniferous fossil plant assemblages and palaeoecology at the
- 1410 Writhlington Nature Reserve. Proceedings of the Geologists' Association 105, 277–286.
- 1411 Pšenička, J., Opluštil, S., 2013. The epifytic plants in the fossil record and its example from
- 1412 in situ tuff from Pennsylvanian of Radnice Basin (Czech Republic). Bulletin of Geosciences1413 88, 401–416.
- 1414 Rashid, I., Mir, S.H., Zurro, D., Dar, R.A., Reshi, Z.A., 2019. Phytoliths as proxies of the
 1415 past. Earth-Science Reviews 194, 234–250.
- 1416 Raup, D.M., Sepkoski, J.J., 1982. Mass extinctions in the marine fossil record. Science 215,
 1417 1501–1503.
- 1418 Reitalu, T., Bjune, A.E., Blaus, A., Giesecke, T. Helm, T., Matthias, I., Peglar, S.M.,
- 1419 Salonen, J.S., Seppä, H., Väli, V., Birks, H.J.B., 2019. Patterns of modern pollen and plant
- richness across northern Europe. Journal of Ecology 107, 1662–1677.
- 1421 Reitalu, T., Kuneš, P., Giesecke, T., 2014. Closing the gap between plant ecology and
- 1422 Quaternary palaeoecology. Journal of Vegetation Science 25, 1188–1194.

- 1423 Rice, J., Rothwell, G.W., Mapes, G., Mapes, R.H., 1996. Suavitas imbricata gen. et sp. nov.,
- 1424 an anatomically preserved seed analogue of putative lycophyte affinities from Upper
- 1425 Pennsylvanian marine deposits. American Journal of Botany 83, 1083–1090.
- 1426 Roopnarine, P.D., Angielczyk, K.D., Weik, A., Dineen, A., 2018. Ecological persistence,
- 1427 incumbency and reorganization in the Karoo Basin during the Permian-Triassic transition.
- 1428 Earth-Science Reviews 189, 244–263.
- 1429 Rößler, R., Barthel, M., 1998. Rotliegend taphocoenoses preservation favoured by rhyolithic
- 1430 explosive volcanism. Freiberger Forschungshefte, Paläontologie, Stratigraphie, Fazies C
 1431 474(6), 59–101.
- 1432 Rothwell, G.W., Mapes, G., Mapes, R.H., 1996. Anatomically preserved vojnovskyalean
- seed plants in Upper Pennsylvanian (Stephanian) marine shales of North America. Journal ofPaleontology 70, 1067-1079.
- Rull, V., 2011. Neotropical biodiversity: timing and potential drivers. Trends in Ecology and
 Evolution 26, 508–513.
- 1437 Rull, V., 2013. Some problems in the study of the origin of neotropical biodiversity using
- palaeoecological and molecular phylogenetic evidence. Systematics and Biodiversity 11,415–423.
- 1440 Rust, J., Singh, H., Rana, R.S., McCann, T., Singh, L., Anderson, K., Sarkar, N.,
- 1441 Nascimbene, P.C., Stebner, F., Thomas, J.C., et al., 2010. Biogeographic and evolutionary
- 1442 implications of a diverse paleobiotain amber from the early Eocene of India. Proceedings of
- 1443 the National Academy of Sciences 107, 18360–18365.
- 1444 Sadowski E.-M., Schmidt A.R., Seyfullah L.J., Kunzmann L., 2017. Conifers of the 'Baltic
- amber forest' and their palaeoecological significance. Stapfia 106, 1–73.

- 1446 Sadowski, E.M., Seyfullah, L.J., Regalado, L., Skadell, L.E., Gehler, A., Gröhn, C., Hoffeins,
- 1447 C., Hoffeins, H.W., Neumann, C., Schneider, H., Schmidt, A.R., 2019. How diverse were
- 1448 ferns in the Baltic amber forest? Journal of Systematics and Evolution 57, 305–328.
- 1449 Saxena, A., Singh, K.J., Cleal, C.J., Chandra, S., Goswami, S., Shabbar, H., 2018.
- 1450 Development of the Glossopteris flora and its end Permian demise in the Tatapani–Ramkola
- 1451 Coalfield, Son–Mahanadi Basin, India. Geological Journal 54, 2472–2494.
- 1452 Saxena, A., Murthy, S., Singh, K.J., 2020. Floral diversity and environment during the early
- 1453 Permian: a case study from Jarangdih Colliery, East Bokaro Coalfield, Damodar Basin, India.
- 1454 Palaeobiodiversity and Palaeoenvironments 100, 33–50.
- 1455 Scheihing, M.H., Pfefferkorn, H.W., 1984. The taphonomy of land plants in the Orinoco
- 1456 Delta: a model for the incorporation of plant parts in clastic sediments of Late Carboniferous
- 1457 age of Euramerica. Review of Palaeobotany and Palynology 41, 205–240.
- 1458 Schmidt, A.R., Ragazzi, E., Coppellotti, O., Roghi, G., 2006. A microworld in Triassic amber
- Amber as old as the first dinosaurs captured the diversity of microbial life 220 million years
 ago. Nature 444, 835–836.
- 1461 Scott, A.C., 1978. Sedimentological and ecological control of Westphalian B plant
- 1462 assemblages from West Yorkshire. Proceedings of the Yorkshire Geological Society 41, 461–1463 508.
- 1464 Scott, A.C., 1979. The ecology of Coal Measure floras from northern Britain. Proceedings of
- 1465 the Geologists' Association 90, 97–116.
- 1466 Sepkoski, J.J., jr, 1978. A kinetic-model of phanerozoic taxonomic diversity I. Analysis of
- 1467 marine orders. Paleobiology 4, 223–251.
- 1468 Sepkoski, J.J., jr 1979. A kinetic-model of phanerozoic taxonomic diversity II. Early
- 1469 phanerozoic families and multiple equilibria. Paleobiology 5, 222–251.

- 1470 Sepkoski, J.J., jr, 1984. A kinetic-model of phanerozoic taxonomic diversity III. Post-
- 1471 Paleozoic families and mass extinctions. Paleobiology 10, 246–267.
- 1472 Sepkoski, J.J., jr, 1988. Alpha, beta, or gamma: where does all the diversity go? Paleobiology1473 14, 221–234.
- 1474 Seppä, H., Bennett, K.D., 2003. Quaternary pollen analysis: recent progress in palaeoecology
- and palaeoclimatology. Progress in Physical Geography 27, 548–579.
- 1476 Serbet, R., Rothwell, G.W., 1992. Characterizing the most primitive seed ferns. I. A
- 1477 reconstruction of *Elkinsia polymorpha*. International Journal of Plant Sciences 153, 602–621.
- 1478 Servais, T., Cascales-Miñana, B., Cleal, C.J., Gerrienne, P., Harper, D.A., Neumann, M.,
- 1479 2019. Revisiting the Great Ordovician Diversification of land plants: Recent data and
- 1480 perspectives. Palaeogeography, Palaeoclimatology, Palaeoecology 534, 13 pp.
- 1481 Silvestro, D., Cascales-Miñana, B., Bacon, C.D., Antonelli, A., 2015. Revisiting the origin
- 1482 and diversification of vascular plants through a comprehensive Bayesian analysis of the fossil
- 1483 record. New Phytologist 207, 425–436.
- 1484 Šimůnek, Z., 2007. New classification of the genus Cordaites from the Carboniferous and
- 1485 Permian of the Bohemian Massif, based on cuticle micromorphology. Sborník Národního
- 1486 Muzea v Praze, Serie B, Přírodní Vidy 62, 97–210.
- 1487 Sjögren, P., van der Knaap, W.O., van Leeuwen, J.F.N., 2015. Pollen dispersal properties of
- 1488 Poaceae and Cyperaceae: first estimates of their absolute pollen productivities. Review of
- 1489 Palaeobotany and Palynology 216, 123–131.
- 1490 Slater, B.J., McLoughlin, S., Hilton, J., 2015. A high-latitude Gondwanan lagerstätte: the
- 1491 Permian permineralised peat biota of the Prince Charles Mountains, Antarctica. Gondwana
- 1492 Research 27, 1446–1473.

- 1493 Smith, A.H.V., 1962. The palaeoecology of Carboniferous peats based on the microspores
- and petrography of bituminous coals. Proceedings of the Yorkshire Geological Society 33,423–474.
- 1496 Smith, A.H.V., 1968. Seam profiles and seam characters, in: Murchison, D.G., Westoll, T.S.
- 1497 (Eds.), Coal and coalbearing strata. Oliver and Boyd, Edinburgh, pp. 31–40.
- 1498 Smith, A.H.V., Butterworth, M.A., 1967. Miospores in the coal seams of the Carboniferous
- 1499 of Great Britain. Palaeontological Association, (Special Papers in Palaeontology 1).
- 1500 Solórzano Kraemer, M.M., Delclòs, X., Clapham, M., Arillo A., Peris, D., Jäger, P., Stebner,
- 1501 F., Peñalver E., 2018. Arthropods in modern resins reveal if amber accurately recorded forest
- arthropod communities. Proceedings of the National Academy of Sciences 115, 6739–6744.
- 1503 Spicer, R.A., 1980. The importance of depositional sorting to the biostratigraphy of plant
- 1504 megafossils, in: Dilcher, D.L., Taylor, T.N. (Eds.), Biostratigraphy of fossil plants. Dowden,
- 1505 Hutchinson and Ross, Stroudsburg PA, pp. 171–183.
- 1506 Spicer, R.A., 1981. The sorting and deposition of allochthonous plant material in a modern
- 1507 environment at Silwood Lake, Silwood Park, Berkshire, England. Professional Papers of the
- 1508 U.S. Geological Survey 1143, 1–77.
- 1509 Steart, D.C., Boon, P.I., Greenwood, D.R., Diamond, N.T., 2002. Transport of leaf litter in
- 1510 upland streams of Eucalyptus and Nothofagus forests in south-eastern Australia. Archiv für
- 1511 Hydrobiologie 156, 43–61.
- 1512 Steemans P., Lepot, K., Marshall, C.P., le Herissé, A., Javaux, E.J., 2010. FTIR
- 1513 characterisation of the chemical composition of Silurian miospores (cryptospores and trilete
- 1514 spores) from Gotland, Sweden. Review of Palaeobotany and Palynology 162, 577–590.

- 1515 Stein, W.E., Berry, C.M., Hernick, L.V., Mannolini, F., 2012. Surprisingly complex
- 1516 community discovered in the mid-Devonian fossil forest at Gilboa. Nature 483, 78.
- 1517 Sternberg, K.M. von, 1820. Versuch einer geognostisch-botanischen Darstellung der Flora
- 1518 der Vorwelt, Vol. I. 1. F. Fleischer, Leipzig
- 1519 Stolle, E., 2007. Regional Permian palynological correlations: Southeast Turkey Northern
- 1520 Iraq. Comunicações Geológicas 94, 125–143.
- 1521 Stolle, E., 2010. Recognition of southern Gondwanan palynomorphs at Gondwana's northern
- 1522 margin and biostratigraphic correlation of Permian strata from SE Turkey and Australia, in:
- 1523 Shen, S.-Z., Henderson, C.M., Somerville, I.D. (Eds.), Lopingian (Late Permian) stratigraphy
- 1524 of the world, major events and environmental change. Geological Journal 45, 336–349.
- 1525 Stolle, E., 2011. Pollen-dominated European palynological assemblages from the Permian of
- 1526 NW Turkey (Asia Minor) palaeogeographical context and microfloral affinities. Geological
- 1527 Quarterly 55, 181–186.
- 1528 Stolle, E., 2012. Co-occurrence of Sinuspores sinuatus (Artüz) Ravn, 1986 with established
- 1529 palynological markers indicating younger strata: AK-1X well section (Pennsylvanian,
- 1530 Zonguldak Basin, NW Turkey) and the correlation to the stratigraphic system. Geologia
- 1531 Croatica 65, 375–385.
- 1532 Stolle, E., 2016. Çakraz Formation, Çamdağ area, NW Turkey: early/mid-Permian age,
- 1533 Rotliegend (Germany) and Southern Alps (Italy) equivalent—a stratigraphic re-assessment
- 1534 via palynological long-distance correlation. Geological Journal 51, 223–235.
- 1535 Stolle, E., Yalçın, M.N., Kozlu, H. 2012. Palynofacies and bulk organic geochemistry of
- 1536 Permian clastics in the eastern Taurids: Implications for hydrocarbon potential, in: Yalçın,
- 1537 M.N., Çorbacıoğlu, H., Aksu, Ö., Bozdoğan, N. (Eds.), Paleozoic of northern Gondwana and

- 1538 its petroleum potential. Turkish Association of Petroleum Geologists, Ankara (Special
- 1539 Publication 6), pp. 119–122.
- 1540 Strömberg, C.A., 2004. Using phytolith assemblages to reconstruct the origin and spread of
- 1541 grass-dominated habitats in the great plains of North America during the late Eocene to early
- 1542 Miocene. Palaeogeography, Palaeoclimatology, Palaeoecology 207, 239–275.
- 1543 Strömberg, C.A., 2011. Evolution of grasses and grassland ecosystems. Annual Review of
- 1544 Earth and Planetary Sciences 39, 517–544.
- 1545 Strömberg, C.A., Dunn, R.E., Crifò, C., Harris, E.B., 2018. Phytoliths in paleoecology:
- 1546 analytical considerations, current use, and future directions, in: Croft, D.A., Su, D., Simpson,
- 1547 S.W. (Eds.), Methods in paleoecology. Springer, Cham, pp. 235–287.
- Strullu-Derrien, C., Kenrick, P., Knoll, A.H., 2019. The Rhynie chert. Current Biology 29,
 1218–1223.
- 1550 Sugita, S.A., 1993. Model of pollen source area for an entire lake surface. Quaternary
- 1551 Research 39, 239–244.
- 1552 Swingland, I.R., 2001. Biodiversity, definition of, in: Levin, S.A. (Ed.), Encyclopedia of
- 1553 biodiversity. Volume 1. Academic Press, San Diego, pp. 377–391.
- 1554 Taylor, T.N., Taylor, E.L., Kings, M., 2009. Paleobiology. The biology and evolution of
- 1555 fossil plants. [Second Edition]. Academic Press, Burlington.
- 1556 Thomas, B.A., 1987. The use of in-situ spores for defining species of dispersed spores.
- 1557 Review of Palaeobotany and Palynology 51, 227–233.
- 1558 Thomas, B.A., 2005. A reinvestigation of Selaginella species from the Asturian (Westphalian
- 1559 D) of the Zwickau coalfield, Germany and their assignment to the new sub genus
- 1560 Hexaphyllum. Zeitschrift der Deutschen Gesellschaft für Geowissenschaften 156, 403–418.

- 1562 Pennsylvanian (Carboniferous) calamitaleans based upon new studies of Calamites
- 1563 Sternberg, 1820 in the Duckmantian at Brymbo, North Wales, UK. Palaeontology 57, 21–36.
- 1564 Thomas, B.A., Cleal, C.J., 2015. Cyclones and the formation of plant beds in late
- 1565 Carboniferous tropical swamps. Palaeobiodiversity and Palaeoenvironments 95, 531–536.
- 1566 Thomas, B.A., Seyfullah, L.J., 2015. Stigmaria Brongniart: a new specimen from
- 1567 Duckmantian (Lower Pennsylvanian) Brymbo (Wrexham, North Wales) together with a
- review of known casts and how they were preserved. Geological Magazine 152, 858–870.
- 1569 Thomas, B.A., Cleal, C.J., 2020. The nomenclature of fossil-taxa representing different
- 1570 preservational states: *Lepidodendron* as a case-study. Taxon, doi.org/10.1002/tax.12291.
- 1571 Thomas, B.A., Dimitrova, T.K., 2017. Ecological changes in Pennsylvanian (Asturian and
- 1572 early Cantabrian) coal floras inferred from lycophyte microspore abundances. Earth Science
- 1573 Reviews 171, 646–662.
- 1574 Thomas, B.A., Appleton, P., Cleal, C.J., Seyfullah, L.J., 2020. The distribution of plant
- 1575 fossils and their palaeoecology in Duckmantian (Bashkirian, Lower Pennsylvanian) strata at
- 1576 Brymbo, North Wales, UK. Geological Journal 55, 3179–3207.
- 1577 Tinner, W., Lotter, A.F., 2006. Holocene expansions of *Fagus sylvatica* and *Abies alba* in
- 1578 Central Europe: where are we after eight decades of debate? Quaternary Science Reviews 25,1579 526–549.
- 1580 Tomescu, A.M., Bomfleur, B., Bippus, A.C., Savoretti, A., 2018. Why are bryophytes so rare
- 1581 in the fossil record? A spotlight on taphonomy and fossil preservation, in: Krings, M.,
- 1582 Harper, C.J., Cuneo, N.R., Rothwell, G.W. (Eds.), Transformative paleobotany. Elsevier,
- 1583 Amsterdam, pp. 375–416.

- 1584 Traverse, A., 1988. Plant evolution dances to a different beat. Plant and animal evolutionary
- 1585 mechanisms compared. Historical Biology 1, 227–301.
- 1586 Traverse, A., 2007. Paleopalynology (2nd edition). Springer, Dordrecht.
- Tuomisto, H., 2012. An updated consumer's guide to evenness and related indices. Oikos
 121, 1203–1218.
- 1589 Uhl, D., 2006. Some considerations on the taphonomy of conifer remains from the Late
- 1590 Permian of Europe. Neues Jahrbuch für Geologie & Paläontologie, Monatshefte 08/2006,
 1591 483–502.
- 1592 Uhl, D. 2015. Preliminary note on fossil flowers and inflorescences from the Late Oligocene
- 1593 of Enspel (Westerwald, W-Germany). Palaeobiodiversity and Palaeoenvironments 95, 47–53.
- 1594 Uhl, D., Cleal, C.J., 2010. Late Carboniferous vegetation change in lowland and intramontane
- 1595 basins. International Journal of Coal Geology 83, 318–328.
- 1596 Uhl, D., Krüger, P.S., Wuttke, M., 2018. Epidermal anatomy of *Glyptostrobus europaeus*
- 1597 (Brongn.) Unger from the late Oligocene of the Westerwald (Rhineland-Palatinate, W-
- 1598 Germany). Fossil Imprint 74, 334–340.
- 1599 Urban, M.A., Nelson, D.M., Jiménez-Moreno, G., Châteauneuf, J.J., Pearson, A., Hu, F.S.,
- 1600 2010. Isotopic evidence of C₄ grasses in southwestern Europe during the Early Oligocene–
- 1601 Middle Miocene. Geology 38, 1091–1094.
- 1602 Vajda, V., Bercovici, A., 2014. The global vegetation pattern across the Cretaceous-
- 1603 Paleogene mass extinction interval: A template for other extinction events. Global and
- 1604 Planetary Change 122, 29–49.
- 1605 van der Knaap, W.O., van Leeuwen, J.F., Finsinger, W., Gobet, E., Pini, R., Schweizer, A.,
- 1606 Valsecchi, V., Ammann, B., 2005. Migration and population expansion of Abies, Fagus,

- 1607 *Picea*, and *Quercus* since 15000 years in and across the Alps, based on pollen-percentage
- 1608 threshold values. Quaternary Science Reviews 24, 645–680.
- 1609 Vellend, M., Cornwell, W.K., Magnuson-Ford, K., Mooers, A.Ø., 2011. Measuring
- 1610 phylogenetic biodiversity, in: McGill, B.J., Magurran, A. (eds.), Biological diversity:
- 1611 frontiers in measurement and assessment. Oxford University Press, Oxford, pp. 194–207.
- 1612 Vellend, M., Baeten, L., Becker-Scarpitta, A., Boucher-Lalonde, V., McCune, J.L., Messier,
- 1613 J., Myers-Smith, I.H., Sax, D.F., 2017. Plant biodiversity change across scales during the
- 1614 Anthropocene. Annual Review of Plant Biology 68, 563–586.
- 1615 Wagner, R.H., 1989. A late Stephanian forest swamp with Sporangiostrobus fossilized by
- 1616 volcanic ash fall in the Puertollano Basin, central Spain. International Journal of Coal
- 1617 Geology 12, 523–552.
- Wagner, R.H., Diez, J.B., 2007. Verdeña (Spain): Life and death of a Carboniferous forest
 community. Compte Rendu Palevol 6, 495–504
- 1620 Wang Deming, Qin Min, Liu Le, Liu Lu, Zhou Yi, Zhang Yingying, Huang Pu, Xue
- 1621 Jinzhuang, Zhang Shuhui, Meng Meicen, 2019. The most extensive Devonian fossil forest
- with small lycopsid trees bearing the earliest stigmarian roots. Current Biology 29, 2604–
 2615.
- 1624 Wang Jun, Pfefferkorn, H.W., Zhang Yi, Feng Zhou, 2012. Permian vegetational Pompeii
- 1625 from Inner Mongolia and its implications for landscape paleoecology and paleobiogeography
- 1626 of Cathaysia. Proceedings of the National Academy of Sciences 109, 4927–4932.
- Wellman, C.H., 2010. The invasion of the land by plants: when and where? New Phytologist188, 306–309.

- 1629 Weng, C., Hooghiemstra, H., Duivenvoorden, J.F., 2006. Challenges in estimating past plant
- 1630 diversity from fossil pollen data: statistical assessment, problems, and possible
- 1631 solutions. Diversity and distributions 12, 310–318.
- 1632 Whittaker, R.H., 1960. Vegetation of the Siskiyou Mountains, Oregon and California.
- 1633 Ecological Monographs 30, 279–338.
- Whittaker, R.H., 1977. Evolution of species diversity in land communities. Evolutionary
 Biology 10, 1–67.
- 1636 Whittaker, R.J., Willis, K.J., Field, R., 2001. Scale and species richness: towards a general,
- 1637 hierarchical theory of species diversity. Journal of Biogeography 28, 453–470.
- 1638 Willard, D.A., 1993. Vegetational patterns in the Springfield coal (Middle Pennsylvanian,
- 1639 Illinois Basin): comparison of miospore and coal-ball records. Geological Society of
- 1640 America, Special Paper 286, 139–152.
- 1641 Willard, D.A., Donders, T.H., Reichgelt, T., Greenwood, D.R., Sangiorgi, F., Peterse, F.,
- 1642 Nierop, K.G., Frieling, J., Schouten, S., Sluijs, A., 2019. Arctic vegetation, temperature, and
- 1643 hydrology during Early Eocene transient global warming events. Global and Planetary
- 1644 Change 178, 139–152.
- 1645 Willard, D.A., Phillips, T.L., Lesnikowska, A.D., DiMichele, W.A., 2007. Paleoecology of
- 1646 the Late Pennsylvanian-age Calhoun coal bed and implications for long-term dynamics of
- 1647 wetland ecosystems. International Journal of Coal Geology 69, 21–54.
- 1648 Williams, J.W., Grimm, E.C., Blois, J.L., Charles, D.F., Davis, E.B., Goring, S.J., Graham,
- 1649 R.W., Smith, A.J., Anderson, M., Arroyo-Cabrales, J., Ashworth, A.C., 2018. The Neotoma
- 1650 Paleoecology Database, a multiproxy, international, community-curated data resource.
- 1651 Quaternary Research 89, 156–177.

- 1652 Willis, K.J., Bennett, K.D., Bhagwat, S.A., Birks, H.J.B., 2010. Perspective 4°C and beyond:
- 1653 What did this mean for biodiversity in the past? Systematics and Biodiversity 8, 3–9.
- 1654 Willis, K.J., Bhagwat, S.L., 2009. Biodiversity and climate change. Science 326, 806–807.
- Willis, K.J., McElwain, J., 2013. The evolution of plants (2nd edition). Oxford UniversityPress, Oxford.
- 1657 Willis, K.J., Niklas, K.J., 2004. The role of Quaternary environmental change in plant
- 1658 macroevolution: the exception or the rule? Philosophical Transactions of the Royal Society B
- 1659 Biological Series 359, 159–72.
- 1660 Willis, K.J., Rudner, E., Sümegi, P., 2000. The full-glacial forests of central and southeastern
- 1661 Europe. Quaternary Research 53, 203–213.
- 1662 Wing, S.L., DiMichele, W.A., 1992. Ecological characterization of fossil plants, in:
- 1663 Behrensmeyer, A.K., Damuth, J.D., DiMichele, W.A., Potts, R., Sues, H.-D., Wing, S.L.
- 1664 (Eds.) Terrestrial ecosystems through time. University of Chicago Press, Chicago, pp. 139–
 1665 180.
- 1666 Wing, S.L., DiMichele, W.A., 1995. Conflict between local and global changes in plant
- 1667 diversity through geological time. Palaios 10, 551–564.
- 1668 Wing, S.L., Harrington, G.J., Bowen, G.J., Koch, P.L., 2003. Floral change during the initial
- 1669 Eocene thermal maximum in the Powder River Basin, Wyoming, in: Wing, S.L., Gingerich,
- 1670 P.D., Schmitz, B., Thomas, E. (Eds.), Causes and Consequences of Globally Warm Climates
- 1671 in the Early Paleogene. Geological Society of America, Boulder, Colorado, (Special Paper
- 1672 369), pp. 425-440.
- 1673 Wolfe, J.A., 1993. A method of obtaining climatic parameters from leaf assemblages. U.S.
- 1674 Geological Survey Bulletin 2040, 1–71.
- 1675 Xing, Y., Gandolfo, M.A., Onstein, R.E., Cantrill, D.J., Jacobs, B.F., Jordan, G.J., Lee, D.E.,
- 1676 Popova, S., Srivastava, R., Su, T., Vikulin, S.V., 2016. Testing the biases in the rich
- 1677 Cenozoic angiosperm macrofossil record. International Journal of Plant Sciences 177, 371–1678 388.
- 1679 Xiong C., Wang D., Wang Q., Benton, M.J., Xue J., Meng M., Zhao Q., Zhang, J., 2013.
- 1680 Diversity dynamics of Silurian–Early Carboniferous land plants in South China. PLoS One

1681 8(9), p.e75706.

- 1682 Yu, J., Broutin, J., Chen, Z.Q., Shi, X., Li, H., Chu, D., Huang, Q., 2015. Vegetation
- 1683 changeover across the Permian–Triassic Boundary in Southwest China: extinction, survival,
- 1684 recovery and palaeoclimate: a critical review. Earth-Science Reviews 149, 203–224.
- 1685 Zavialova, N., Kustatscher, E., van Konijnenburg-van Cittert, J.H.A., 2010. Spore
- 1686 ultrastructure of Selaginellites leonardii and diversity of Selaginellalean spores. Geo.Alp 7,

1687 1–17.

1688

1689 Figure captions

- 1690 Fig. 1. Variation in productivity of different organs of a plant as illustrated by the pollen,
- 1691 flowers, leaves, shoots, stem and roots of a hypothetical modern-day angiosperm tree.
- 1692 Redrawn and adapted from Hughes (1976, fig. 3.6) and Cleal and Thomas (2019, fig. 10.4).
- 1693 Fig. 2. Three types of diversity that can be recognised in the plant fossil record, using the
- 1694 Middle Pennsylvanian (c. 310 Ma) swamp vegetation of Variscan Euramerica, based on Cleal1695 et al. (2012).
- 1696 Fig. 3. Morphological similarity of late Permian gigantopterid leaves (A, C, Gigantonoclea
- 1697 hallei (Halle) Wang) and modern angiosperm leaves (B, Castanea sativa Miller; D, Quercus
- 1698 robur L.). All scale bars = 10 mm. A, C, Naturhistoriska Riksmuseet (NRM), Stockholm (A,
- 1699 NRM S128498; C, NRM S128494), B, D, Royal Botanic Gardens Edinburgh herbarium
- 1700 (RBGE). Adapted from Glasspool et al. (2004).
- 1701 Fig. 4. Partial reconstruction of the Late Devonian seed plant *Elkinsia* based on associated
- 1702 fronds, ovulate structures and anatomically preserved stems. Drawn from by Annette
- 1703 Townsend (based on Serbet and Rothwell 1992).
- 1704 Fig. 5. Examples of the differences in the fossil-genera represented by Carboniferous
- arborescent lycopsids and sphenopsids. Adapted from Cleal and Thomas (2019).
- 1706 Fig. 6. Spores of fern Weischelia reticulata (Stokes and Webb) Fontaine showing different
- 1707 maturation stages; Escucha Formation (Albian), Escucha, northern Teruel Province, Spain. A,
- 1708 General view of a soral cluster up to 2 mm in diameter showing tightly packed peltate
- 1709 indusia. B, Tightly-packed spores grouped inside a receptaculum. C, Inaperturate, discoidal
- 1710 spores with smooth exine and lacking trilete mark. D, Packed spores showing different
- 1711 ontogenetic stages. D, E, Fully-developed tetragonal spores with well-rounded corners and

1712 clear trilete scar. Original unpublished material from the study in Diez et al (2005) with1713 permission of the authors.

1714 Fig. 7. T⁰ fossil or submerged forests of arborescent lycopsids in the Carboniferous of the

1715 UK. A, Fossil trees rooted in a coal seam being exposed at Brymbo, north Wales (Appleton et

1716 al. 2010). B, Excavated trees in the Victoria Park, Glasgow (Thomas and Seyfullah 2015).

1717 Fig. 8. Palaeozoic wetland vegetation preserved in the lower unit of the early Moscovian

1718 Whetstone Horizon (Bělka tuff), Ovčin, Central Bohemia, Czech Republic. A, Remains of

1719 cordaites and arborescent lycopsids plotted out on an exposed area of the tuff divided into 1

1720 m² quadrats; the small number against each specimen represents the height above the base of

1721 the tuff that the fossil occurred. B, Reconstruction of forest based on the type of plots shown

1722 in Fig. 7A. From Opluštil et al. (2014).

1723 Fig. 9. Comparison of palynological and macrofloral spectra obtained from roof-shales

1724 overlying four Moscovian-age coal seams in South Wales, UK, between the Daren Ddu Seam

at the base and the Llantwit No. 1 Seam at the top. Redrawn from Dimitrova et al. (2005, fig.4).

1727 Fig. 10. Range of basinal and extra-basinal vegetation represented in Moscovian (late

1728 Carboniferous) palynospectra from the Sydney Coalfield, Cape Breton, Canada (Dimitrova et1729 al. 2011).

1730 Fig. 11. The Evolutionary Floras model of vegetation evolution based on a factor analysis of

1731 global plant-family distribution through the Phanerozoic (Cleal and Cascales-Miñana 2014).

1732 Fig. 12. Factors that affect how we interpret past vegetation diversity from the macrofloral

and palynological records, demonstrating the importance of integrating the two sets of data.

Table 1. Fossil-genera assigned to different parts of six representative plants from the main groups in the late Carboniferous tropical coal swamps. The fossil-genera selected to represent each plant group in taxonomic diversity studies (e.g. Cleal et al. 2012) designated by an asterisk (*). This does not include the pollen/spores produced by these plants.

		Lycospids	Calamites	Sphenophylls	Marattialeans	Medullosaleans	Cordaites
Stems		Lepidophloios*	Calamites	. Sphenophyllum*	Caulopteris	-	Artisia
Foliage		Cyperites	Annularia*			Alethopteris*	Cordaites*
Reproductive structures	Female	Lepidocarpon	Calamostachys	Bowmanites	Cyathocarpus*	Trigonocarpus	Cardiocarpus
		Lepidostrobophyllum					
	Male	Lepidostrobus				Whittleseya	Cordaitanthus
Rooting structures		Stigmaria	Pinnularia	-	-	-	-
Total fossil-genera		6	4	2	2	3	4