

Packing k -partite k -uniform hypergraphs

Mycroft, Richard

DOI:

[10.1016/j.jcta.2015.09.007](https://doi.org/10.1016/j.jcta.2015.09.007)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Mycroft, R 2016, 'Packing k -partite k -uniform hypergraphs', *Journal of Combinatorial Theory, Series A*, vol. 138, pp. 60-132. <https://doi.org/10.1016/j.jcta.2015.09.007>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

PACKING k -PARTITE k -UNIFORM HYPERGRAPHS

RICHARD MYCROFT

ABSTRACT. Let G and H be k -graphs (k -uniform hypergraphs); then a perfect H -packing in G is a collection of vertex-disjoint copies of H in G which together cover every vertex of G . For any fixed H let $\delta(H, n)$ be the minimum δ such that any k -graph G on n vertices with minimum codegree $\delta(G) \geq \delta$ contains a perfect H -packing. The problem of determining $\delta(H, n)$ has been widely studied for graphs (*i.e.* 2-graphs), but little is known for $k \geq 3$. Here we determine the asymptotic value of $\delta(H, n)$ for all complete k -partite k -graphs H , as well as a wide class of other k -partite k -graphs. In particular, these results provide an asymptotic solution to a question of Rödl and Ruciński on the value of $\delta(H, n)$ when H is a loose cycle. We also determine asymptotically the codegree threshold needed to guarantee an H -packing covering all but a constant number of vertices of G for any complete k -partite k -graph H .

1. INTRODUCTION

1.1. Basic notions. A k -uniform hypergraph, or k -graph H consists of a vertex set $V(H)$ and an edge set $E(H)$, where every $e \in E(H)$ is a set of precisely k vertices of H . So a 2-graph is a simple graph. We often identify H with its edge set, for example writing $e \in H$ to mean that e is an edge of H , or $|H|$ to denote the number of edges of H .

If G and H are k -graphs, then an H -packing in G (also known as an H -tiling or H -matching) is a collection of vertex-disjoint copies of H in G . This is a generalisation of a matching in G , which is the case of an H -packing when H is the k -graph with k vertices and one edge. We say that a matching or H -packing in G is *perfect* if it covers every vertex of G . Clearly G can only contain a perfect H -packing if $|V(H)|$ divides $|V(G)|$.

We focus mainly on the case when H is a fixed k -graph and $|V(G)|$ is much larger than $|V(H)|$. Our general question is then: what minimum degree conditions on G are sufficient to guarantee that G contains a perfect H -packing? There are various notions of minimum degree for k -graphs, but we shall consider here only one, namely the *codegree*. Let G be a k -graph on n vertices. For any set $S \subseteq V(G)$, the *degree* $\deg_G(S)$ of S is the number of edges of G which contain S as a subset. The *minimum codegree* $\delta(G)$ of G is then the minimum of $\deg_G(S)$ taken over all sets S of $k - 1$ vertices of G . Note that this coincides with the standard notion of degree for graphs.

For any fixed k -graph H and any integer n we define $\delta(H, n)$ to be the smallest integer δ such that any k -graph G on n vertices with minimum codegree $\delta(G) \geq \delta$ contains a perfect H -packing. As noted earlier, this is only defined for those n which are divisible by $|V(H)|$; we shall only consider these values of n . A major problem in extremal graph theory is to determine the behaviour of $\delta(H, n)$ for large n .

1.2. Perfect packings in graphs. In the case when H is a graph, this question has been widely studied, and the value of $\delta(H, n)$ has been determined up to an additive constant in all cases. Indeed, the celebrated Hajnal-Szemerédi theorem [10] determined that $\delta(K_r, n) = (r - 1)n/r$, and Komlós, Sárközy and Szemerédi [20] showed that for any graph H there exists a constant C such that $\delta(H, n) \leq (1 - 1/\chi(H))n + C$. This confirmed a conjecture of Alon and Yuster [3], who had previously established the weaker result with $o(n)$ in place of C , and who

observed that the constant C cannot be removed completely. Finally Kühn and Osthus [22] determined the value of $\delta(H, n)$ up to an additive constant for any graph H using the critical chromatic number $\chi_{\text{cr}}(H)$ first introduced by Komlós [19].

Since there are multiple similarities between the results of Kühn and Osthus for graphs [22] and the results of this paper for k -graphs, we shall state their results in some detail. Let H be a graph on m vertices, and let $\chi(H)$ denote the chromatic number of H , which we assume here to be greater than two (the behaviour in the bipartite case is somewhat different, but is less closely related to the k -graph results considered in this paper). So assume that $\chi(H) = r \geq 3$, and define $\mathcal{D}(H) := \bigcup_c \{|X_i^c| - |X_j^c| : i, j \in [r]\}$, where the union is taken over all proper r -colourings c of H , and X_1^c, \dots, X_r^c denote the colour classes of c . We then define $\gcd(H)$ to be the greatest common factor of $\mathcal{D}(H)$, unless $\mathcal{D}(H) = \{0\}$, in which case $\gcd(H)$ is undefined. Also, we define $\sigma(H) := \min_{c,j} |X_j^c|/m$, so $\sigma(H)$ is the smallest possible size of a colour class of a proper r -colouring of H , expressed as a proportion of the number of vertices of H . Kühn and Osthus [22] demonstrated that there exists a constant C such that

$$\left(1 - \frac{1}{\chi^*(H)}\right)n \leq \delta(H, n) \leq \left(1 - \frac{1}{\chi^*(H)}\right)n + C,$$

where

$$\chi^*(H) = \begin{cases} \chi_{\text{cr}}(H) := \frac{\chi(H)-1}{1-\sigma(H)} & \text{if } \gcd(H) = 1, \\ \chi(H) & \text{otherwise.} \end{cases}$$

1.3. Perfect packings in hypergraphs: known results. For k -graphs H with $k \geq 3$ much less is known. Indeed, the only cases for which $\delta(H, n)$ is known even asymptotically are the cases when H is a 3-graph on 4 vertices and the case of a perfect matching (*i.e.* when $H = K_k^k$ consists of k vertices and one edge). The first bounds for the latter case were given by Daykin and Häggkvist [7], and later Rödl, Ruciński and Szemerédi [31] showed that $n/2 - k \leq \delta(K_k^k, n) \leq n/2$ for all sufficiently large n (indeed, they actually determined the exact value of $\delta(K_k^k, n)$ for large values of n). Beyond this, the value of $\delta(H, n)$ is known for three other 3-graphs, all on four vertices. Let $K_4^3 - 2e$, $K_4^3 - e$ and K_4^3 denote the 3-graphs on 4 vertices with 2, 3 and 4 edges respectively. The value of $\delta(K_4^3 - 2e, n)$ was found to be $n/4 + o(n)$ by Kühn and Osthus [21]; recently Czygrinow, DeBiasio and Nagle [4] found the exact value for large n to be either $n/4$ or $n/4 + 1$ according to the parity of $n/4$. Lo and Markström [25, 27] showed that $\delta(K_4^3 - e, n) = n/2 + o(n)$ and that $\delta(K_4^3, n) = 3n/4 + o(n)$. Simultaneously with the latter, Keevash and Mycroft [16] showed that the exact value of $\delta(K_4^3, n)$ for large n is $3n/4 - 1$ or $3n/4 - 2$, again according to the parity of $n/4$; these results confirmed a conjecture of Pikhurko [29], who had previously shown that $\delta(K_4^3, n) \leq 0.8603n$, and who gave the construction which establishes the lower bound on $\delta(K_4^3, n)$. The exact value of $\delta(K_4^3 - e, n)$ for large n remains an open problem. The cases listed above comprise all the k -graphs H with no isolated vertices for which the value of $\delta(H, n)$ was previously known even asymptotically (if H contains an isolated vertex then the behaviour is somewhat different, as we can restate the problem as asking for non-perfect packing of a smaller k -graph).

Other conditions, such as different notions of degree, which guarantee a perfect H -packing in a large k -graph G have also been considered; see the survey by Rödl and Ruciński [30] for a full account of these. In particular, in recent years there has been much study of the case of a perfect matching, see e.g. [1, 2, 6, 11, 14, 16, 17, 18, 23, 26, 28, 29, 35, 36]. For perfect H -packings other than a perfect matching, results are much more sparse. Lo and Markström [27] found the asymptotic values of $\delta_1(K_3^3(m), n)$ and $\delta_1(K_4^4(m), n)$, where $\delta_1(H, n)$ denotes the smallest integer δ such that any k -graph G on n vertices with $\deg_G(\{x\}) \geq \delta$ for any $x \in V(G)$ contains a perfect H -packing, and $K_r^r(m)$ denotes the complete r -partite r -graph (defined

below) whose vertex classes each have size m . More recently, Han and Zhao [12] gave the exact value of $\delta_1(K_4^3 - 2e, n)$ for large n , whilst Lenz and Mubayi [24] proved that for any linear k -graph F (meaning that any two edges of F intersect in at most one vertex), any sufficiently large ‘quasirandom’ k -graph with linear density contains a perfect F -packing. However, in general our knowledge of conditions which guarantee a perfect H -packing in a k -graph G remains very limited.

1.4. Perfect packings in hypergraphs: new results. In this paper we determine the asymptotic value of $\delta(K, n)$ for all complete k -partite k -graphs, as well as a large class of non-complete k -partite k -graphs K . Let K be a k -graph on vertex set U with at least one edge (if K has no edges then trivially $\delta(K, n) = 0$). Then a k -partite realisation of K is a partition of U into vertex classes U_1, \dots, U_k so that for any $e \in K$ and $1 \leq j \leq k$ we have $|e \cap U_j| = 1$. Equivalently, we colour all vertices of K with k colours so that no edge contains two vertices of the same colour; the vertex classes are then the colour classes. Note in particular that we must have $|U_j| \geq 1$ for every $1 \leq j \leq k$. We say that K is k -partite if it admits a k -partite realisation. The complete k -partite k -graph with vertex classes U_1, \dots, U_k is the k -graph on $U = U_1 \cup \dots \cup U_k$ in which every set $e \subseteq U$ with $|e \cap U_j| = 1$ for each $1 \leq j \leq k$ is an edge. Observe that a complete k -partite k -graph has only one k -partite realisation up to permutations of the vertex classes U_1, \dots, U_k .

Our first theorem states that for any k -partite k -graph K we have $\delta(K, n) \leq n/2 + o(n)$.

Theorem 1.1. *Let K be a k -partite k -graph on b vertices. Then for any $\alpha > 0$ there exists n_0 such that if G is a k -graph on $n \geq n_0$ vertices for which b divides n and $\delta(G) \geq n/2 + \alpha n$, then G contains a perfect K -packing.*

Theorem 1.1 could also be proved by the so-called ‘absorbing method’ by using similar arguments and results to those of Lo and Markström [27]. However, our methods also give stronger bounds for many k -partite k -graphs K , for this we make the following definitions. Let K be a k -partite k -graph on vertex set U . Then we define

$$\mathcal{S}(K) := \bigcup_{\chi} \{|U_1|, \dots, |U_k|\} \text{ and } \mathcal{D}(K) := \bigcup_{\chi} \{||U_i| - |U_j|| : i, j \in [k]\},$$

where in each case the union is taken over all k -partite realisations χ of K into vertex classes U_1, \dots, U_k of K . The *greatest common divisor* of K , denoted $\gcd(K)$, is then defined to be the greatest common divisor of the set $\mathcal{D}(K)$ (if $\mathcal{D}(K) = \{0\}$ then $\gcd(K)$ is undefined). So for any given k -partite realisation of K , the difference in size of any two vertex classes of this realisation must be divisible by $\gcd(K)$. However, it is not true that a k -partite k -graph K must have some k -partite realisation in which the greatest common factor of the differences of vertex class sizes is $\gcd(K)$. To see this, take disjoint sets A, B, C, D and E of size one, one, two, two and six respectively. Form a 3-graph K on $A \cup B \cup C \cup D \cup E$ whose edges are any triple $\{x, y, z\}$ with $x \in A, y \in C$ and $z \in E$ or with $x \in B, y \in D$ and $z \in E$. Then, up to permutation of the vertex classes, K has two distinct 3-partite realisations, one with vertex classes $A \cup B, C \cup D$ and E of sizes two, four and six (so the highest common factor of the differences of class sizes is two), and the other with vertex classes $A \cup D, B \cup C$ and E of sizes three, three and six (whose differences have highest common factor three). So $\gcd(K) = 1$ in this case, but the differences between sizes of vertex classes of any single k -partite realisation of K have a larger common factor.

We also define the *smallest class ratio* of K , denoted $\sigma(K)$, by

$$\sigma(K) := \frac{\min_{S \in \mathcal{S}(K)} S}{|V(K)|}.$$

Property of K	Type	Upper bound on $\delta(K, n)$
$\mathcal{S}(K) = \{1\}$	0	$n/2 + o(n)$
$\gcd(\mathcal{S}(K)) > 1$	0	$n/2 + o(n)$
$\gcd(K) = 1$	1	$\sigma(K)n + o(n)$
$\gcd(\mathcal{S}(K)) = 1, \gcd(K) = d > 1$	d	$\max\left(\sigma(K)n, \frac{n}{p(d)}\right) + o(n)$

TABLE 1. A summary of the upper bounds on $\delta(K, n)$ provided by Theorems 1.1, 1.2 and 1.3, according to the *type* of K as defined in Section 2. In the final row, $p(d)$ denotes the smallest prime factor of d .

So each vertex class of any k -partite realisation of K has size at least $\sigma(K)|V(K)|$. The parameter $\sigma(K)$ therefore provides a measure of how ‘lopsided’ K can be. Note in particular that $\sigma(K) \leq 1/k$, with equality if and only if we have $|U_1| = |U_2| = \dots = |U_k|$ for any k -partite realisation of K with vertex classes U_1, \dots, U_k .

Observe that the definitions of the parameters $\gcd(K)$ and $\sigma(K)$ of a k -partite k -graph K bear a strong resemblance to those of the parameters $\gcd(H)$ and $\sigma(H)$ defined earlier for a graph H with chromatic number $r = k$. We saw that Kühn and Osthus showed that these parameters determine $\delta(H, n)$ for any such H ; similarly $\gcd(K)$ and $\sigma(K)$ play an extensive role in determining $\delta(K, n)$ for a k -partite k -graph K . Indeed, our next theorem strengthens Theorem 1.1 for k -partite k -graphs K with $\gcd(K) = 1$, by stating that $\delta(K, n) \leq \sigma(K)n + o(n)$ for such graphs.

Theorem 1.2. *Let K be a k -partite k -graph on b vertices, and suppose that $\gcd(K) = 1$. Then for any $\alpha > 0$ there exists n_0 such that if G is a k -graph on $n \geq n_0$ vertices for which b divides n and $\delta(G) \geq \sigma(K)n + \alpha n$, then G contains a perfect K -packing.*

Our third theorem improves on Theorem 1.1 for k -partite k -graphs K with $\gcd(\mathcal{S}(K)) = 1$ such that $\gcd(K) > 1$ is odd.

Theorem 1.3. *Let K be a k -partite k -graph on b vertices such that $\gcd(\mathcal{S}(K)) = 1$ and $\gcd(K) > 1$, and let p be the smallest prime factor of $\gcd(K)$. Then for any $\alpha > 0$ there exists n_0 such that if G is a k -graph on $n \geq n_0$ vertices for which b divides n and*

$$\delta(G) \geq \max\left\{\sigma(K)n + \alpha n, \frac{n}{p} + \alpha n\right\},$$

then G contains a perfect K -packing.

For convenience, the upper bounds on $\delta(K, n)$ provided by Theorems 1.1, 1.2 and 1.3 are summarised in Table 1. In Section 2 we shall see that these upper bounds are best possible up to the error term for a large class of k -partite k -graphs K ; in particular, this is true for all complete k -partite k -graphs. So for any complete k -partite k -graph K the asymptotic value of $\delta(K, n)$ is determined by the parameters $\gcd(K)$ and $\sigma(K)$, in the same way that the corresponding parameters determined $\delta(G, n)$ for any r -chromatic graph G . The same is true for many incomplete k -partite k -graphs K ; we discuss these k -graphs in Section 9, as well as discussing the value of $\delta(K, n)$ for those k -partite k -graphs for which the correct asymptotic value remains unknown (of course, Theorems 1.1, 1.2 and 1.3 still provide an upper bound on $\delta(K, n)$ in these cases). For those k -partite k -graphs K for which we do determine the correct asymptotic value of $\delta(K, n)$, it is natural to ask whether the $o(n)$ error term can be removed. We conjecture that in fact each of Theorems 1.1, 1.2 and 1.3 is still valid if the αn error term is replaced by a sufficiently large constant C which depends only on K .

One special case of these results answers a question of Rödl and Ruciński [30, Problem 3.15], who asked for the value of $\delta(C_s^3, n)$, where C_s^3 denotes the loose cycle 3-graph of length s . (For general k , the loose cycle k -graph of length s , denoted C_s^k , is defined for any $s > 1$ to have $s(k-1)$ vertices $\{1, \dots, s(k-1)\}$ and s edges $\{j(k-1)+1, \dots, j(k-1)+k\}$ for $0 \leq j < s$, with addition taken modulo k .) In Section 8 we shall see that $\gcd(C_s^k) = 1$ for any $k \geq 3$ and $s \geq 2$ except for the case $k = 3$ and $s = 3$, in which case $\mathcal{S}(C_s^k) = \{2\}$ and so $\gcd(K)$ is undefined. So in all cases except for $k = s = 3$ the k -graph C_s^k satisfies the condition of Theorem 1.2, and in fact we will show that furthermore C_s^k belongs to the class of k -graphs for which Theorem 1.2 is best possible up to the error term. By modifying our arguments to handle the case $k = s = 3$ separately, we obtain the following theorem.

Theorem 1.4. *For integers $k \geq 3$ and $s \geq 2$ we have*

$$\delta(C_s^k, n) = \begin{cases} \frac{n}{2(k-1)} + o(n) & \text{if } s \text{ is even, and} \\ \frac{s+1}{2s(k-1)}n + o(n) & \text{otherwise.} \end{cases}$$

Note that C_2^3 is identical to the 3-graph $K_4^3 - 2e$; as described earlier, the result above was proved in this case (that is, for $k = 3$ and $s = 2$) by Kühn and Osthus [21], and more recently Czygrinow, DeBiasio and Nagle [4] gave the exact value of $\delta(C_2^3, n)$ for large n .

The final results of this paper, in Section 9, concern the problem of finding a K -packing covering all but a constant number of vertices of a large k -graph H . By adapting the methods used for our results on perfect packings, we find that the minimum codegree requirement of Theorem 1.2 (which applied only to k -partite k -graphs K with $\gcd(K) = 1$) is sufficient to ensure such a K -packing for any k -partite k -graph K . More specifically, we have the following theorem.

Theorem 1.5. *Let K be a k -partite k -graph. Then there exists a constant $C = C(K)$ such that for any $\alpha > 0$ there exists $n_0 = n_0(K, \alpha)$ such that any k -graph H on $n \geq n_0$ vertices with $\delta(H) \geq \sigma(K)n + \alpha n$ admits a K -packing covering all but at most C vertices of H .*

By modifying a construction from Section 2, we will further see that Theorem 1.5 is asymptotically best possible for a large class of k -partite k -graphs which includes all complete k -partite k -graphs.

The results of this paper are significant as they provide the first cases other than that of a perfect matching for which the value of the well-studied parameter $\delta(H, n)$ is known even asymptotically for a k -graph H on more than four vertices. Furthermore, the diverse behaviour of this parameter over different k -partite k -graphs, according to the divisibility properties of the different vertex class sizes, is interesting in itself and increases our understanding of the extensive role such divisibility conditions play in a wide variety of problems involving the embedding of a spanning subgraph in a large k -graph H (see [16] for further discussion of this point). The proofs in this paper also demonstrate techniques for making use of the recent hypergraph blow-up lemma of Keevash [13], particularly the techniques used in Section 6.3 to delete copies of K so as to meet certain divisibility conditions.

1.5. Layout of the paper. In Section 2 we give constructions which show that the lowest upper bound provided by Theorems 1.1, 1.2 and 1.3 is asymptotically best possible for all complete k -partite k -graphs. Then, in Section 3 we state Lemmas 3.1 and 3.2 which are similar to Theorems 1.1, 1.2 and 1.3, but which pertain only to certain complete k -partite k -graphs. We deduce Theorems 1.1, 1.2 and 1.3 from these lemmas; having done so, we can focus solely on these complete k -partite k -graphs in proving Lemmas 3.1 and 3.2 (complete k -partite k -graphs are simpler to deal with as they have only one k -partite realisation). In

Section 4 we outline how the proofs of Lemmas 3.1 and 3.2 will proceed. These proofs make extensive use of hypergraph regularity; in particular, we use the recent hypergraph blow-up lemma due to Keevash [13]. The necessary background for the use of these tools is given in Section 5. Section 6 then gives a number of auxiliary lemmas which will be needed in the proofs of Lemmas 3.1 and 3.2, after which we prove these lemmas in Section 7. In Section 8 we turn to the problem of a loose cycle packing, proving Theorem 1.4, and the final section, Section 9 consists of concluding remarks. Firstly, we consider non-complete k -partite k -graphs, identifying large classes of such k -graphs for which the bounds of Theorems 1.1, 1.2 and 1.3 are asymptotically best possible. Following this we consider the question of finding a K -packing covering all but a constant number of vertices of a large k -graph G , proving Theorem 1.5. Finally, we briefly discuss the problem of finding $\delta(H, n)$ for k -graphs H which are not k -partite.

1.6. Notation. Throughout this paper, when we speak of ‘deleting’ a k -graph K from a k -graph H , we mean that both the vertices and edges of K are deleted from H , so what remains is the subgraph of H induced by the undeleted vertices. Also, for a k -graph H we define the *adjacency graph* $\text{Adj}(H)$ to be the graph on $V(H)$ where there is an edge between two vertices i and j if and only if some edge of H contains both i and j . We say that H is *connected* if $\text{Adj}(H)$ is connected.

We write vectors in bold font, and write, for example, v_j for the j th coordinate of \mathbf{v} . We write \mathbf{u}_j for the unit vector whose j th coordinate is one and whose other coordinates are all zero (the dimension of \mathbf{u}_j will always be clear from the context). Whenever we speak of a partition of a set, we implicitly fix an order of the parts of this partition. We write $[r]$ to denote the set of integers from 1 to r . For a set A , we use $\binom{A}{k}$ to denote the collection of subsets of A of size k , and similarly $\binom{A}{\leq k}$ to denote the collection of subsets of A of size at most k . We write $x = y \pm z$ to mean that $y - z \leq x \leq y + z$, and write $o(n)$ to denote a function which tends to zero as $n \rightarrow \infty$. Also, we use $x \ll y$ to mean for any $y \geq 0$ there exists $x_0 \geq 0$ such that for any $x \leq x_0$ the following statement holds, and similar statements with more constants are defined similarly. Finally, we omit floors and ceilings throughout this paper whenever they do not affect the argument.

2. EXTREMAL EXAMPLES

In this section we shall give constructions which demonstrate that the upper bound on $\delta(K, n)$ provided by Theorems 1.1, 1.2 and 1.3 is asymptotically best possible for all complete k -partite k -graphs, and many others besides.

For ease of discussion, we divide all k -partite k -graphs into *types*. Indeed, let K be a k -partite k -graph with at least one edge. Then we say that K is *type 0* if $\gcd(\mathcal{S}(K)) > 1$ or if K consists of k vertices and one edge (in which case a K -packing is a matching). Note that the latter condition is equivalent to saying that $\mathcal{S}(K) = \{1\}$. If K is not type 0, then for any $d \geq 1$ we say that K is *type d* if $\gcd(\mathcal{S}(K)) = 1$ and $\gcd(K) = d$. Observe that every k -partite k -graph K with at least one edge falls into precisely one of these types, since $\gcd(K)$ is defined for any K which is not type 0. Also note that the definitions of $\mathcal{S}(K)$ and $\gcd(K)$ immediately imply that $\gcd(\mathcal{S}(K))$ divides $\gcd(K)$, so we cannot have $\gcd(\mathcal{S}(K)) > 1$ and $\gcd(K) = 1$. Table 1 displays the asymptotic upper bounds provided for k -partite k -graphs of each type by Theorems 1.1, 1.2 and 1.3. The results of this section will show that for a large class of k -partite k -graphs, which includes all complete k -partite k -graphs, these bounds are best possible up to the $o(n)$ error terms.




FIGURE 1. The construction of Proposition 2.2, shown for $p = 2, k = 3$ on the left and for $p = 3, k = 4$ on the right. The left hand construction is also used for Proposition 2.1 for $k = 3$. In each case the k -graph of the construction has all k -tuples of the forms shown as edges (so, for example, the edges of the 3-graph on the left are all 3-tuples with either 1 or 3 vertices in V_1).

Our first construction is well-known and gives a condition (P1) on a k -partite k -graph K which is sufficient to ensure that the bound given by Theorem 1.1 is asymptotically tight. The left hand part of Figure 1 gives an illustration of this construction.

Proposition 2.1. *Let $p > 1$, and let K be a k -partite k -graph on vertex set U such that*

(P1) *each set $A \subseteq U$ for which $|e \cap A|$ is even for every $e \in K$, has size divisible by p .*

Then for any n there exists a k -graph G on n vertices with $\delta(G) \geq n/2 - k$ such that G does not contain a perfect K -packing.

Proof. Let V_1 and V_2 be disjoint sets of vertices with $|V_1 \cup V_2| = n$ such that $|V_1|, |V_2| \geq n/2 - 1$ and p does not divide $|V_2|$. Let G be the k -graph on vertex set $V_1 \cup V_2$ whose edges are all k -tuples $e \in \binom{V_1 \cup V_2}{k}$ such that $|V_2 \cap e|$ is even. Then $\delta(G) \geq \min\{|V_1|, |V_2|\} - (k-1) \geq n/2 - k$. Moreover, for any copy K' of K in G , every edge $e \in K'$ is an edge of G , and so $|V_2 \cap e|$ is even. By our assumption on K this implies that p divides $|V_2 \cap V(K')|$, and so the number of vertices of V_2 covered by any K -packing in G is divisible by p . Since p does not divide $|V_2|$, we conclude that G does not contain a perfect K -packing. \square

Suppose that K is a complete k -partite k -graph of type 0 with vertex classes U_1, \dots, U_k . If K consists of just one edge then (P1) is trivially satisfied for $p = 2$; otherwise there exists some $p > 1$ such that p divides $|U_j|$ for every $j \in [k]$. Let $A \subseteq U := \bigcup_{i \in [k]} U_i$ be such that $|e \cap A|$ is even for every $e \in K$. For any $i \in [k]$ and any $x, y \in U_i$ we may choose vertices $u_j \in U_j$ for each $j \neq i$, and since K is complete both $\{x\} \cup \{u_j : j \neq i\}$ and $\{y\} \cup \{u_j : j \neq i\}$ are edges of K . Both these sets therefore have an even number of vertices in A , so we must have $x, y \in A$ or $x, y \notin A$. It follows that for each $i \in [k]$ we have either $U_i \subseteq A$ or $U_i \cap A = \emptyset$, and therefore that p divides $|A|$. We conclude that any complete k -partite k -graph K of type 0 satisfies (P1), and so Theorem 1.1 is asymptotically best possible for any such K .

Our next construction generalises the construction used in Proposition 2.1. For this we need the following definitions. For any integer $p \geq 2$, let \mathcal{V}_p be the $(p-1)$ -dimensional sublattice of \mathbb{Z}_p^p generated by the vectors $\mathbf{v}_1, \dots, \mathbf{v}_{p-1}$, where for each $1 \leq j \leq p-1$ we define

$$\mathbf{v}_j = \mathbf{u}_j + (j-1)\mathbf{u}_p = (\overbrace{0, \dots, 0}^{j-1}, 1, \overbrace{0, \dots, 0}^{p-j-1}, j-1).$$

The key property of the lattice \mathcal{V}_p is that, working in \mathbb{Z}_p^p ,

(†) for any $\mathbf{x} \in \mathbb{Z}_p^p$ there is precisely one $j \in [p]$ such that $\mathbf{x} + \mathbf{u}_j \in \mathcal{V}_p$.

That is, for any vector $\mathbf{x} \in \mathbb{Z}_p^p$ there is precisely one coordinate of \mathbf{x} which, if incremented by one (modulo p), yields a vector $\mathbf{y} \in \mathcal{V}_p$. To see this, let $\mathbf{x} \in \mathbb{Z}_p^p$. Then it follows immediately from the definition of \mathcal{V}_p that there is $\mathbf{y}' \in \mathcal{V}_p$ such that \mathbf{x} and \mathbf{y}' differ only in their last coordinates x_p and y'_p (or do not differ at all). Let $d \in [p]$ be such that $x_p - y'_p \equiv d - 1$ modulo p , and define

$$\mathbf{y} := \begin{cases} \mathbf{y}' + \mathbf{v}_d & \text{if } d \leq p - 1 \\ \mathbf{y}' & \text{if } d = p. \end{cases}$$

Then in any case we have $\mathbf{y} \in \mathcal{V}_p$. Moreover, if $d = p$ then $x_p - y_p = x_p - y'_p \equiv -1$ modulo p , so $\mathbf{y} = \mathbf{x} + \mathbf{u}_p$. On the other hand, if $d \neq p$ then $y_p \equiv y'_p + (\mathbf{v}_d)_p \equiv y'_p + d - 1 \equiv x_p$ modulo p , so \mathbf{y} and \mathbf{x} differ only in coordinate d , with $y_d \equiv x_d + 1$ modulo p , and therefore $\mathbf{y} = \mathbf{x} + \mathbf{u}_d$. This proves that for any $\mathbf{x} \in \mathbb{Z}_p^p$ there is at least one $j \in [p]$ such that $\mathbf{x} + \mathbf{u}_j \in \mathcal{V}_p$. If for some $\mathbf{x} \in \mathbb{Z}_p^p$ there were $j \neq j'$ such that $\mathbf{x} + \mathbf{u}_j \in \mathcal{V}_p$ and $\mathbf{x} + \mathbf{u}_{j'} \in \mathcal{V}_p$, then we would obtain $(\mathbf{x} + \mathbf{u}_j) - (\mathbf{x} + \mathbf{u}_{j'}) = \mathbf{u}_j - \mathbf{u}_{j'} \in \mathcal{V}_p$. However, it is easily checked that $\mathbf{u}_j - \mathbf{u}_{j'} \notin \mathcal{V}_p$ for any $j \neq j'$, proving (\dagger) .

If \mathcal{P} is a partition of a set X into parts X_1, \dots, X_p , for any $S \subseteq X$ we define the *index vector of S with respect to \mathcal{P}* , denoted $\mathbf{i}_{\mathcal{P}}(S)$, to be the vector in \mathbb{Z}_p^p whose j -th coordinate is $|S \cap X_j|$ modulo p ; this is well-defined since we consider \mathcal{P} to include an order on its parts. We sometimes omit the subscript \mathcal{P} and write simply $\mathbf{i}(S)$ if \mathcal{P} is clear from the context.

Proposition 2.2. *Let $p \geq 2$ and let K be a k -partite k -graph on vertex set U such that*

- (P2) *for any partition \mathcal{P} of U into p parts such that $\mathbf{i}(e) \in \mathcal{V}_p$ for every $e \in K$ we must also have $\mathbf{i}(U) \in \mathcal{V}_p$.*

Then for any n there exists a k -graph G on n vertices with $\delta(G) \geq n/p - k$ such that G does not contain a perfect K -packing.

Proof. Let V be a set of n vertices, and choose a partition \mathcal{P} of V into parts V_1, \dots, V_p such that $\mathbf{i}(V) \notin \mathcal{V}_p$ and $|V_j| \geq n/p - 1$ for each $j \in [p]$. Let G be the k -graph on vertex set V such that a k -tuple $e \in \binom{V}{k}$ is an edge of G precisely if $\mathbf{i}(e) \in \mathcal{V}_p$ (see Figure 1 for two illustrations of this construction). Then for any $(k-1)$ -tuple $e' \in \binom{V}{k-1}$ we can choose $j \in [p]$ by (\dagger) such that $\mathbf{i}(e') + \mathbf{u}_j \in \mathcal{V}_p$, and then adding any of the $|V_j \setminus e'| \geq n/p - k$ vertices $v \in V_j \setminus e'$ to e' gives a k -tuple $e := e' \cup \{v\}$ such that $\mathbf{i}(e) = \mathbf{i}(e') + \mathbf{u}_j \in \mathcal{V}_p$, that is, an edge $e \in G$. So $\delta(G) \geq n/p - k$. Now, for any copy K' of K in G , every edge $e \in K'$ is an edge of G and so has the property that $\mathbf{i}(e) \in \mathcal{V}_p$. By (P2) it follows that $\mathbf{i}(V(K')) \in \mathcal{V}_p$. So if \mathcal{F} is a K -packing in G , then $\mathbf{i}(V(\mathcal{F})) = \sum_{K' \in \mathcal{F}} \mathbf{i}(V(K')) \in \mathcal{V}_p$; since $\mathbf{i}(V) \notin \mathcal{V}_p$ this implies that \mathcal{F} is not perfect. \square

Note in particular that for $p = 2$ the k -graph G constructed in the proof of Proposition 2.2 is the same as that in Proposition 2.1. A similar argument as above shows that any complete k -partite k -graph K of type $d \geq 2$ satisfies (P2) for any p which divides d . Indeed, let U_1, \dots, U_k be the vertex classes of K , and suppose that sets V_1, \dots, V_p partition $V(K)$ such that (taking index vectors with respect to this partition) $\mathbf{i}(e) \in \mathcal{V}_p$ for every $e \in K$. Fix any $j \in [k]$ and any $u, v \subseteq U_j$; then we may choose edges $e, e' \in K$ such that $u \in e$, $v \in e'$ and $e \setminus \{u\} = e' \setminus \{v\} =: e^*$. Since $\mathbf{i}(e^*) \in \mathbb{Z}_p^p$, by (\dagger) there is precisely one $i \in [k]$ such that adding a vertex of V_i to e^* gives an edge whose index vector lies in \mathcal{V}_p . Since $\mathbf{i}(e), \mathbf{i}(e') \in \mathcal{V}_p$, we must have $u, v \in V_i$ for this i , and so we conclude that every vertex class U_j of K must be a subset of some V_i . Since K has type d we have that p divides $\gcd(K) = d$, and so each vertex class U_j has equal size b_1 modulo p . So we must have $\mathbf{i}(V(K)) = b_1 \mathbf{i}(e) \in \mathcal{V}_p$, proving that K indeed satisfies property (P2). So, up to the $o(n)$ error term, the bound of $n/p + o(n)$ given in Theorem 1.3 is best possible for any complete k -partite k -graph of type $d \geq 2$.




FIGURE 2. The construction of Proposition 2.3, illustrated for $k = 3$. The edges of this 3-graph are all 3-tuples which intersect A .

The final construction we use is well-known. For this, we write $\tau(K)$ to denote the proportion of vertices of K contained in a smallest vertex cover of K . That is, $\tau(K) = |S|/b$, where $b = |V(K)|$ and $S \subseteq V(K)$ is a set of minimum size with the property that every edge of K contains a vertex of S (we express $\tau(K)$ as a proportion for comparison with $\sigma(K)$).

Proposition 2.3. *For any k -graph K and any n there exists a k -graph G on n vertices with $\delta(G) = \lceil \tau(K)n \rceil - 1$ such that G does not contain a perfect K -packing.*

Proof. Write $b = |V(K)|$ and $\tau = \tau(K)$. Let A and B be disjoint sets of vertices such that $|A| = \lceil \tau n \rceil - 1$ and $|A \cup B| = n$, and let G be the k -graph on vertex set $A \cup B$ whose edges are all k -tuples $e \in \binom{A \cup B}{k}$ such that $|e \cap A| \geq 1$ (this construction is illustrated in Figure 2). Then $\delta(G) = |A| = \lceil \tau n \rceil - 1$. Let K' be a copy of K in G . Then any edge $e \in K'$ must contain a vertex of A . So $A \cap V(K')$ is a vertex cover of K' , so $|A \cap V(K')| \geq \tau b$. Any K -packing in G therefore has size at most $|A|/\tau b < n/b$, so is not perfect. \square

This means that for any k -graph K with $\tau(K) = \sigma(K)$ we have $\delta(K, n) \geq \sigma(K)n$. Together with Theorems 1.2 and 1.3 this determines $\delta(K, n)$ asymptotically for any such k -partite k -graph K which has type 1, or which has type $d \geq 2$ and also satisfies property (P2) with p being the smallest prime factor of d . Note that in particular we have $\tau(K) = \sigma(K)$ for any complete k -partite k -graph K . To see this, let K be a complete k -partite k -graph on b vertices with vertex classes U_1, \dots, U_k . Since K is complete a subset $S \subseteq V(K)$ is a vertex cover if and only if $U_j \subseteq S$ for some $j \in [k]$. So $\tau(K) = \min_{j \in [k]} |U_j|/b = \sigma(K)$, as required.

In conclusion, these examples show that the bound given by Theorems 1.1, 1.2 and 1.3 is best possible up to the $o(n)$ error term for all complete k -partite k -graphs. We consider non-complete k -partite k -graphs further in Section 9.

3. REDUCTION TO COMPLETE k -PARTITE k -GRAPHS

For simplicity, we would like to restrict our attention to complete k -partite k -graphs alone, as these have only one k -partite realisation (up to permutations of the vertex classes) and so are easier to work with. Clearly if U_1, \dots, U_k are the vertex classes of a k -partite realisation of a k -graph K , then K is a spanning subgraph of the complete k -partite k -graph K' on the same vertex classes, and so if G contains a K' -packing then G contains a K -packing also. However, we cannot deduce from this that proving Theorems 1.1, 1.2 and 1.3 for complete k -partite k -graphs would imply that these theorems hold for all k -partite k -graphs, as it may well be the case that $\gcd(K) \neq \gcd(K')$. Instead, in this section we shall state two lemmas (Lemmas 3.1 and 3.2) which essentially say that Theorems 1.1, 1.2 and 1.3 hold for certain complete k -partite k -graphs. We shall then deduce Theorems 1.1, 1.2 and 1.3 from these two lemmas, showing that it is indeed sufficient to consider only these complete k -partite k -graphs for the rest of the paper (in which we prove Lemmas 3.1 and 3.2).

Lemma 3.1. *Let K be the complete k -partite k -graph whose vertex classes have sizes b_1, \dots, b_k , where these sizes are not all equal, and suppose that $\gcd(K)$ and b_1 are coprime. Then for any $\alpha > 0$ there exists $n_0 = n_0(K, \alpha)$ such that the following statement holds. Let H be a k -graph on $n \geq n_0$ vertices such that*

- (a) $b := b_1 + \dots + b_k$ divides n ,
- (b) $\delta(H) \geq \sigma(K)n + \alpha n$, and
- (c) if $\gcd(K) > 1$, then $\delta(H) \geq n/p + \alpha n$, where p is the smallest prime factor of $\gcd(K)$.

Then H contains a perfect K -packing.

Lemma 3.2. *Let K be the complete k -partite k -graph whose vertex classes each have size b_1 . Then for any $\alpha > 0$ there exists $n_0 = n_0(K, \alpha)$ such that if $n \geq n_0$ is divisible by $b_1 k$ and H is a k -graph on n vertices with $\delta(H) \geq n/2 + \alpha n$ then H contains a perfect K -packing.*

For the rest of this section we seek to deduce Theorems 1.1, 1.2 and 1.3 from Lemmas 3.1 and 3.2. For this we shall need the following fact of elementary number theory.

Fact 3.3. *For any positive integers r_1, \dots, r_k there exist integers a_1, \dots, a_k such that $a_1 r_1 + \dots + a_k r_k = \gcd(\{r_1, \dots, r_k\})$.*

Let \mathcal{F} be a collection of k -graphs. Then we say that a k -graph G contains an \mathcal{F} -packing if G can be packed with members of \mathcal{F} . More precisely, an \mathcal{F} -packing in G is a collection of pairwise vertex-disjoint subgraphs F_1, \dots, F_r of G so that each F_j is in \mathcal{F} (that is, F_j is isomorphic to a member of \mathcal{F}). We say that an \mathcal{F} -packing of G is *perfect* if it covers every vertex of G . This naturally generalises the notion of an H -packing for a k -graph H , as an H -packing of G and an $\{H\}$ -packing of G are identical. The following elementary proposition implies that to demonstrate that G contains a perfect H -packing it is sufficient to show that G contains a perfect \mathcal{F} -packing for some family \mathcal{F} such that every $F \in \mathcal{F}$ contains a perfect H -packing (we omit the simple proof).

Proposition 3.4. *Suppose that G and H are k -graphs and that \mathcal{F} is a collection of k -graphs such that*

- (i) G contains a perfect \mathcal{F} -packing, and
- (ii) every $F \in \mathcal{F}$ contains a perfect H -packing.

Then G contains a perfect H -packing.

To deduce Theorems 1.1, 1.2 and 1.3 from Lemmas 3.1 and 3.2, we make use of the following complete k -partite k -graphs, which will also play important roles later on in the paper.

Definition 3.5. *Fix an integer $k \geq 3$.*

- (i) *For any integer m , we define the balanced k -partite k -graph $\mathcal{B}(m)$ to be the complete k -partite k -graph on vertex classes W_1, \dots, W_k , where $|W_1| = |W_2| = \dots = |W_k| = m$. So $\mathcal{B}(m)$ has km vertices.*
- (ii) *Likewise, for integers m and d with $d < m$, we define the d -unbalanced k -partite k -graph $\mathcal{U}(m, d)$ to be the complete k -partite k -graph on vertex classes W_1, \dots, W_k , where*

$$|W_1| = m - d, |W_2| = m + d, \text{ and } |W_3| = \dots = |W_k| = m.$$

So $\mathcal{U}(m, d)$ has km vertices also.

- (iii) *Finally, let m be an integer and $0 < \sigma < 1$. Then we define the σ -lopsided k -partite k -graph $\mathcal{L}(m, \sigma)$ to be the complete k -partite k -graph on vertex classes W_1, \dots, W_k , where*

$$|W_1| = \sigma m, \text{ and } |W_2| = \dots = |W_k| = \frac{(1 - \sigma)m}{k - 1},$$

provided that these vertex class sizes are each integers (otherwise $\mathcal{L}(m, \sigma)$ is undefined). So $\mathcal{L}(m, \sigma)$ has m vertices.

Note that the definitions of $\mathcal{B}(m)$, $\mathcal{U}(m, d)$ and $\mathcal{L}(m, \sigma)$ each depend on k ; this dependence is suppressed in our notation as k will always be clear from the context.

Given a k -partite k -graph K , some special cases of the above definition will be of particular importance; we therefore define $\mathcal{B}(K) := \mathcal{B}(b)$ and $\mathcal{L}(K) := \mathcal{L}((k-1)!b, \sigma(K))$, where b denotes the number of vertices of K . Finally, if $\gcd(K)$ is defined, then we define $\mathcal{U}_s(K) := \mathcal{U}(sb, \gcd(K))$ for those integers s for which the k -graph $\mathcal{U}_s(K)$ so defined admits a perfect K -packing; the next proposition tells us that this is the case for any sufficiently large s .

We note for future reference that $\mathcal{B}(K)$, $\mathcal{U}_s(K)$ and $\mathcal{L}(K)$ have kb , kbs and $(k-1)!b$ vertices respectively, that $\sigma(\mathcal{L}(K)) = \sigma(K)$, and that if $\gcd(K)$ divides s then $\gcd(\mathcal{U}_s(K)) = \gcd(K)$. Crucially, each of these k -partite k -graphs admits a perfect K -packing, as shown by the following proposition.

Proposition 3.6. *Let K be a k -partite k -graph on b vertices, and let $\sigma := \sigma(K)$. Then the k -graphs $\mathcal{B}(K) = \mathcal{B}(b)$ and $\mathcal{L}(K) = \mathcal{L}((k-1)!b, \sigma)$ each contain a perfect K -packing. Furthermore, if $d := \gcd(K)$ is defined then there exists $s_0 = s_0(K)$ such that for any $s \geq s_0$ the k -graph $\mathcal{U}_s(K) = \mathcal{U}(sb, d)$ contains a perfect K -packing.*

Proof. Let U_1, \dots, U_k be the vertex classes of a k -partite realisation of K . We form a k -partite k -graph K^* with vertex classes W_1, \dots, W_k as follows. Initially take W_1, \dots, W_k to be empty sets, and then add k vertex-disjoint copies of K to K^* , so that the vertices of U_i in the j th copy of K are added to W_{i+j} (with addition taken modulo k). That is, each vertex class of K^* receives the vertices of one copy of U_1 , one copy of U_2 , and so forth. So each vertex class of K^* has size b . We conclude that K^* is a spanning subgraph of $\mathcal{B}(b)$. By construction K^* contains a perfect K -packing, so $\mathcal{B}(b)$ contains a perfect K -packing also.

A similar argument holds for $\mathcal{L}((k-1)!b, \sigma)$. Indeed, by the definition of $\sigma(K)$ we may assume that $|U_1| = \sigma b$. Then we form a k -partite k -graph K^* consisting of $(k-1)!$ vertex-disjoint copies of K : for each permutation ρ of $[k]$ with $\rho(1) = 1$ we add a copy of K to K^* in which the vertices of U_j are included in $W_{\rho(j)}$ for each $j \in [k]$. So K^* has $(k-1)!b$ vertices in total; the first vertex class of K^* has size $(k-1)!|U_1| = (k-1)!b\sigma$, whilst each other vertex class of K^* has equal size

$$\frac{(k-1)!b - (k-1)!b\sigma}{k-1} = (k-2)!b(1-\sigma).$$

So K^* is a spanning subgraph of $\mathcal{L}((k-1)!b, \sigma)$. As before, since K^* contains a perfect K -packing by construction, $\mathcal{L}((k-1)!b, \sigma)$ contains a perfect K -packing also.

Finally we come to $\mathcal{U}(sb, d)$. For this we must consider all possible k -partite realisations χ of K ; let \aleph be the set formed by all such χ . We write $U_1^\chi, \dots, U_k^\chi$ for the vertex classes of the realisation χ . Note that we consider all possible realisations, not simply all possible realisations up to permutations of the vertex classes. In particular, this means that the number of realisations $N := |\aleph|$ is divisible by $k!$. Note also that $N \leq k^b$, and that by symmetry we have $\sum_{\chi \in \aleph} |U_j^\chi| = bN/k$ for each $j \in [k]$. In addition, recall that

$$d := \gcd(K) := \gcd(\{|U_1^\chi| - |U_2^\chi| : \chi \in \aleph\}).$$

So by Fact 3.3 we may choose integers a_χ for each k -partite realisation χ of K such that

$$\sum_{\chi \in \aleph} a_\chi (|U_1^\chi| - |U_2^\chi|) = d.$$

Let $a := \max_{\chi \in \mathbb{N}} a_\chi$. We now form a k -partite k -graph K^* similarly as before, with vertex classes W_1, \dots, W_k which we initially take to be empty sets. Then, for each realisation χ of K , add $a - a_\chi$ vertex-disjoint copies of K to K^* , with the vertices of U_j^χ added to W_j for each $j \in [k]$, and also add a_χ vertex-disjoint copies of K to K^* , with the vertices of U_1^χ added to W_2 , the vertices of U_2^χ added to W_1 , and the vertices of U_j^χ added to W_j for each $j \geq 3$. Then the total number of vertices added to W_1 is

$$\sum_{\chi \in \mathbb{N}} ((a - a_\chi)|U_1^\chi| + a_\chi|U_2^\chi|) = \sum_{\chi \in \mathbb{N}} a|U_1^\chi| - \sum_{\chi \in \mathbb{N}} a_\chi(|U_1^\chi| - |U_2^\chi|) = \frac{aNb}{k} - d.$$

In the same way the number of vertices added to W_2 is

$$\sum_{\chi \in \mathbb{N}} (a - a_\chi)|U_2^\chi| + a_\chi|U_1^\chi| = \frac{aNb}{k} + d,$$

and the number of vertices added to W_j for each $j \geq 3$ is $\sum_{\chi \in \mathbb{N}} a|U_j^\chi| = aNb/k$. So we may take $s_0 = aN/k$. Then K^* contains a perfect K -packing by construction, and is a spanning subgraph of $\mathcal{U}(s_0b, d)$, from which we conclude that $\mathcal{U}(s_0b, d)$ contains a perfect K -packing. Finally, for any $s \geq s_0$ observe that $\mathcal{U}(sb, d)$ admits a $\{\mathcal{B}(b), \mathcal{U}(s_0b, d)\}$ -packing consisting of $s - s_0$ copies of $\mathcal{B}(b)$ and one copy of $\mathcal{U}(s_0b, d)$; since we have already seen that both $\mathcal{B}(b)$ and $\mathcal{U}(s_0b, d)$ contain perfect K -packings it follows that $\mathcal{U}(sb, d)$ contains a perfect K -packing by Proposition 3.4. \square

We now have the definitions we need to derive Theorems 1.1, 1.2 and 1.3 from Lemmas 3.1 and 3.2. We shall also need the following weak version of a theorem of Erdős [8], which states that the Turán density of any k -partite k -graph is zero.

Theorem 3.7 ([8]). *For any k -partite k -graph K and any $\alpha > 0$ there exists n_0 such that any k -graph G on $n \geq n_0$ vertices with at least $\alpha \binom{n}{k}$ edges contains a copy of K .*

Proof of Theorem 1.1. We may assume that $1/n \ll 1/k, 1/b, \alpha$. By repeated application of Theorem 3.7 we may delete at most k vertex-disjoint copies of K from G to obtain a subgraph H such that kb divides $n' := |V(H)|$. Since we deleted at most $kb \leq \alpha n/2$ vertices in forming H we have $\delta(H) \geq n/2 + \alpha n/2 \geq n'/2 + \alpha n'/2$. So H contains a perfect $\mathcal{B}(K)$ -packing by Lemma 3.2 (applied with $\mathcal{B}(K), n'$ and $\alpha/2$ in place of K, n and α respectively). Together with the deleted copies of K this gives a perfect $\{\mathcal{B}(K), K\}$ -packing of G , and G therefore contains a perfect K -packing by Propositions 3.4 and 3.6. \square

Proof of Theorems 1.2 and 1.3. We may assume that $1/n \ll 1/k, 1/b, \alpha$. Introduce new constants m and s with $1/n \ll 1/m \ll 1/s \ll 1/k, 1/b, \alpha$ such that both m and s are divisible by $\gcd(K)$. Then we may assume that s is large enough for $\mathcal{U}_s(K)$ to be defined and so to contain a perfect K -packing by Proposition 3.6.

Our first step is to form a complete k -partite k -graph K_1 with vertex class sizes b_1, \dots, b_k such that K_1 admits a perfect K -packing and such that b_1 and $\gcd(K)$ are coprime. For Theorem 1.2 we have $\gcd(K) = 1$ so we may simply take any k -partite realisation of K and add edges to form a complete k -partite k -graph K_1 . So assume that $\gcd(K) > 1$ and $\gcd(\mathcal{S}(K)) = 1$, as in Theorem 1.3, and let s_1, \dots, s_t be the elements of $\mathcal{S}(K)$. By Fact 3.3 we may choose integers $a_1, \dots, a_t \geq 0$ such that $a_1 s_1 + \dots + a_t s_t$ is coprime to $\gcd(K)$, and since $1/s \ll 1/k, 1/b$ we may do this so that $a_1 + \dots + a_t \leq s$. By definition of $\mathcal{S}(K)$, for each $i \in [t]$ we may choose a k -partite realisation χ_i of K whose k vertex classes have sizes $b_1^i, b_2^i, \dots, b_k^i$ with $b_1^i = s_i$. Let K_1 be the complete k -partite k -graph whose j th vertex class

has size $b_j := \sum_{i \in [t]} a_i b_j^i$. Then by construction K_1 has a perfect K -packing consisting of a_1 copies of K with realisation χ_1 , a_2 copies of K with realisation χ_2 , and so forth. Moreover, $b_1 = \sum_{i \in [t]} a_i b_1^i = \sum_{i \in [t]} a_i s_i$, so b_1 is coprime to $\gcd(K)$. So K_1 has the desired properties; note also that K_1 has at most sb vertices.

We next form a complete k -partite k -graph K_2 with vertex class sizes b'_1, \dots, b'_k such that K_2 admits a perfect $\{\mathcal{U}_s(K), K_1\}$ -packing, $\gcd(K_2) = \gcd(K)$ and b'_1 and $\gcd(K)$ are coprime. For this observe that, since K_1 has a perfect K -packing, the definition of $\gcd(K)$ implies that $b_i - b_j$ is divisible by $\gcd(K)$ for any $i, j \in [k]$. Without loss of generality we may assume that $b_2 \geq b_3$. Fix $d := (b_2 - b_3)/\gcd(K)$, and define

$$\begin{aligned} b'_1 &:= b_1 + (d+1)(bs + \gcd(K)), \\ b'_2 &:= b_2 + (d+1)(bs - \gcd(K)), \text{ and} \\ b'_i &:= b_i + (d+1)bs \text{ for } 3 \leq i \leq k. \end{aligned}$$

So in particular $b'_i \leq 3b^3s$ for any $i \in [k]$. Let K_2 be the complete k -partite k -graph with vertex class sizes b'_1, \dots, b'_k . Then K_2 admits a perfect $\{\mathcal{U}_s(K), K_1\}$ -packing consisting of $d+1$ copies of $\mathcal{U}_s(K)$ and one copy of K_1 . Also, since b_1 and $\gcd(K)$ were coprime, and $\gcd(K)$ divides s , we find that b'_1 and $\gcd(K)$ are coprime. Finally, observe that $b'_3 - b'_2 = b_3 - b_2 + (d+1)\gcd(K) = \gcd(K)$. So $\gcd(K_2) \leq \gcd(K)$, and from the definition of b'_i for $i \in [k]$ and the fact that $\gcd(K)$ divides s we see that $\gcd(K)$ divides $b'_i - b'_j$ for any $i, j \in [k]$, from which we conclude that $\gcd(K_2) = \gcd(K)$. So K_2 has the desired properties.

Define b''_i for $i \in [k]$ by

$$\begin{aligned} b''_1 &= (k-1)!b\sigma(K)m + b'_1, \text{ and} \\ b''_i &= (k-2)!b(1 - \sigma(K))m + b'_i \text{ for } 2 \leq i \leq k, \end{aligned}$$

and let K_3 be the complete k -partite k -graph with vertex class sizes b''_1, \dots, b''_k . Then K_3 admits a perfect $\{\mathcal{L}(K), K_2\}$ -packing consisting of one copy of K_2 and m copies of $\mathcal{L}(K)$; by Propositions 3.4 and 3.6 and the fact that K_2 admits a perfect $\{\mathcal{U}_s(K), K_1\}$ -packing it follows that K_3 admits a perfect K -packing. Also, since b'_1 and $\gcd(K)$ were coprime, and $\gcd(K)$ divides m , we find that b''_1 and $\gcd(K)$ are coprime. Furthermore, $b''_3 - b''_2 = b'_3 - b'_2 = \gcd(K)$, and since $\gcd(K)$ divides m we deduce that $\gcd(K_3) = \gcd(K)$ also. Finally, $|V(K_3)| = (k-1)!bm + |V(K_2)| \geq (k-1)!bm$, and so

$$\sigma(K_3) \leq \frac{b''_1}{|V(K_3)|} \leq \frac{(k-1)!b\sigma(K)m + b'_1}{(k-1)!bm} = \sigma(K) + \frac{3b^3s}{(k-1)!bm} \leq \sigma(K) + \frac{\alpha}{3}.$$

By Theorem 3.7 we may arbitrarily choose and delete from G at most $|V(K_3)|/b = |V(\mathcal{L}(K))|m/b + |V(K_2)|/b \leq (k-1)!m + 3kb^2s \leq \alpha n/3b$ copies of K so that the set $V' \subseteq V(G)$ of undeleted vertices is such that $|V(K_3)|$ divides $|V'|$. Also $H := G[V']$ has $\delta(H) \geq \delta(G) - \alpha n/3 \geq \sigma(K)n + 2\alpha n/3 \geq \sigma(K_3)n + \alpha n/3$. Similarly, if $\gcd(K) > 1$ then $\delta(H) \geq n/p + 2\alpha n/3$, where p is the smallest prime factor of $\gcd(K) = \gcd(K_3)$. So we may apply Lemma 3.1 with $K_3, \alpha/3$ and $|V'|$ in place of K, α and n respectively to obtain a perfect K_3 -packing in H . Together with the deleted copies of K this gives a perfect $\{K_3, K\}$ -packing of G , and G therefore contains a perfect K -packing by Proposition 3.4. \square

4. OUTLINE OF THE PROOFS

The proofs of Lemmas 3.1 and 3.2 use strong hypergraph regularity and the recent hypergraph blow-up lemma due to Keevash. The broad outline of how these are used will be familiar to those acquainted with the use of the blow-up lemma in graphs, but this method




FIGURE 3. The left hand diagram illustrates Φ in the case $k = 3$, whilst the diagram on the right shows a copy of Φ located within four vertex classes V_1, V_2, V_3 and V_4 . If there are many copies of Φ of the form shown, then we say that the triple $(1, S, 4)$ is Φ -dense, where $S = \{2, 3\}$.

remains relatively novel for hypergraphs (for which there are many additional technicalities and subtleties). In this section we give a rough outline of how these proofs proceed.

4.1. Proof outline for Lemma 3.1.

The proof proceeds through the following steps.

Apply the Regular Approximation Lemma: The first step is to apply the Regular Approximation Lemma (Theorem 5.1) to H . This returns both a partition of $V(H)$ into ‘clusters’ U_1, \dots, U_m , and a k -graph G on $V(H)$, with the following properties. Firstly, G is close to H , meaning that almost all edges of G are edges of H , and vice versa. Secondly, this partition is regular for G , meaning (loosely speaking) that for the purposes of embedding small subgraphs in G , the k -partite subgraph of G induced by any k -tuple of clusters behaves like a random k -graph of similar density. We write $Z = G \triangle H$, so Z is a sparse graph which contains all the ‘bad’ edges of G which are not edges of H ; we will often choose copies of K in $G \setminus Z$, and by definition of Z these copies are also in H .

Having obtained G , Z and the partition of $V(H)$ into m clusters, we define a ‘reduced k -graph’ \mathcal{R} on $[m]$ (Definition 5.8). This has m vertices, one corresponding to each cluster, and the edges of \mathcal{R} are those k -tuples S for which the corresponding clusters induce a dense k -partite subgraph of G and a sparse k -partite subgraph of Z . Defined in this way, \mathcal{R} ‘almost inherits’ the minimum codegree condition of H , meaning that almost all $(k-1)$ -tuples of vertices in \mathcal{R} have almost the same degree (proportionately) as $(k-1)$ -tuples in H (Lemma 5.11).

One k -graph which will play an important role in our proof is the k -graph on $k+1$ vertices with two edges, which we denote by Φ (see Figure 3). We call the $k-1$ vertices in the intersection of these edges the *central vertices* of Φ , and the remaining two vertices of Φ are the *end vertices*. For vertices i, j of \mathcal{R} and a $(k-1)$ -tuple S of vertices of \mathcal{R} , we then say that the triple (i, S, j) is Φ -dense if there are many copies of Φ in G whose end vertices lie in the clusters U_i and U_j and whose central vertices lie in the clusters U_ℓ for $\ell \in S$, and the triple is Z -sparse if there are few edges of Z in either of the k -tuples of clusters corresponding to edges of Φ . We are particularly interested in Φ -dense and Z -sparse triples in \mathcal{R} , since we will be able to choose copies of K within these triples with some flexibility over the number of vertices which are embedded in U_i and U_j (indeed, $k-1$ of the vertex classes of K will be contained in the clusters U_ℓ for $\ell \in S$, whilst we will be able to choose how many vertices from the remaining vertex class are contained in each of U_i and U_j). Also, to keep track of these useful triples we define a graph \mathcal{S} on $[m]$, where each vertex corresponds to a cluster, and an edge ij indicates the existence some S for which the triple (i, S, j) is Φ -dense and Z -sparse. In Lemma 5.11 we show that the condition $\delta(H) \geq n/p + o(n)$ yields a minimum codegree

condition $\delta(\mathcal{S}) > m/p$ on \mathcal{S} . It follows that \mathcal{S} has fewer than p connected components if $\gcd(K) > 1$, a fact which plays a crucial role later in the proof.

Refine the regularity partition into ‘lopsided groups’: Our next step is to find an almost-perfect packing of \mathcal{R} with a specific k -graph $\mathcal{A}_{p,q}^k$ (where p and q are chosen depending on $\sigma(K)$). Lemma 6.1 shows that the degree condition on \mathcal{R} which is ‘inherited’ from the condition $\delta(H) \geq \sigma(K) + \alpha n$ is sufficient to guarantee such a packing. For each copy \mathcal{A} of $\mathcal{A}_{p,q}^k$ in this packing we then use Lemma 6.2 to partition the clusters covered by \mathcal{A} into kt ‘subclusters’ V_j^i , which are labelled so that the subclusters V_1^i, \dots, V_k^i are taken from clusters corresponding to an edge of \mathcal{A} (i.e. an edge of \mathcal{R}). This guarantees that the k -partite subgraphs G^i and Z^i induced by these k subclusters are dense and sparse respectively. Furthermore, whilst it is unavoidable that these k subclusters may have different sizes, the definition of $\mathcal{A}_{p,q}^k$ will allow us to ensure that each G^i has $\sigma(G^i) > \delta(H) - o(n) > \sigma(K) + o(n)$. That is, each G^i is ‘less lopsided’ than K . This is the point in the argument where this part of the minimum codegree assumption is used; a weaker condition would not suffice to guarantee the existence of subclusters with $\sigma(G^i) > \sigma(K)$.

Having obtained the kt subclusters V_j^i , we define a k -graph \mathcal{R}' and a graph \mathcal{S}' on vertex set $[t] \times [k]$ to correspond to \mathcal{R} and \mathcal{S} . Indeed, the vertex (i, j) of \mathcal{R}' and \mathcal{S}' corresponds to the subcluster (i, j) , and k subclusters form an edge of \mathcal{R}' if the k clusters from which these subclusters were taken form an edge of \mathcal{R} . Similarly, two subclusters form an edge of \mathcal{S}' if the two clusters from which these subclusters were taken form an edge of \mathcal{S} . This allows us to retain information about the regularity partition when working with the subclusters. It follows from the definition of \mathcal{S}' that the connected components of \mathcal{S}' correspond to those of \mathcal{S} , so \mathcal{S}' also has fewer than p connected components if $\gcd(K) > 1$.

Obtain robustly-universal complexes: Next, for each i we delete a small number of vertices from each subcluster V_j^i so that the k -partite k -graph $G^i \setminus Z^i$ restricted to the remaining vertices is ‘robustly universal’. This means that, even after the removal of a few more vertices, we can find any k -partite k -graph of bounded maximum degree in $G^i \setminus Z^i \subseteq H$ which we can find in the complete k -partite k -graph on the same vertex set. These deletions are achieved by Theorem 5.4, a result of Keevash [13] which conceals the use of the hypergraph blow-up lemma. We also ‘put aside’ a randomly-chosen set X consisting of a small number of vertices from each subcluster; these vertices are immune from deletion over the next two steps, and ensure that every vertex which is not deleted lies in many edges which are not deleted, which is a requirement for the application of robust universality.

Delete a K -packing covering bad vertices: At this point we deal with the small number of ‘bad vertices’, meaning those vertices in clusters which were not covered by our $\mathcal{A}_{p,q}^k$ -packing, as well as those vertices which were deleted to make the k -graphs $G^i \setminus Z^i$ robustly universal. For this, Lemma 6.4 shows that for any vertex v of H there is a copy of K in H which contains v ; this is a straightforward corollary of Theorem 3.7 (the well-known result of Erdős that k -partite k -graphs have Turán density zero). Using this, we greedily choose and delete copies of K in H which cover all the bad vertices but which only cover a small number of vertices from each subcluster. Following these deletions, all remaining vertices of H lie in $G^i \setminus Z^i$ for some i .

Delete a K -packing to ensure divisibility of cluster sizes: We now delete further copies of K in H so that, following these deletions, the number of vertices remaining in each subcluster is divisible by $bk \gcd(K)$ (recall that b is the order of K). Lemma 6.7 states that we can do this; loosely speaking, this is achieved by deleting a series of K -packings in H to achieve successively stronger divisibility conditions on the subcluster sizes (a more detailed outline of

the proof of this lemma is given in Section 6.3). If $\gcd(K) > 1$, then it is crucial for this that, as stated above, \mathcal{S}' has fewer than p components. For example, the k -graph G constructed for Proposition 2.2 would yield a graph \mathcal{S}' with p components corresponding to the parts V_1, \dots, V_p , and the point of the construction is that it is not possible to delete a K -packing in G so that every part has size divisible by p . Recall that \mathcal{S}' has at most p components since \mathcal{S} had minimum codegree $\delta(\mathcal{S}) > m/p$, and that this in turn was inherited from the minimum codegree condition $\delta(H) \geq n/p + \alpha n$ of H . This is the point in the argument where this part of the minimum codegree assumption is used, and a weaker condition would not suffice. However, if $\gcd(K) = 1$ then we do not need this part of the minimum codegree assumption.

Blow-up a perfect K -packing in the remaining k -graph: Certainly a K -packing has bounded vertex degree, so our robustly universal k -graphs $G^i \setminus Z^i$ each contain a perfect K -packing if and only if the complete k -partite k -graph on the same vertex set does also. To this end, Corollary 6.13 shows that for those k -partite k -graphs K which meet the conditions of Lemma 3.1, two properties are sufficient to ensure such a packing: firstly that $G^i \setminus Z^i$ should be ‘less lopsided’ than K , and secondly that each vertex class of $G^i \setminus Z^i$ should have size divisible by $b_k \gcd(K)$. Our partition into subclusters was chosen so that the first condition holds, whilst the final round of deletions described above ensures that the second condition holds also. We can therefore find a perfect K -packing in $G^i \setminus Z^i \subseteq H$ for each i ; these K -packings, together with the deleted copies of K , form a perfect K -packing in H . This completes the outline of the proof of Lemma 3.1.

4.2. Proof outline for Lemma 3.2. In this lemma K is instead a complete k -partite k -graph K whose vertex classes each have the same size b_1 . The proof of this lemma proceeds through the same steps as the proof of Lemma 3.1, though there are two principal differences. Firstly, rather than finding an $\mathcal{A}_{p,q}^k$ -packing in \mathcal{R} , we can now find simply a matching $M_{\mathcal{R}}$ in \mathcal{R} which covers almost all of the vertices of \mathcal{R} . In consequence, there is no need to divide the clusters into subclusters, or to define \mathcal{R}' and \mathcal{S}' ; we simply continue working with the clusters and the k -graph \mathcal{R} and graph \mathcal{S} . The second principal difference is that a complete k -partite k -graph G contains a perfect K -packing if and only if every vertex class of G has equal size and this common size is divisible by b_1 . So to find a perfect K -packing in our robustly universal k -graphs $G^i \setminus Z^i$ in the final step of the proof, it is not sufficient to delete copies of K in the penultimate step such that every cluster of $G^i \setminus Z^i$ has size divisible by $b_k \gcd(K)$; we must now ensure also that these clusters have the same size for any i . As a consequence we must be more precise in our definition of \mathcal{S} . Indeed, we now define a directed graph \mathcal{S}^+ whose vertices correspond to clusters, and whereas before an edge $ij \in \mathcal{S}$ indicated the existence of some $(k-1)$ -tuple S for which (i, S, j) is Φ -dense and Z -sparse, we now only have an edge $i \rightarrow j$ of \mathcal{S}^+ if this is true for $S = e(i) \setminus \{i\}$, where e is the edge of $M_{\mathcal{R}}$ which contains i . Then, similarly as before, the minimum degree condition $\delta(H) \geq n/2 + \alpha n$ implies that \mathcal{S}^+ has minimum outdegree $\delta^+(\mathcal{S}^+) > m/2$.

It would be possible to proceed by considering the directed graph \mathcal{S}^+ , but there are a number of additional problems which would arise in this case. Instead, we make use of the notion of ‘irreducibility’ of k -graphs containing a perfect matching, which was introduced by Keevash and Mycroft [16], and is presented in Section 5.6. Using the stronger minimum degree condition of Lemma 3.2, we can insist that the reduced k -graph \mathcal{R} is irreducible on the matching $M_{\mathcal{R}}$. Then, in Section 6.4 we show that, under this assumption, we need only consider the undirected base graph \mathcal{S} of \mathcal{S}^+ for the purposes of deleting K -packings to adjust cluster sizes. Using this, we prove Lemma 6.10, which shows that it is indeed possible to delete a K -packing in H so that, following these deletions, the k clusters corresponding to any edge of $M_{\mathcal{R}}$ have equal size. We then use Lemma 6.10 in place of Lemma 6.7 in the

proof of Lemma 3.2; all other steps of the proof proceed roughly as before. Note, however, that our use of irreducibility requires that almost all $(k-1)$ -tuples S of vertices of \mathcal{R} have $\deg_{\mathcal{R}}(S) > m/k$ (this condition is inherited from the minimum codegree of H), so it would not be possible to use this approach in the proof of Lemma 3.1.

5. REGULARITY AND THE BLOW-UP LEMMA

Much of the notation introduced in this section was first introduced by Rödl and Skokan [34] and by Rödl and Schacht [32, 33].

5.1. Hypergraphs, complexes and partitions. A *hypergraph* H consists of a vertex set $V(H)$ and an edge set $E(H)$, where every edge of H is a set of vertices of H . So a k -graph (as defined in Section 1) is a hypergraph in which all the edges have size k . As with k -graphs we frequently identify a hypergraph H with the set of its edges. So, for example, $e \in H$ means that e is an edge of H , and $|H|$ is the number of edges in H . Likewise, if G and H are hypergraphs on a common vertex set V then the hypergraph $G \setminus H$ is the hypergraph on V formed by removing from G any edge which also lies in H . For any hypergraph H and any $U \subseteq V(H)$, the *restriction of H to U* , denoted $H[U]$ is the hypergraph on vertex set U whose edges are those edges of H which are subsets of U . Also, recall that if H is a hypergraph with vertex set V , the *degree* $\deg_H(S)$ of a set $S \subseteq V(H)$ is defined to be the number of edges of H which contain S as a subset. The *maximum vertex degree* of H , denoted $\Delta_{\text{vertex}}(H)$, is then defined to be the maximum of $\deg_H(\{v\})$ taken over all vertices $v \in V(H)$; so every vertex of H is contained in at most $\Delta_{\text{vertex}}(H)$ edges of H . For any set of vertices X , we write $K(X)$ for the complete hypergraph on vertex set X , that is, the edges of $K(X)$ are all subsets of X .

Now let X be a set of vertices, and let \mathcal{Q} be a partition of X into r parts X_1, \dots, X_r . We say that a subset $S \subseteq X$ is \mathcal{Q} -*partite* if $|S \cap X_i| \leq 1$ for any $i \in [r]$. Similarly, we say that a hypergraph H on X is \mathcal{Q} -*partite* if every edge of H is \mathcal{Q} -partite, and we refer to the parts X_i of \mathcal{Q} as the *vertex classes* of H . We say that H is r -*partite* if it is \mathcal{Q} -partite for some partition \mathcal{Q} of X into r parts. For any \mathcal{Q} -partite set $S \subseteq X$ we define the *index* of S to be $i(S) := \{i \in [r] : |S \cap X_i| = 1\}$. So S intersects precisely those X_i for which $i \in i(S)$. Likewise, for any $A \subseteq [r]$ we write $X_A := \bigcup_{i \in A} X_i$, and define H_A to be the $|A|$ -graph with vertex set X_A whose edges are all edges of H of index A (note that H_A is naturally $|A|$ -partite with vertex classes X_i for $i \in A$). In particular, $K(X)_A$ is the complete $|A|$ -partite $|A|$ -graph with vertex classes X_i for $i \in A$.

A k -*complex* J is a hypergraph in which every edge has size at most k and which has the property that if $e_1 \in J$ and $e_2 \subseteq e_1$ then $e_2 \in J$ (so the edges of J form a simplicial complex). We refer to edges of size i as i -*edges*, and write J_i for the i -graph on $V(J)$ formed by the i -edges of J . Informally, it may be helpful to think of a k -complex J as consisting of ‘layers’ J_i for $0 \leq i \leq k$. So any edge e in the ‘ i th layer’ J_i of J lies ‘above’ i edges of J in the ‘ $(i-1)$ th layer’, namely those subsets of e of size $i-1$. The ‘top layer’ of a k -complex J will play a particularly important role; due to this we often write $J_=_$ in place of J_k to emphasise that this is the ‘top layer’. So $J_=_$ is a k -graph on $V(J)$. For any k -graph H we can naturally generate a k -complex H^{\leq} on $V(H)$, whose edges are all subsets of edges of H . Observe in particular that $(H^{\leq})_=_ = H$, and also that if H is \mathcal{Q} -partite for some partition \mathcal{Q} of $V(H)$ then H^{\leq} is \mathcal{Q} -partite also.

Now suppose again that \mathcal{Q} partitions a set of vertices X into r parts X_1, \dots, X_r , and that J is a hypergraph on X . For any $A \subseteq [r]$, the *absolute density of J at A* , denoted $d(J_A)$, is

the proportion of edges of $K(X)_A$ which are also edges of J_A . So

$$d(J_A) := \frac{|J_A|}{|K(X)_A|} = \frac{|J_A|}{\prod_{i \in A} |X_i|}.$$

If J is a k -complex then we also have the notion of *relative density*. Indeed, the *relative density of J at A* is the proportion of those edges which could feasibly be in J_A (in the sense that they are supported by ‘lower levels’ of J) which are actually edges of J_A . More precisely, we write J_A^* for the set of all edges $e \in K(X)_A$ such that every proper subset $e' \subset e$ is an edge of J . So J_A^* is the set of edges which could feasibly be in J_A (given the ‘lower levels’ of J), and we define the relative density of J at index A to be

$$d_A(J) := \frac{|J_A|}{|J_A^*|}.$$

(If the set J_A^* is empty then we instead define $d_A(J)$ to be zero.)

5.2. Partition complexes. Loosely speaking, the Regular Approximation Lemma states that any k -graph is close to another k -graph which can be split into pieces, each of which forms the ‘top level’ of a regular k -complex. For a graph G , this split involves simply a partition of the vertex set into a number of ‘clusters’, whereupon the edges between any pair of clusters form a regular bipartite graph. However, for a k -graph H (for $k \geq 3$) we must not only partition the vertices of H , but also the pairs of vertices of H , the triples of vertices of H , and so forth, up to $(k-1)$ -tuples of vertices of H . To keep track of these partitions we need the notion of a *partition complex*, which we now introduce.

Let X be a set of vertices, and let \mathcal{Q} partition X into parts X_1, \dots, X_r . Recall that for any $A \subseteq [r]$, $K(X)_A$ consists of all $|A|$ -tuples of vertices of X with index A . A *partition k -system \mathcal{P} on X* consists of a partition \mathcal{P}_A of the edges of $K(X)_A$ for each $A \subseteq [r]$ with $|A| \leq k$. We refer to the partition classes of \mathcal{P}_A as *cells*. So every edge of $K(X)_A$ is contained in precisely one cell of \mathcal{P}_A . We say that \mathcal{P} is *a-bounded* if for each A the partition \mathcal{P}_A has at most a cells. Also, for any $j \in [k]$ we write

$$\mathcal{P}^{(j)} = \bigcup_{A \in \binom{[r]}{j}} \mathcal{P}_A,$$

so $\mathcal{P}^{(j)}$ is a partition of the set of all \mathcal{Q} -partite j -tuples of vertices of X . Note in particular that $\mathcal{P}^{(1)}$ is a partition of the vertex set X which refines \mathcal{Q} . We refer to the cells of $\mathcal{P}^{(1)}$ as *clusters* of \mathcal{P} , so each cluster is a subset of some X_i , and every vertex of X lies in some cluster of \mathcal{P} . We say that \mathcal{P} is *vertex-equitable* if every cluster of \mathcal{P} has equal size. Also, for any \mathcal{Q} -partite set $S \subseteq X$ with $|S| \leq k$, we write $\text{Cell}(S)$ to denote the cell of \mathcal{P} which contains S .

We say that \mathcal{P} is a *partition k -complex on X* if it is a partition k -system on X with the additional property that for any edges $S, S' \in K(X)_A$ with $\text{Cell}(S) = \text{Cell}(S')$ and any subset $B \subseteq A$ we have $\text{Cell}(S \cap X_B) = \text{Cell}(S' \cap X_B)$. That is, if two sets lie in the same cell of \mathcal{P} , then their subsets of any given index also lie in the same cell of \mathcal{P} . To illustrate this definition, consider the following example of a partition 3-complex, where we slightly abuse notation in subscripts by writing, for example, \mathcal{P}_{12} rather than $\mathcal{P}_{\{1,2\}}$. Take $X = X_1 \cup X_2 \cup X_3$, and let the vertex classes X_1, X_2 and X_3 also be the clusters of \mathcal{P} (but bear in mind it is also possible for each vertex class to be partitioned into several clusters). Then the partition $\mathcal{P}^{(1)}$ is simply the partition of X into the clusters X_1, X_2 and X_3 . Next, \mathcal{P}_{12} is a partition of the set of all pairs $\{x_1, x_2\}$ with $x_1 \in X_1$ and $x_2 \in X_2$. That is, the cells C_{12}^i of \mathcal{P}_{12} are edge-disjoint bipartite graphs with vertex classes X_1 and X_2 , whose union is the complete bipartite graph on X_1 and X_2 . Similarly, the cells C_{13}^j of \mathcal{P}_{13} are bipartite graphs with vertex classes X_1 and

X_3 , and the cells C_{23}^ℓ of \mathcal{P}_{23} are bipartite graphs with vertex classes X_2 and X_3 . Now, for any choice of cells C_{12}^i , C_{13}^j and C_{23}^ℓ from \mathcal{P}_{12} , \mathcal{P}_{13} and \mathcal{P}_{23} respectively, the union of these cells is a tripartite graph; let $\Delta_{ij\ell}$ be the set of triangles in this tripartite graph. Observe that these sets $\Delta_{ij\ell}$ partition the set of all triples $\{x_1, x_2, x_3\}$ with $x_1 \in X_1$, $x_2 \in X_2$ and $x_3 \in X_3$; indeed any triple $\{x_1, x_2, x_3\}$ appears only in the $\Delta_{ij\ell}$ such that $\{x_1, x_2\} \in C_{12}^i$, $\{x_1, x_3\} \in C_{13}^j$ and $\{x_2, x_3\} \in C_{23}^\ell$. Finally, \mathcal{P}_{123} is also a partition of the set of all triples $\{x_1, x_2, x_3\}$ with $x_1 \in X_1$, $x_2 \in X_2$ and $x_3 \in X_3$; the requirement that \mathcal{P} is a partition k -complex requires that \mathcal{P}_{123} is a refinement of the partition into sets $\Delta_{ij\ell}$.

Suppose that \mathcal{P} is a partition k -complex on X . For any \mathcal{Q} -partite set $e \in \binom{X}{k}$, define $\mathcal{P}(e) := \bigcup_{e' \subseteq e} \text{Cell}(e')$. Then the fact that \mathcal{P} is a partition k -complex implies that $\mathcal{P}(e)$ is a k -partite k -complex with vertex classes X_j for $j \in i(e)$. Loosely speaking, the Regular Approximation Lemma will provide us with a partition k -complex so that all these k -complexes $\mathcal{P}(e)$ are regular complexes (as defined in the next section). Now suppose instead that \mathcal{P} is a partition $(k-1)$ -complex on X , and recall that X is partitioned into r parts X_1, \dots, X_r . Then for any $A \in \binom{[r]}{k}$ the cells of \mathcal{P} naturally generate a partition $\hat{\mathcal{P}}_A$ of the edges of $K(X)_A$. Indeed, we say that edges S and S' in $K(X)_A$ are *weakly equivalent* if $\text{Cell}(S_B) = \text{Cell}(S'_B)$ for any $B \subsetneq A$. This defines an equivalence relation on $K(X)_A$; we take the equivalence classes of this relation to be the parts of $\hat{\mathcal{P}}_A$. We can then extend \mathcal{P} to a partition k -complex $\hat{\mathcal{P}}$ on X by adding the partitions $\hat{\mathcal{P}}_A$ for $A \in \binom{[r]}{k}$ to \mathcal{P} . That is, for any $A \subseteq [r]$ with $|A| < k$ the cells of $\hat{\mathcal{P}}_A$ are the cells of \mathcal{P} of index A , and for any $A \in \binom{[r]}{k}$ the cells of $\hat{\mathcal{P}}_A$ are the equivalence classes of the weak equivalence relation on $K(X)_A$. We refer to $\hat{\mathcal{P}}$ as the *partition generated from \mathcal{P} by weak equivalence*. In particular, if \mathcal{P} is a -bounded, then \mathcal{P}_A has at most a cells for each $A \in \binom{[r]}{k-1}$, so $\hat{\mathcal{P}}$ is a^k -bounded. In a similar manner, for any \mathcal{Q} -partite k -graph G on X we can generate a partition k -complex $G[\hat{\mathcal{P}}]$ on X from $\hat{\mathcal{P}}$ by refining the partitions $\hat{\mathcal{P}}_A$ for each $A \in \binom{[r]}{k}$. Indeed, for each such A and each cell C of $\hat{\mathcal{P}}_A$ we have two cells of $G[\hat{\mathcal{P}}]_A$, namely $G \cap C$ and $C \setminus G$, whilst for any $A \in \binom{[r]}{\leq k-1}$, the cells of $G[\hat{\mathcal{P}}]_A$ are the same as those of $\hat{\mathcal{P}}_A$.

5.3. Hypergraph regularity. We now have all of the notation that we need to explain the notion of a regular complex and state the Regular Approximation Lemma we shall use. The concept of regularity with which we shall work was first introduced in the k -uniform case by Rödl and Skokan [34], but we shall consider it in the form used by Rödl and Schacht [32, 33].

Roughly speaking, an r -partite k -complex J is ε -regular if whenever we restrict J to those edges supported by a large subcomplex of $J \setminus J_-$ (that is, J minus its ‘top layer’), the resulting k -complex has similar densities to J . To demonstrate this, we shall first consider graphs (*i.e.* 2-graphs). If G is a bipartite graph with vertex classes V_1 and V_2 , then the standard definition of ε -regularity of G is that for any $V'_1 \subseteq V_1$ and $V'_2 \subseteq V_2$ with $|V'_1| > \varepsilon|V_1|$ and $|V'_2| > \varepsilon|V_2|$ we have $d(G[V'_1 \cup V'_2]) = d(G) \pm \varepsilon$. However, the definition of regularity which we generalise to hypergraphs is subtly different. Indeed, we say that G is ε -regular if for any $V'_1 \subseteq V_1$ and $V'_2 \subseteq V_2$ with $|V'_1||V'_2| > \varepsilon|V_1||V_2|$ we have $d(G[V'_1 \cup V'_2]) = d(G) \pm \varepsilon$ (note that this is equivalent to the previous definition in the sense that ε -regularity in the former implies ε -regularity in the latter, whilst ε^2 -regularity in the latter implies ε -regularity in the former). Now consider the 2-partite 2-complex J with edge set $\{\emptyset\} \cup \{\{v\} : v \in V(G)\} \cup G$, so the ‘layers’ of J are $\{\emptyset\}$, $\{\{v\} : v \in V(G)\}$ and G . Then saying that G is ε -regular (under the latter definition) is equivalent to saying that J is ε -regular under the following definition: J is ε -regular if, for any subcomplex $L \subseteq J$ with $|L_{\{1,2\}}^*| \geq \varepsilon|J_{\{1,2\}}^*|$, we have $|J_{\{1,2\}} \cap L_{\{1,2\}}^*|/|L_{\{1,2\}}^*| = d_{\{1,2\}}(J) \pm \varepsilon$.

Indeed, using the correspondence $V'_j = \{v \in V_j : \{v\} \in L_{\{j\}}\}$ for $j \in \{1, 2\}$ we find that the two definitions are equivalent, since then $|L_{\{1,2\}}^*| = |V'_1||V'_2|$ and $|J_{\{1,2\}}^*| = |V_1||V_2|$.

In general, let \mathcal{Q} partition a set X into r parts X_1, \dots, X_r , and let J be a \mathcal{Q} -partite k -complex. Then we generalise the definition above as follows: for any $A \in \binom{[r]}{\leq k}$ we say that J is ε -regular at A if for any subcomplex $L \subseteq J$ with $|L_A^*| \geq \varepsilon|J_A^*|$ we have

$$\frac{|J_A \cap L_A^*|}{|L_A^*|} = d_A(J) \pm \varepsilon.$$

We say J is ε -regular if J is ε -regular at A for every $A \in \binom{[r]}{\leq k}$. Now suppose that \mathcal{P} is a partition k -complex on X . Recall that for any \mathcal{Q} -partite set $e \in \binom{X}{k}$ the partition k -complex \mathcal{P} naturally yields a k -partite k -complex $\mathcal{P}(e)$ with vertex classes X_j for $j \in i(e)$; we say that \mathcal{P} is ε -regular if $\mathcal{P}(e)$ is ε -regular for any \mathcal{Q} -partite set $e \in \binom{X}{k}$.

Let G and H be r -partite k -graphs with common vertex classes X_1, \dots, X_r , and let $X := X_1 \cup \dots \cup X_r$. Then we say that G and H are ξ -close if $|G_A \triangle H_A| < \xi|K_A(X)|$ for every $A \in \binom{[r]}{\leq k}$. The Regular Approximation Lemma states that for any r -partite k -graph H there is an r -partite k -graph G on $V(H)$ (with the same vertex partition as H) and a partition $(k-1)$ -complex \mathcal{P} on $V(H)$ such that G is ξ -close to H and the partition k -complex $G[\hat{\mathcal{P}}]$ is ε -regular. This will suffice for our purposes as we shall avoid using any edge of $G \setminus H$ whilst working with G , so any edge we do use will be an edge of H . There are other regularity lemmas for k -graphs which give information on H itself (see [9, 34]) but the regular complexes yielded by these are not sufficiently dense to apply the blow-up lemma (see [13, Section 3] for further discussion of this point). The next theorem is the Regular Approximation Lemma; this is a slight restatement of a result of Rödl and Schacht (Theorem 14 of [32]).

Theorem 5.1 (Regular Approximation Lemma, [32]). *Suppose that integers n, a, r, k and reals ε, ξ satisfy $1/n \ll \varepsilon \ll 1/a \ll \xi, 1/r, 1/k$ and that $a!r$ divides n . Let \mathcal{Q} partition a set X of n vertices into r parts of equal size, and let H be a \mathcal{Q} -partite k -graph on X . Then there is an a -bounded ε -regular vertex-equitable partition $(k-1)$ -complex \mathcal{P} on X and a \mathcal{Q} -partite k -graph G with vertex set X such that G is ξ -close to H and $G[\hat{\mathcal{P}}]$ is ε -regular.*

One useful property of regularity is that if G is regular and dense, then the restriction of G to any not-too-small subsets of its vertex classes is also regular and dense. The following lemma (a weakened version of Theorem 6.18 in [13]) states this more precisely.

Lemma 5.2 (Regular restriction, [13]). *Suppose that $1/n \ll \varepsilon \ll d, c, 1/k$. Let J be an ε -regular k -partite k -complex with vertex classes X_1, \dots, X_k such that $d(J_{[k]}) \geq c$, $d_{[k]}(J) \geq d$ and $|X_j| \geq n$ for each $j \in [k]$. Also, for each $j \in [k]$, let $X'_j \subseteq X_j$ have $|X'_j| \geq \varepsilon^{1/2k}|X_j|$, and let $J' := J[X'_1 \cup \dots \cup X'_k]$. Then J' is $\sqrt{\varepsilon}$ -regular, $d(J'_{[k]}) \geq c/2$ and $d_{[k]}(J') \geq d/2$.*

5.4. Robustly universal complexes. Another vital tool in the proofs of Lemmas 3.1 and 3.2 is the recent hypergraph blow-up lemma of Keevash [13]. This states that if an r -partite k -complex J is ‘super-regular’ (a stronger property than regularity), then J contains a copy of any r -partite k -complex L with the same vertex classes and small maximum vertex degree. Another result in [13] shows that any regular and dense r -partite k -complex J can be made super-regular by the deletion of a few vertices from each vertex class. However, the notion of hypergraph super-regularity is very technical, so we shall avoid these technicalities through the related notion of ‘robust universality’, also from [13]. Roughly speaking, we say that an r -partite k -complex J' is robustly universal if even after the deletion of many vertices of J' , the resulting complex J has the property that one can find in J a copy of any

r -partite k -complex L with the same vertex classes as J and small maximum vertex degree. The next definition states this property formally; for this we make the following definitions. Let R be a k -complex on vertex set $[r]$, and let \mathcal{Q} partition a set X into parts X_1, \dots, X_r . Then a \mathcal{Q} -partite k -complex J on X is R -indexed if every edge $e \in J$ has $i(e) \in R$ (recall that $i(e)$ denotes the index of e). Also, for any $S \in R_+$, any $j \in S$ and $v \in X_j$, we write $J_S(v)$ for the $(k-1)$ -partite $(k-1)$ -complex with vertex set $\bigcup_{j \in S \setminus \{i\}} X_j$ and whose edges are those $(k-1)$ -tuples e' of vertices such that $e' \cup \{v\} \in J_S$.

Definition 5.3 (Robustly universal complexes, [13]). *Suppose that R is a k -complex on vertex set $[r]$, and that J' is an r -partite k -complex with vertex classes V'_1, \dots, V'_r such that $|J'_{\{i\}}| = |V'_i|$ for each $i \in [r]$. Then we say that*

- *J' is D -universal on R if for any R -indexed r -partite k -complex L with vertex classes U_1, \dots, U_r such that $|U_j| \leq |V'_j|$ for all $j \in [r]$ and $\Delta_{\text{vertex}}(L) \leq D$, there is a copy of L in J' in which the vertices of U_j correspond to the vertices of V'_j .*
- *J' is η -robustly D -universal on R if for any sets $V_j \subseteq V'_j$ such that $|V_j| \geq \eta|V'_j|$ for any $j \in [r]$ and $|J_S(v)| \geq \eta|J'_S(v)|$ for any $S \in R_+$ and $v \in V_S$, where $J = J'[\bigcup_{j \in [r]} V_j]$, the r -partite k -complex J is D -universal on R .*

In the case where R has k vertices and is formed by the downwards closure of a single edge, we omit ‘on R ’ and write simply ‘ D -universal’ or ‘ η -robustly D -universal’.

Note that the definition of robust universality given here is weaker than that from [13] in two ways. Firstly, the definition there allows R to be a so-called ‘multicomplex’, allowing us to distinguish between edges of J with the same index. Secondly, the definition in [13] also permits us to choose for a small number of vertices $v \in V(L)$ a small ‘target set’ into which v is to be embedded; an additional parameter c_0 governs how small these ‘target sets’ can be. However, we do not need either of these strengthenings.

Clearly robust universality is a very strong property, and so the main difficulty in the use of robust universality lies in obtaining robustly universal complexes in the first place. For this purpose we have the following theorem, which is a weakened version of Theorem 6.32 in [13] (to correspond to our weakened definition of robustly universal complexes). It states that if J is a regular k -complex which is dense on edges of R , and Z is a k -graph which has few edges in common with J_+ , then we may delete a small number of vertices from each vertex class of J so that the subcomplex of $J \setminus Z$ induced by the remaining vertices is robustly universal on R . Our use of the blow-up lemma is therefore concealed in this theorem, which we have slightly restated from the form in [13] in that the statements in (i) apply to $J' \setminus Z$ rather than to J' . The proof of this theorem in [13] in fact gives this altered result; alternatively, it can be derived by first deleting vertices of J which lie in atypically few edges of J_+ or in atypically many edges of Z , and then applying the form of the theorem stated in [13] (although the deletion step here is redundant, since these atypical vertices are deleted in the proof of this theorem in [13]).

Theorem 5.4 ([13]). *Suppose that*

$$1/n \ll 1/r' \ll \varepsilon \ll d^* \ll d_a \ll \nu \ll d, \eta, 1/k, 1/D, 1/C, 1/D_R,$$

and that $r \leq r'$. Let R be a k -complex on $[r]$ with $\Delta_{\text{vertex}}(R) \leq D_R$, and let J be an r -partite k -complex with vertex classes V_1, \dots, V_r , such that $n \leq |J_{\{j\}}| = |V_j| \leq Cn$ for every $j \in [r]$. Also let Z be a k -graph on $V(J)$, and suppose that

- (a) *J is ε -regular,*
- (b) *$d_S(J) \geq d$ and $d(J_S) \geq d_a$ for any $S \in R_+$,*

(c) $|Z \cap J_S| \leq \nu |J_S|$ for any $S \in R_-$.

Then we can delete at most $2\nu^{1/3}|V_j|$ vertices from each set V_j to obtain subsets V'_j so that, writing $V' = V'_1 \cup \dots \cup V'_r$ and $J' = J[V']$, we have

- (i) $d((J' \setminus Z)_S) > d^*$ and $|(J' \setminus Z)_S(v)| > d^*|(J' \setminus Z)_S|/|V'_j|$ for every $S \in R_-$, $j \in S$ and $v \in V'_j$, and
- (ii) $J' \setminus Z$ is η -robustly D -universal on R .

Having obtained a robustly universal k -partite k -complex J , we will then delete further vertices of J , and we wish these deletions to preserve the property that J is universal. The following proposition allows us to do this without difficulty; we shall only delete vertices which do not lie in the sets X_j .

Proposition 5.5. *Suppose that $1/n \ll d^* \ll \eta \ll \beta, 1/D, 1/k$. Let J be a k -partite k -complex with vertex classes W_1, \dots, W_k which is η -robustly D -universal and which satisfies $d(J_{[k]}) > d^*$ and $|J_{[k]}(v)| > d^*|J_{[k]}|/|W_j|$ for every $j \in [k]$ and $v \in W_j$. Suppose also that $\beta n \leq s_j \leq |W_j| \leq n$ for each $j \in [k]$ and some integers s_j . For each $j \in [k]$ choose a subset $X_j \subseteq W_j$ of size s_j uniformly at random and independently of all other choices. Then with probability $1 - o(1)$ we have the property that for any sets Y_j with $X_j \subseteq Y_j \subseteq W_j$ for each $j \in [k]$, the induced k -complex $J[\bigcup_{j \in [k]} Y_j]$ is D -universal.*

Proof. Observe that for any such sets Y_j we have $|Y_j| \geq |X_j| = s_j \geq \beta n \geq \eta|W_j|$ for any $j \in [k]$, and that $|J_{[k]}(v)| \geq |J_{[k]}(X \cup \{v\})(v)|$ for every $v \in Y$, where we define $Y := \bigcup_{j \in [k]} Y_j$ and $X = \bigcup_{j \in [k]} X_j$. So by definition of a η -robustly D -universal complex it suffices to show that with probability $1 - o(1)$ we have the property that $|J_{[k]}(X \cup \{v\})(v)| \geq \eta|J_{[k]}(v)|$ for every $v \in W := \bigcup_{j \in [k]} W_j$. In fact, Lemma 4.4 of [15], which was proved by a straightforward application of Azuma's inequality, states that for any $v \in W$ this inequality holds with probability at least $1 - 1/n^2$, so taking a union bound over all vertices of W proves the result. \square

5.5. The reduced k -graph. In this section we introduce the idea of the *reduced k -graph*, for which we make use of the k -graph Φ defined in Section 4. Recall that Φ has vertex set $[k+1]$, and has two edges, $\{1, \dots, k\}$ and $\{2, \dots, k+1\}$. Also recall that the vertices 1 and $k+1$ are the end vertices of Φ , and the vertices $2, \dots, k$ are the central vertices of Φ . Our definition of the reduced k -graph R will enable us, given a copy of Φ in R , to find a $(k+1)$ -partite k -complex which is universal on this copy of Φ . The next proposition shows that within this k -complex we can find many copies of $\Phi(m)$, the m -fold blowup of Φ , which is the $(k+1)$ -partite k -graph with vertex classes L_1, \dots, L_{k+1} of size m and whose edges are all k -tuples of vertices whose index is an edge of Φ . Copies of $\Phi(m)$ are particularly useful since we have flexibility over how a k -partite k -graph K can be embedded within $\Phi(m)$: we can embed $k-1$ of the vertex classes of K in the central vertex classes of $\Phi(m)$ (that is, those vertex classes corresponding to central vertices of Φ), and then the vertices of the remaining vertex class of K can be distributed as we choose among the two end vertex classes of $\Phi(m)$ (that is, those vertex classes corresponding to end vertices of Φ).

Proposition 5.6. *Let J be a $(k+1)$ -partite k -complex with vertex classes X_1, \dots, X_{k+1} which is D -universal on Φ , where $D \geq 2^{(k+1)m}$, and suppose that $|X_i| \geq n$ for each $i \in [k+1]$. Then there are at least $\lfloor n/m \rfloor$ vertex-disjoint copies of $\Phi(m)$ in $J_{[k+1]}$ whose end vertex classes lie in X_1 and X_{k+1} , and whose central vertex classes lie in X_2, \dots, X_k .*

Proof. Let L be the $(k+1)$ -partite k -complex formed by the downwards closure of $\lfloor n/m \rfloor$ vertex-disjoint copies of $\Phi(m)$; equivalently, L consists of $\lfloor n/m \rfloor$ vertex-disjoint copies of $\Phi(m)^\leq$. Since each copy of $\Phi(m)$ has $(k+1)m$ vertices we have $\Delta_{\text{vertex}}(L) \leq 2^{(k+1)m}$. Together with the fact that J is D -universal on Φ , it follows that J contains a copy of L in which the end vertex classes of each copy of $\Phi(m)^\leq$ lie in X_1 and X_{k+1} and whose central vertex classes lie in X_2, \dots, X_k . Then $L_\leq \subseteq J_\leq$ consists of the desired copies of $\Phi(m)$. \square

For notational simplicity, for the rest of this section we work within the following setup.

Setup 5.7. Fix integers n, a, r, D and k and constants $\varepsilon, d^*, \xi, \nu, \mu, c, \eta, \theta$ and γ with

$$1/n \ll \varepsilon \ll d^* \ll 1/a \ll 1/r, \xi \ll \nu \ll \mu \ll c, \eta \ll \theta \ll \gamma, 1/D, 1/k.$$

Let X be a set of n vertices, and let \mathcal{Q} be a partition of X into r parts T_1, \dots, T_r of equal size. Let \mathcal{P} be an a -bounded ε -regular vertex-equitable partition $(k-1)$ -complex on X such that the partition $\mathcal{P}^{(1)}$ of X into clusters X_1, \dots, X_m refines \mathcal{Q} . Assume that the number of clusters m satisfies $r \leq m \leq ar$, and let $n_1 = n/m$ be the common size of each cluster. Finally let G and Z be \mathcal{Q} -partite k -graphs on X , such that the partition k -complex $G[\hat{\mathcal{P}}]$ is ε -regular.

We can now give our definition of the reduced k -graph R . Similarly as in previous applications of hypergraph regularity, R has vertices corresponding to the clusters of \mathcal{P} , and edges corresponding to k -tuples of clusters which support many edges of G and few edges of Z . However, we also add a third condition, which we will use for Lemma 5.11 to show that any edge of R can be extended to a copy of Φ in R whose corresponding vertex classes support many copies of Φ in G .

Definition 5.8 (Reduced k -graph, Φ -dense, Z -sparse). Under Setup 5.7, the reduced k -graph of G and Z (with parameters c and ν) is the k -graph R on vertex set $[m]$ in which vertex i corresponds to the cluster X_i , and where $e \in \binom{[m]}{k}$ is an edge of R if

- (i) $|G[\bigcup_{i \in e} X_i]| \geq cn_1^k$,
- (ii) $|Z[\bigcup_{i \in e} X_i]| \leq \nu n_1^k$, and
- (iii) for any $e' \in \binom{[m]}{k-1}$ there are at most $\nu^2 mn_1^k$ edges of Z which intersect all of the clusters X_i with $i \in e'$.

Furthermore, for any $i, j \in [m]$ and $S \in \binom{[m]}{k-1}$, we say that the triple (i, S, j) is Φ -dense if there are at least $c^2 n_1^{k+1}$ copies of Φ in G which have an end vertex in each of X_i and X_j and a central vertex in X_ℓ for each $\ell \in S$, and we say that (i, S, j) is Z -sparse if each of $Z_{S \cup \{i\}}$ and $Z_{S \cup \{j\}}$ contains at most νn_1^k edges.

Note that under Setup 5.7, the partition \mathcal{Q} of X naturally induces a partition of $[m] = V(R)$ into r parts of equal size; we denote this partition by \mathcal{Q}_R . So i and j are in the same part of \mathcal{Q}_R if and only if the clusters X_i and X_j are subsets of the same part of \mathcal{Q} . The next lemma shows that within any Φ -dense and Z -sparse triple we can obtain a k -complex which is D -universal on Φ , to which we can gainfully apply Proposition 5.6.

Lemma 5.9. Adopt Setup 5.7, and suppose that sets $A, B \in \binom{[m]}{k}$ satisfy $|A \cap B| = k-1$. Let i and j be the elements of $A \setminus B$ and $B \setminus A$ respectively and suppose that the triple $(i, A \cap B, j)$ is Φ -dense and Z -sparse. Suppose also that we have subsets $Y_\ell \subseteq X_\ell$ with $|Y_\ell| \geq \eta n_1$ for each $\ell \in A \cup B$. Then there exist subsets $W_\ell \subseteq Y_\ell$ with $|W_\ell| \geq (1-\mu)|Y_\ell|$ for each $\ell \in A \cup B$ and a $(k+1)$ -partite k -complex J whose vertex classes are W_ℓ for $\ell \in A \cup B$ such that $J_\leq \subseteq G \setminus Z$ and J is D -universal on Φ (where we here consider A and B to be the edges of Φ , so i and j are the ends of Φ).

Proof. Introduce new constants d_a, ν' and ν'' with $d^* \ll d_a \ll 1/a$ and $\nu \ll \nu' \ll \nu'' \ll \mu$. Recall that for each \mathcal{Q}_R -partite set $S \in \binom{[m]}{k}$, $\hat{\mathcal{P}}$ partitions the n_1^k edges of $K(X)_S$ into at most a^k cells. We call such a cell C a *good* cell if it satisfies

- (a) $|C| \geq c^2 n_1^k / 5a^k$,
- (b) $|C \cap G| \geq c^2 |C| / 5$, and
- (c) $|C \cap Z| \leq \nu^{1/2} |C|$;

otherwise, C is a *bad* cell. Consider the copies of Φ in G whose edges e and f have indices $i(e) = A$ and $i(f) = B$. Since $(i, A \cap B, j)$ is Φ -dense, there are at least $c^2 n_1^{k+1}$ such copies of Φ in G . We will show that at least one of these copies of Φ must have the property that both of its edges are contained in good cells of $\hat{\mathcal{P}}$. For this, first note that since $\hat{\mathcal{P}}$ partitions $K(X)_A$ into at most a^k cells, at most $c^2 n_1^k / 5$ edges of $K(X)_A$ lie in cells C which fail (a). Likewise, since $(i, A \cap B, j)$ is Z -sparse we have $|Z_A| \leq \nu n_1^k$, and so at most $\nu^{1/2} n_1^k$ edges of $K(X)_A$ lie in cells C which fail (c). Finally, the number of edges of G_A which lie in cells C which fail (b) is

$$\sum_C |C \cap G| < \sum_C c^2 |C| / 5 \leq c^2 n_1^k / 5,$$

where the sum is taken over all cells C of index A which fail (b). We deduce that at most $(c^2/5 + \nu^{1/2} + c^2/5) n_1^k \cdot n_1 < c^2 n_1^{k+1} / 2$ of the copies of Φ we counted have the edge of index A in a cell which fails (a), (b) or (c). The same argument shows that fewer than $c^2 n_1^{k+1} / 2$ of the copies of Φ we counted have the edge of index B in a cell which fails (a), (b) or (c).

We may therefore fix a copy of Φ in G , whose edges e and f have indices $i(e) = A$ and $i(f) = B$ respectively, such that $\text{Cell}(e)$ and $\text{Cell}(f)$ each satisfy (a), (b) and (c). Recall that $G[\hat{\mathcal{P}}](e)$ is defined to be the k -partite k -complex with vertex classes X_i for $i \in A$ and whose edge set is

$$(G_A \cap \text{Cell}(e)) \cup \bigcup_{e' \subsetneq e} \text{Cell}(e'),$$

and that $G[\hat{\mathcal{P}}](f)$ is defined similarly. We define a $(k+1)$ -partite k -complex J^1 with vertex classes X_i for $i \in A \cup B$ to have edge set

$$J^1 := G[\hat{\mathcal{P}}](e) \cup G[\hat{\mathcal{P}}](f).$$

So the ‘top level’ of J^1 consists of all edges of G in the same cell as either e or f , whilst the lower levels of J^1 are comprised of the cells of \mathcal{P} which lie ‘below’ these cells. The crucial observation is that since e and f are the edges of a copy of Φ in G , for any $e' \subseteq e$ and $f' \subseteq f$ with $i(e') = i(f')$ we have $e' = f'$, and so J^1 includes only one cell of this index. That is, $J^1[X_A] = G[\hat{\mathcal{P}}](e)$, and $J^1[X_B] = G[\hat{\mathcal{P}}](f)$. Since $G[\hat{\mathcal{P}}]$ is ε -regular, $G[\hat{\mathcal{P}}](e)$ and $G[\hat{\mathcal{P}}](f)$ are ε -regular, and so it follows from the previous observation that J^1 is ε -regular also. Furthermore, we have

$$d_A(J^1) = \frac{|J_A^1|}{|(J_A^1)^*|} = \frac{|G \cap \text{Cell}(e)|}{|\text{Cell}(e)|} \geq \frac{c^2}{5},$$

and similarly $d_B(J^1) \geq c^2/5$. Also

$$d(J_A^1) = \frac{|J_A^1|}{|K(X)_A|} = \frac{|G \cap \text{Cell}(e)|}{|\text{Cell}(e)|} \cdot \frac{|\text{Cell}(e)|}{n_1^k} \geq \frac{c^2}{5} \cdot \frac{c^2}{5a^k} > 2d_a,$$

and similarly $d(J_B^1) > 2d_a$. Finally, observe that

$$|Z \cap J_A^1| \leq |Z \cap \text{Cell}(e)| \leq \nu^{1/2} |\text{Cell}(e)| \leq \nu^{1/2} \frac{|G \cap \text{Cell}(e)|}{c^2/5} \leq \nu' |J_A^1|,$$

and similarly $|Z \cap J_B^1| \leq \nu' |J_B^1|$.

Let $Y := \bigcup_{\ell \in A \cup B} Y_\ell$, and define $J^2 := J^1[Y]$. So J^2 is a $(k+1)$ -partite k -complex with vertex classes Y_ℓ for $\ell \in A \cup B$. By Lemma 5.2 applied to $J^2[Y_A]$ and $J^2[Y_B]$ in turn, we find that J^2 is $\sqrt{\varepsilon}$ -regular, that $d_A(J^2), d_B(J^2) \geq c^2/10$, and that $d(J_A^2) \geq d(J_A^1)/2 \geq d_a$ and $d(J_B^2) \geq d(J_B^1)/2 \geq d_a$. In particular, the fact that $d(J_A^2) \geq d(J_A^1)/2$, together with our assumption that $|Y_\ell| \geq \eta |X_\ell|$ for each $\ell \in A$, implies that $|J_A^2| \geq \eta^k |J_A^1|/2$. So

$$|Z \cap J_A^2| \leq |Z \cap J_A^1| \leq \nu' |J_A^1| \leq \frac{2\nu' |J_A^2|}{\eta^k} \leq \nu'' |J_A^2|,$$

and similarly $|Z \cap J_B^2| \leq \nu'' |J_B^2|$. So we may apply Theorem 5.4 with J^2 , Φ and the sets Y_ℓ in place of J , R and the sets V_ℓ respectively, and with $\eta n_1, 1/\eta, \nu''$ and $c^2/10$ in place of n, C, ν and d respectively. This yields subsets $W_\ell \subseteq Y_\ell$ with $|W_\ell| \geq (1 - 2(\nu'')^{1/3}) |Y_\ell| \geq (1 - \mu) |Y_\ell|$ for each $\ell \in A \cup B$ such that, writing $J := J^2[\bigcup_{\ell \in A \cup B} W_\ell] \setminus Z$, we have that J is η -robustly D -universal on Φ (so in particular J is D -universal on Φ), and that $J_- \subseteq G \setminus Z$. \square

Note that the application of Theorem 5.4 at the end of the proof also yields the facts that $d(J_A) > d^*$ and $|J_A(v)| > d^* |J_A|/|W_j|$ for any $j \in A$ and $v \in W_j$. We do not need these facts when applying Lemma 5.9, but we do need the analogous results when applying the next lemma, whose proof is similar to but simpler than that of Lemma 5.9, so we omit it (a comparable result was also proved for a slightly different definition of reduced k -graph in [15, Section 5.1.5], by a similar argument).

Lemma 5.10. *Adopt Setup 5.7, let R be the reduced k -graph of G and Z , and let A be an edge of R . Then for any subsets $Y_i \subseteq X_i$ with $|Y_i| \geq \eta |X_i|$ for each $i \in A$, there exist subsets $W_i \subseteq Y_i$ with $|W_i| \geq (1 - \mu) |Y_i|$ for each $i \in A$ and a k -partite k -complex J with vertex classes W_i for $i \in A$ such that J is η -robustly D -universal, $J_- \subseteq G \setminus Z$, $d(J_A) \geq d^*$ and $|J_A(v)| > d^* |J_A|/|W_j|$ for every $j \in A$ and $v \in W_j$.*

Our final lemma shows if all \mathcal{Q} -partite $(k-1)$ -tuples have large degree in $G \cup Z$, then this degree condition is ‘almost’ inherited by the reduced k -graph R , in that almost all $(k-1)$ -tuples of R satisfy a comparable condition. Furthermore, we also find that any edge of R can be extended to many Φ -dense and Z -sparse triples. To prove this latter result we make use of the unusual condition (iii) in Definition 5.8; this is the purpose of that condition.

Lemma 5.11. *Adopt Setup 5.7, and suppose that every \mathcal{Q} -partite $(k-1)$ -tuple e of vertices of G has $\deg_{G \cup Z}(e) \geq \gamma n$, and also that $|Z| \leq \xi n^k$. Then*

- (i) *there are at most θm^{k-1} many $(k-1)$ -tuples $S' \in \binom{[m]}{k-1}$ with $\deg_R(S') < (\gamma - \theta)m$.*
- (ii) *Furthermore, for any edge $S \in R$ and any $i \in S$ there are at least $(\gamma - \theta)m$ choices for $j \in [m] \setminus S$ such that the triple $(i, S \setminus \{i\}, j)$ is Φ -dense and Z -sparse.*

Proof. Let \mathcal{S}' consist of all sets $S' \in \binom{[m]}{k-1}$ such that

- (a) S' is \mathcal{Q}_R -partite,
- (b) at most $\nu^2 m n_1^k$ edges of Z intersect all of the clusters X_ℓ with $\ell \in S'$, and
- (c) for any $S'' \in \binom{S'}{k-2}$ at most $\nu^3 m^2 n_1^k$ edges of Z intersect all of the clusters X_ℓ with $\ell \in S''$.

We will show that every $S' \in \mathcal{S}'$ has $\deg_R(S') \geq (\gamma - \theta)m$. To see this, fix some $S' \in \mathcal{S}'$, and let

$$\mathcal{S} := \{S' \cup \{i\} : i \in [m] \setminus S'\}.$$

Since S' is \mathcal{Q}_R -partite, any $(k-1)$ -tuple e' which consists of one vertex of X_ℓ for each $\ell \in S'$ has $\deg_{G \cup Z}(e') \geq \gamma n$ by assumption. Since $G \cup Z$ is \mathcal{Q} -partite, any edge $e \in G \cup Z$ with $e' \subseteq e$ must have $e \in (G \cup Z)_S$ for some $S \in \mathcal{S}$, and so we conclude that there are at least

$n_1^{k-1}\gamma n = \gamma n_1^k m$ edges $e \in G \cup Z$ whose index $i(e)$ is a member of \mathcal{S} . By (b), at most $\nu^2 m n_1^k$ of these edges lie in Z , and furthermore at most $c n_1^k m$ of these edges lie in G_S for some $S \in \mathcal{S}$ with $|G_S| < c n_1^k$. This leaves at least $(\gamma - c - \nu^2) n_1^k m$ edges which lie in G_S for some $S \in \mathcal{S}$ with $|G_S| \geq c n_1^k$. Since $|G_S| \leq n_1^k$ for any $S \in \mathcal{S}$, we conclude that there are at least $(\gamma - c - \nu^2) m$ sets $S \in \mathcal{S}$ such that $|G_S| \geq c n_1^k$. Now observe that there can be at most νm sets $S \in \mathcal{S}$ such that $|Z_S| > \nu n_1^k$. Indeed, if there were more, then taking the union of these Z_S we would obtain more than $\nu^2 m n_1^k$ edges of Z which meet X_ℓ for each $\ell \in S'$, contradicting (b). Similarly, there can be at most $2k\nu m$ sets $S \in \mathcal{S}$ for which some subset $T' \in \binom{S}{k-1}$ has the property that least $\nu^2 n_1^k m$ edges of Z meet X_ℓ for every $\ell \in T'$. Indeed, if there were more, then some $S'' \in \binom{S'}{k-2}$ would be a subset of at least $2\nu m$ of the subsets T' , implying that more than $\nu^3 m^2 n_1^k$ edges of Z meet X_ℓ for every $\ell \in S''$, contradicting (c). We conclude that there are at least $(\gamma - c - \nu^2 - \nu - 2k\nu) m \geq (\gamma - \theta) m$ sets $S \in \mathcal{S}$ such that $|G_S| \geq c n_1^k$, $|Z_S| \leq \nu n_1^k$, and no subset $T' \in \binom{S}{k-1}$ has the property that at least $\nu^2 n_1^k m$ edges of Z meet X_ℓ for every $\ell \in T'$; any S with these three properties is an edge of R . So we do indeed have $\deg_R(S') \geq (\gamma - \theta) m$.

It remains to prove that there are at most θm^{k-1} sets $S' \in \binom{[m]}{k-1}$ such that $S' \notin \mathcal{S}'$, that is, which fail either (a), (b) or (c). For this, first note that at most m^{k-1}/r sets $S' \in \binom{[m]}{k-1}$ are not \mathcal{Q}_R -partite. Writing N for the number of sets $S' \in \binom{[m]}{k-1}$ such that more than $\nu^2 m n_1^k$ edges of Z meet X_ℓ for each $\ell \in S'$, the fact that $|Z| \leq \xi n^k$ implies that $N \nu^2 m n_1^k \leq k \xi n^k$, so $N \leq k \xi m^{k-1}/\nu^2$. Finally, write N' for the number of sets $S' \in \binom{[m]}{k-1}$ such that some subset $S'' \in \binom{S'}{k-2}$ has the property that there are more than $\nu^3 m^2 n_1^k$ edges of Z which meet X_ℓ for every $\ell \in S''$. The number of sets S'' with this property is then at least N'/m , so we obtain $(N'/m) \nu^3 m^2 n_1^k \leq k^2 \xi n^k$, that is, $N' \leq k^2 \xi m^{k-1}/\nu^3$. We conclude that, as claimed, the number of sets $S' \in \binom{[m]}{k-1}$ such that $S' \notin \mathcal{S}'$ is at most

$$m^{k-1}/r + N + N' \leq m^{k-1}/r + k \xi m^{k-1}/\nu^2 + k^2 \xi m^{k-1}/\nu^3 \leq \theta m^{k-1}.$$

For the ‘furthermore’ part, fix any $S \in R$ and $i \in S$, and write $S' := S \setminus \{i\}$. Since $S \in R$ we know that there are at most $\nu^2 m n_1^k$ edges of Z which meet X_ℓ for every $\ell \in S'$. So at most νn_1^{k-1} edges $e' \in K(X)_{S'}$ have $\deg_Z(e') \geq \nu n$. Now, for any edge $e \in G_S$ we have a $(k-1)$ -tuple $e' := e \setminus X_i \in K(X)_{S'}$; since G is \mathcal{Q} -partite our minimum codegree assumption implies that $\deg_{G \cup Z}(e') \geq \gamma n$. Each $(k-1)$ -tuple e' is formed in this way from at most n_1 edges of G_S , so we conclude that there are at least $|G_S| - \nu n_1^k$ edges $e \in G_S$ for which $\deg_G(e') \geq (\gamma - \nu) n$. Since at most $k n_1 \leq \nu n$ vertices lie in the sets X_ℓ for $\ell \in S$, there are at least $(|G_S| - \nu n_1^k)(\gamma - 2\nu) n$ copies of Φ in G whose edges have indices S and $S' \cup \{j\}$ for some $j \in [m] \setminus S$. Since for any $j \notin S$ at most $|G_S| n_1$ of these copies have a vertex in X_j , we conclude that the triple (i, S', j) is Φ -dense for at least

$$\frac{(|G_S| - \nu n_1^k)(\gamma - 2\nu) n - c^2 n_1^{k+1} m}{|G_S| n_1} \geq \frac{|G_S|(\gamma - 2\nu) m - (\gamma \nu + c^2) n_1^k m}{|G_S|} \geq \left(\gamma - \frac{\theta}{2} \right) m$$

choices of $j \in [m] \setminus S$, where we used the fact that $|G_S| \geq c n_1^k$ since $S \in R$. So to complete the proof it suffices to show that the triple (i, S', j) is Z -sparse for all but at most $\theta m/2$ choices of $j \in [m] \setminus S$. For this, recall that $|Z_S| \leq \nu n_1^k$ since S is an edge of R , so if (i, S', j) is not Z -sparse then $|Z_{S' \cup \{j\}}| > \nu n_1^k$. Furthermore, since $S \in R$ there are at most $\nu^2 m n_1^k$ edges of Z which meet X_ℓ for every $\ell \in S'$. So, writing N'' for the number of choices of j for which the triple is not Z -sparse, we have $N'' \nu n_1^k \leq \nu^2 m n_1^k$, and so $N'' \leq \nu m \leq \theta m/2$, as required. \square

5.6. Degree sequences and irreducibility. Let J be a k -complex. Then the *degree sequence* of J is the sequence $\delta(J) = (\delta_0(J), \delta_1(J), \dots, \delta_{k-1}(J))$, where for any $i \in [k]$ we define

$$\delta_{i-1}(J) := \min_{e \in J_{i-1}} \deg_{J_i}(e).$$

So every edge $e \in J_{i-1}$ is a subset of at least $\delta_{i-1}(J)$ edges of J_i , or in other words there are at least $\delta_{i-1}(J)$ vertices $v \in V(J)$ such that $e \cup \{v\} \in J$. Inequalities between degree sequences should always be interpreted pointwise. Note also that we only defined the minimum codegree $\delta(H)$ for k -graphs H , and the degree sequence $\delta(J)$ for k -complexes J , so there should be no confusion.

The following lemma states that if almost all $(k-1)$ -tuples of vertices of a k -graph H have high degree, then we can find a k -complex J which covers almost all of the vertices of H , such that J has a useful degree sequence and the ‘top level’ J_+ of J is a subgraph of H . The k -partite form of this lemma was given by Keevash, Knox and Mycroft [14, Lemma 7.3] with a straightforward proof. The proof of the form given below is identical except for the simplification of not having to handle multiple vertex classes, so we omit it (this form is also implicit in [16]).

Lemma 5.12. *Suppose that $1/m \ll \theta \ll \beta, 1/k$, and let H be a k -graph on a vertex set V of size m in which at most θm^{k-1} sets $S \in \binom{V}{k-1}$ have $\deg_H(S) \leq D$. Then there exists a k -complex J with $V(J) \subseteq V$ such that $J_+ \subseteq H$, $m' := |V(J)| \geq (1 - \sqrt{\theta})m$ and $\delta(J) \geq (m', (1 - \beta)m', \dots, (1 - \beta)m', D - \beta m')$.*

Now let H be a k -graph on n vertices which admits a perfect matching M . Using the terminology of Keevash and Mycroft [16] we say that H is (C, L) -irreducible on M if for any $u, v \in V(H)$ with $u \neq v$ there exist multisets S and T of edges of H and M respectively, so that $|S|, |T| \leq L$ and, counting with multiplicity, for some $c \leq C$ the vertex u appears in precisely c more edges of S than of T , the vertex v appears in precisely c more edges of T than of S , and every other vertex of H appears equally often in S as in T . The next lemma, a special case of a result of Keevash and Mycroft [16, Lemma 5.6], gives a sufficient degree sequence condition on a k -complex J for J_+ to be irreducible on a perfect matching in J_+ .

Lemma 5.13. *Suppose that $1/m \ll 1/C, 1/L \ll \alpha, 1/k$, and let J be a k -complex on m vertices with $\delta(J) \geq (m, (k-1)m/k + \alpha m, (k-2)/k + \alpha m, \dots, m/k + \alpha m)$ such that J_+ admits a perfect matching M . Then J_+ is (C, L) -irreducible on M .*

6. INGREDIENTS OF THE PROOF

6.1. Partitioning clusters into lopsided groups. As described in Section 4, we will find an almost-perfect packing of the reduced k -graph \mathcal{R} with a particular k -partite k -graph $\mathcal{A}_{p,q}^k$. The k -graph $\mathcal{A}_{p,q}^k$ which we use is defined as follows. The vertex set $V(\mathcal{A}_{p,q}^k)$ is the union of disjoint sets A_1, \dots, A_{q-p} and B , where $|A_j| = k-1$ for each $j \in [q-p]$ and $|B| = p(k-1)$. Then any k -tuple of the form $\{x\} \cup A_j$ with $j \in [q-p]$ and $x \in B$ is an edge of $\mathcal{A}_{p,q}^k$ (see Figure 4 for an illustration). In particular we have $|V(\mathcal{A}_{p,q}^k)| = q(k-1)$.

The next lemma shows that if G is a k -graph on m vertices in which almost all sets of $k-1$ vertices have degree slightly greater than pm/q , then G contains an almost-perfect $\mathcal{A}_{p,q}^k$ -packing, that is, one which covers almost all vertices of G . We will apply this result with the reduced k -graph \mathcal{R} in place of G and with p and q chosen so that $p/q \approx \sigma(K)$. The degree condition needed will then follow from Lemma 5.11 and our assumption in Lemma 3.1 that $\delta(H) \geq \sigma(K)n + \alpha n$. This lemma was previously proved for $k=3, p=1, q=4$ by Kühn




FIGURE 4. The k -graph $\mathcal{A}_{p,q}^k$ in the case $k = 3, p = 1, q = 4$; the edges between B and A_1 are shown, and there are similar edges between B and A_2 and between B and A_3 , giving six edges in total.

and Osthus [21] and then for $p = 1, q = 2k - 2$ by Keevash, Kühn, Mycroft and Osthus [15]; the proof given here is essentially identical, but is included for completeness.

Lemma 6.1. *Suppose that $1/m \ll \theta \ll \psi \ll 1/q, 1/p, 1/k$ and that G is a k -graph on vertex set $[m]$ such that $\deg_G(S) > (\frac{p}{q} + \theta)m$ for all but at most θm^{k-1} sets $S \in \binom{[m]}{k-1}$. Then G admits an $\mathcal{A}_{p,q}^k$ -packing \mathcal{F} such that $|V(\mathcal{F})| \geq (1 - \psi)m$ and $G[V(\mathcal{F})]$ is connected (where $V(\mathcal{F})$ denotes the set of vertices covered by \mathcal{F}).*

Proof. Let \mathcal{F} be a maximal $\mathcal{A}_{p,q}^k$ -packing in G , and let $X := V(G) \setminus V(\mathcal{F})$. We will show that $|X| \leq \psi m/2$; to do this, we suppose for a contradiction that $|X| > \psi m/2$. For any $(k-1)$ -tuple S of vertices of G , we write $N(S)$ to denote the set $\{v \in V(G) : S \cup \{v\} \in G\}$ of neighbours of S , so $|N(S)| = \deg_G(S)$. We also write $\deg(S)$, $N_X(S)$ and $\deg_X(S)$ for $\deg_G(S)$, $N(S) \cap X$ and $|N_X(S)|$ respectively. Note that since $\theta \ll \psi$ we can greedily form a collection of at least $2\theta m$ disjoint $(k-1)$ -tuples $S \in \binom{X}{k-1}$ which each satisfy $\deg(S) \geq pm/q + \theta m$.

Suppose first that for some $r \geq \theta m$ there exist disjoint sets $S_1, \dots, S_r \in \binom{X}{k-1}$ such that $\deg_X(S_i) \geq \theta m/2$ for any $i \in [r]$. In this case, we count the pairs (i, B) such that $i \in [r]$ and $B \subseteq N_X(S_i)$ has size $p(k-1)$. By our choice of the sets S_1, \dots, S_r , the number of such pairs is at least

$$r \binom{\theta m/2}{p(k-1)} \geq \theta m \binom{\theta m/2}{p(k-1)} \geq (q-p) \binom{m}{p(k-1)} \geq (q-p) \binom{|X|}{p(k-1)}.$$

So there must be some set $B \in \binom{X}{p(k-1)}$ which lies in at least $q-p$ such pairs; the corresponding $q-p$ sets S_i together with this set B form a copy of $\mathcal{A}_{p,q}^k$ contained in $G[X]$, contradicting the maximality of \mathcal{F} .

Since there are at least $2\theta m$ disjoint $(k-1)$ -tuples $S \in \binom{X}{k-1}$ which each satisfy $\deg(S) \geq pm/q + \theta m$, it follows that we may choose a family of $r \geq \theta m$ subsets $S_1, \dots, S_r \in \binom{X}{k-1}$ such that each S_i satisfies $\deg(S_i) \geq pm/q + \theta m$ and $\deg_X(S_i) < \theta m/2$. Having fixed this family, we say that a copy $\mathcal{A} \in \mathcal{F}$ is *good* for S_j if $|V(\mathcal{A}) \cap N(S_j)| > p(k-1)$. Note that each set S_j has at least $pm/q + \theta m/2$ neighbours in $V(\mathcal{F})$, and at most

$$|\mathcal{F}|p(k-1) \leq \frac{p(k-1)m}{|V(\mathcal{A}_{p,q}^k)|} = pm/q$$

of these neighbours lie in members of \mathcal{F} which are not good for S_j . So the number of copies $\mathcal{A} \in \mathcal{F}$ which are good for S_j is at least $\theta m/2 |V(\mathcal{A}_{p,q}^k)| = \theta m/2q(k-1)$.

We now count the number of pairs (j, \mathcal{T}) where $j \in [r]$ and $\mathcal{T} \subseteq \mathcal{F}$ consists of $p(k-1)$ copies $\mathcal{A} \in \mathcal{F}$, each of which is good for S_j . By the above calculation, this number is at least

$$r \binom{\theta m/2q(k-1)}{p(k-1)} \geq \theta m \binom{\theta m/2q(k-1)}{p(k-1)} \geq \sqrt{m} \binom{m}{p(k-1)} \geq \sqrt{m} \binom{|\mathcal{F}|}{p(k-1)}.$$

We can therefore choose a collection \mathcal{T} of $p(k-1)$ copies $\mathcal{A} \in \mathcal{F}$ and a subset $R \subseteq [r]$ of size $|R| \geq \sqrt{m}$ such that \mathcal{A} is good for S_j for any $j \in R$ and $\mathcal{A} \in \mathcal{T}$. This means that for each $j \in R$ and each $\mathcal{A} \in \mathcal{T}$ we may choose a subset $L_j^{\mathcal{A}} \subseteq N(S_j) \cap V(\mathcal{A})$ of size $p(k-1) + 1$. Having done so, the fact that $|R| \geq \sqrt{m}$ implies that we may choose a subset $R' \subseteq R$ of size $(p(k-1) + 1)(q-p)$ so that for any fixed $\mathcal{A} \in \mathcal{T}$, $L_j^{\mathcal{A}}$ is the same set for every $j \in R'$. We write $L^{\mathcal{A}}$ for this common value of $L_j^{\mathcal{A}}$.

Arbitrarily partition R' into $p(k-1) + 1$ sets $R'_1, \dots, R'_{p(k-1)+1}$ of size $(q-p)$, and label the vertices of each $L^{\mathcal{A}}$ as $\{v_1^{\mathcal{A}}, v_2^{\mathcal{A}}, \dots, v_{p(k-1)+1}^{\mathcal{A}}\}$. Then for each $s \in [p(k-1) + 1]$, the sets S_j for $j \in R'_s$ and the set $\{v_s^{\mathcal{A}} : \mathcal{A} \in \mathcal{T}\}$ together form a copy of $\mathcal{A}_{p,q}^k$. This produces $p(k-1) + 1$ vertex-disjoint copies of $\mathcal{A}_{p,q}^k$ which are contained in $X \cup V(\mathcal{T})$, so we may enlarge \mathcal{F} by replacing the members of \mathcal{T} with these copies, giving another contradiction.

This proves that $|X| \leq \psi m/2$, so \mathcal{F} covers at least $(1 - \psi/2)m$ vertices of G . Note that $G[\mathcal{A}]$ is connected for any $\mathcal{A} \in \mathcal{F}$. Let $\mathcal{F}' \subseteq \mathcal{F}$ be of maximum size such that $G[V(\mathcal{F}')] is connected, and suppose for a contradiction that $|V(\mathcal{F}')| < (1 - \psi)m$, so $|V(\mathcal{F}) \setminus V(\mathcal{F}')| > \psi m/2$. We first observe that some vertex of $V(\mathcal{F})$ must lie in some $(k-1)$ -tuple $S \in \binom{V(\mathcal{F})}{k-1}$ with $\deg_G(S) \geq pm/q + \theta m$, and so has at least $pm/2q$ neighbours in $V(\mathcal{F})$, so by maximality of \mathcal{F}' we have $|V(\mathcal{F}')| \geq pm/2q$. Therefore, the number of sets $S \in \binom{V(\mathcal{F})}{k-1}$ which contain a vertex $x \in V(\mathcal{F}) \setminus V(\mathcal{F}')$ and a vertex $y \in V(\mathcal{F}')$ is at least$

$$\frac{1}{(k-1)!} \cdot \frac{\psi m}{2} \cdot \frac{pm}{2q} \cdot ((1 - \psi/2)m)^{k-3} > \theta m^{k-1}.$$

It follows that some such S has degree at least $pm/q > \psi m/2$, and so can be extended to an edge of $G[V(\mathcal{F})]$. But then the member of \mathcal{F} containing x can be added to \mathcal{F}' to give a larger subpacking $\mathcal{F}'' \subseteq \mathcal{F}$ such that $G[V(\mathcal{F}'')] is connected, a contradiction. This proves that $|V(\mathcal{F}')| \geq (1 - \psi)m$, so \mathcal{F}' is the desired $\mathcal{A}_{p,q}^k$ -packing. $\square$$

Having obtained an almost-perfect $\mathcal{A}_{p,q}^k$ -packing in the reduced k -graph \mathcal{R} , we will proceed to partition the clusters corresponding to copies of $\mathcal{A}_{p,q}^k$, and then to rearrange the parts obtained into groups of k subclusters which support regular and dense complexes. This partition is effected in the following way.

Lemma 6.2. *Suppose that $pk \leq q$, and that for each vertex $u \in V(\mathcal{A}_{p,q}^k)$ we have a set V_u of n vertices such that the sets V_u are pairwise-disjoint. Let $V = \bigcup_{u \in V(\mathcal{A}_{p,q}^k)} V_u$, and suppose also that $(q-p)p(k-1)$ divides n . Then we may partition V into sets X_j^i with $j \in [k]$ and $i \in [(q-p)p(k-1)]$ such that*

- (i) $|X_1^i| = \frac{n}{q} \sum_{j \in [k]} |X_j^i|$ for each i ,
- (ii) $n/(q-p) = |X_1^i| \leq |X_2^i| = |X_3^i| = \dots = |X_k^i|$ for each i ,
- (iii) for each i and j there exists $f(i, j) \in V(\mathcal{A}_{p,q}^k)$ so that $X_j^i \subseteq V_{f(i,j)}$, and
- (iv) for each fixed i the set $\{f(i, j) : j \in [k]\}$ is an edge of $\mathcal{A}_{p,q}^k$.




FIGURE 5. An illustration of the division of clusters implemented in Lemma 6.2 for the case $k = 3, p = 1, q = 4$. The dashed lines join subclusters which form part of the same group.

Proof. Let A_1, \dots, A_{q-p} and B be as in the definition of the k -graph $\mathcal{A}_{p,q}^k$. So these sets are pairwise-disjoint and their union is $V(\mathcal{A}_{p,q}^k)$; also, $|B| = p(k-1)$ and $|A_a| = k-1$ for each $a \in [q-p]$. Arbitrarily order the vertices of each of these sets, and for $i \in [k-1]$ and $a \in [q-p]$ write $u(i, A_a)$ for the i th vertex of A_a , and similarly for $j \in [p(k-1)]$ write $v(j, B)$ for the j th vertex of B . Next, for every $j \in [q-p]$ and every $u \in A_j$, partition the set V_u into $p(k-1)$ parts $V_u^1, \dots, V_u^{p(k-1)}$ of equal size. Similarly, for each $v \in B$ partition V_v into $q-p$ parts V_v^1, \dots, V_v^{q-p} of equal size. Then for each $a \in [q-p]$ and $b \in [p(k-1)]$ define

$$X_1^{a,b} = V_{v(b,B)}^a \text{ and } X_j^{a,b} = V_{u(j-1,A_a)}^b \text{ for } 2 \leq j \leq k.$$

Relabelling these sets (that is, replacing the superscript (a, b) by an integer in $[(q-p)p(k-1)]$) gives the desired sets.

Property (iii) is immediate from the construction, and since any set of the form $A_a \cup \{v\}$ with $v \in B$ is an edge of $\mathcal{A}_{p,q}^k$, (iv) is satisfied also. Finally, observe that for each i , $|X_1^i| = n/(q-p)$ and $|X_2^i| = \dots = |X_k^i| = n/p(k-1)$. So our assumption that $pk \leq q$ implies that $|X_1^i| \leq |X_2^i|$, proving (ii), and

$$\frac{p}{q} \cdot \sum_{j \in [k]} |X_j^i| = \frac{p}{q} \left(\frac{n}{q-p} + (k-1) \cdot \frac{n}{p(k-1)} \right) = \frac{n}{q-p} = |X_1^i|,$$

so (i) holds also. \square

In the proof of Lemma 3.2 we will find a matching $M_{\mathcal{R}}$ in the reduced k -graph \mathcal{R} , rather than an $\mathcal{A}_{p,q}^k$ -packing. Observe for this that $\mathcal{A}_{1,k}^k$ contains a perfect matching, so it suffices to find an $\mathcal{A}_{1,k}^k$ -packing in \mathcal{R} , which we can do by applying Lemma 6.1 with $p = 1$ and $q = k$. However, in the proof of Lemma 3.2 we also require an additional assumption, namely that the restriction of \mathcal{R} to any large submatching $M'_{\mathcal{R}} \subseteq M_{\mathcal{R}}$ of this matching is irreducible on $M'_{\mathcal{R}}$. The following corollary states that we can do this.

Corollary 6.3. *Suppose that $1/m \ll \theta \ll \psi \ll 1/C, 1/L \ll \alpha, 1/k$, and that G is a k -graph on vertex set $[m]$ such that $\deg_G(S) > (1/k + \alpha)m$ for all but at most θm^{k-1} sets*

$S \in \binom{[m]}{k-1}$. Then G admits a matching M with $|V(M)| \geq (1 - \psi)m$ such that $G[V(M')]$ is (C, L) -irreducible on M' for any $M' \subseteq M$ with $|M'| \geq (1 - \alpha/2)|M|$.

Proof. Introduce a new constant β with $\theta \ll \beta \ll \psi$. By Lemma 5.12 there exists a k -complex J with $V(J) \subseteq [m]$ such that $J_- \subseteq G$, $m_1 := |V(J)| \geq (1 - \sqrt{\theta})m$, and $\delta(J) \geq (m_1, (1 - \beta)m_1, \dots, (1 - \beta)m_1, (1/k + \alpha - \beta)m_1)$. Then at least $(1 - \beta)^{k-2}m_1^{k-1}/(k-1)! \geq (1 - k\beta)\binom{m_1}{k-1}$ many $(k-1)$ -tuples $S \in \binom{V(J)}{k-1}$ are edges of J_{k-1} , and so have $\deg_{J_-}(S) \geq (1/k + \alpha - \beta)m_1 \geq (1/k + k\beta)m_1$. So we can apply Lemma 6.1 to J_- with $p = 1$ and $q = k$, and with $k\beta$ and $\psi/2$ in place of θ and ψ respectively. Since $\mathcal{A}_{1,k}^k$ admits a perfect matching, this yields a matching M in $J_- \subseteq G$ with $|V(M)| \geq (1 - \psi/2)m_1 \geq (1 - \psi)m$. Now fix any $M' \subseteq M$ of size $|M'| \geq (1 - \alpha/2)|M|$, and define $m' := |V(M')|$, so $m_1 - m' \leq 2\alpha m_1/3$. It follows that $\delta(J[V(M')]) \geq (m', (1 - \alpha)m_1, \dots, (1 - \alpha)m_1, (1/k + \alpha/4)m_1)$, and so $J_-[V(M')]$ is (C, L) -irreducible on M' by Lemma 5.13 (with $\alpha/4$ in place of α). Since $J_- \subseteq G$ it follows that $G[V(M')]$ is (C, L) -irreducible on M' . \square

6.2. Incorporating exceptional vertices. We will need to be able to remove a small number of ‘bad’ vertices of H . Our strategy here will be to find a copy of K which contains the vertex to be removed, and to delete that copy of K from H . This copy of K will ultimately form part of the perfect K -packing of H which we construct. The next lemma allows us to do this by demonstrating that any vertex of a k -graph H with high codegree must lie in some copy of K in H .

Lemma 6.4. *Suppose that $1/n \ll \alpha, 1/b$. Let K be a k -partite k -graph on b vertices, and let H be a k -graph on n vertices with $\delta(H) \geq \alpha n$. Then for any vertex $u \in V(H)$ there is a copy of K in H which contains u .*

Proof. Partition the vertices of H into parts V_1 and V_2 by assigning u to V_1 and randomly assigning each other vertex of H to V_1 with probability $1/2$ and V_2 otherwise, where these assignments are independent for each vertex. Let $H' \subseteq H$ be the k -graph on vertex set $V(H)$ whose edge set is

$$\{e \in H : |e \cap V_1| = 1 \text{ and } \{u\} \cup (e \cap V_2) \in K\}.$$

So an edge of H is an edge of H' if it has precisely $k-1$ vertices in V_2 and these $k-1$ vertices together with u also form an edge of H . It suffices to show that for some outcome of our random selection the k -graph H' has at least $2^{-k}\alpha^2\binom{n}{k}$ edges. Indeed, by Theorem 3.7, H' must then contain a copy of $\mathcal{B}(K)$ (recall from Definition 3.5 that this is the complete k -partite k -graph with k vertex classes each of size b). Together with u , this gives a subgraph of H which contains as a subgraph a copy of K containing u .

Now, if we choose vertices x_1, \dots, x_{k-1} in turn to form an edge $\{u, x_1, \dots, x_{k-1}\}$ of H , then we have $n-j$ choices for x_j for $1 \leq j \leq k-2$ and at least $\delta(H) \geq \alpha n$ choices for x_{k-1} . Since this process will count each edge $(k-1)!$ times, we find that u lies in at least $\alpha\binom{n}{k-1}$ edges of H . For any such edge $\{u, x_1, \dots, x_{k-1}\}$ there are at least $\delta(H) \geq \alpha n$ choices of y such that $\{y, x_1, \dots, x_{k-1}\}$ is an edge $e \in H$. Each such edge may be formed by up to k different choices of x_1, \dots, x_{k-1} , so we find that there are at least $\alpha^2\binom{n}{k}$ edges $e \in H$ for which there is some $y \in e$ such that $\{u\} \cup e \setminus \{y\}$ is an edge of H . For each such edge, the probability that y is assigned to V_1 and all vertices of $e \setminus \{y\}$ are assigned to V_2 is at least 2^{-k} . So the expected number of edges e with this form whose vertices are assigned in this way is at least $2^{-k}\alpha^2\binom{n}{k}$, and every such edge is an edge of H' . There must therefore be some outcome of our random partition of $V(H)$ for which H' has at least this many edges, as required. \square

6.3. Ensuring divisibility of subcluster sizes. As described in Section 4, a key step in the proof of Lemma 3.1 is to delete a K -packing in H such that, following these deletions, the size of each subcluster is divisible by $bk \gcd(K)$ (recall that b is the order of the k -graph H). This allows us to complete the proof by finding a perfect K -packing in each of our robustly universal k -partite k -graphs $G^i \setminus Z^i$. In this section we prove Lemma 6.7, which states that we can indeed do this. We begin with the following lemma.

Lemma 6.5. *Let G be a t -partite k -graph with vertex classes X_1, \dots, X_t . Fix any integer d , and let d' be a factor of d such that d' divides $|X_j|$ for any $j \in [t]$. Also fix a k -graph K on b vertices, and suppose that \mathcal{S} is a graph on vertex set $[t]$ such that*

- (i) *for any connected component C of \mathcal{S} , $\sum_{j \in V(C)} |X_j|$ is divisible by d , and*
- (ii) *for any edge $uv \in \mathcal{S}$ there are at least bdt^2 vertex-disjoint copies K' of K in G such that for each $j \in [t]$ we have*

$$|V(K') \cap X_j| \equiv \begin{cases} -d' \pmod{d} & \text{if } j = u \\ d' \pmod{d} & \text{if } j = v \\ 0 \pmod{d} & \text{otherwise} \end{cases}$$

Then G contains an K -packing M of size at most dt^2 so that d divides $|X_j \setminus V(M)|$ for any $j \in [t]$.

Proof. We prove the lemma by repeatedly choosing an K -packing in G , deleting its vertices from G , and adding its members to M (which is initially taken to be empty). This ensures that M will indeed be an K -packing in G . After each deletion we continue to write X_j for the vertices in X_j which were not deleted, and G for the k -graph which remains (that is, the restriction of G to the undeleted vertices). We will also ensure that each deletion preserves the properties that d divides $\sum_{j \in V(C)} |X_j|$ for any component C of \mathcal{S} and that d' divides $|X_j|$ for any $j \in [t]$.

The deletion step is as follows: suppose that there is some $u \in [t]$ such that $|X_u| \not\equiv 0 \pmod{d}$, and let $x \in [d-1]$ satisfy $xd' \equiv |X_u| \pmod{d}$ (this is possible since d' is a factor of d which divides $|X_u|$). Let C be the component of \mathcal{S} containing u ; since d divides $\sum_{j \in V(C)} |X_j|$ by (i) there must be some $v \in V(C)$ such that $v \neq u$ and $|X_v| \not\equiv 0 \pmod{d}$. Also, since C is a component of \mathcal{S} we may choose a path P from u to v in \mathcal{S} . Let $u = w_0, w_1, \dots, w_p = v$ be the vertices of P (in order), so $p \leq t$. Now, for each $\ell \in [p]$, $w_{\ell-1}w_\ell$ is an edge of \mathcal{S} , so by (ii) we may choose x copies of K in G such that the intersection of each copy of K with the vertex class X_j has size equal to d' modulo d if $j = w_{\ell-1}$, equal to $-d'$ modulo d if $j = w_\ell$, and equal to 0 modulo d otherwise. We do this so that the chosen copies of K are pairwise vertex-disjoint (we shall see shortly that we can simply choose copies of K greedily to ensure this). Delete the vertices of each chosen copy of K from G and add these copies to M . The effect of these deletions is to reduce $|X_u|$ by xd' modulo d , to increase $|X_v|$ by xd' modulo d , and to leave the size of each other vertex class unchanged modulo d . So we now have $|X_u| \equiv 0 \pmod{d}$, that is, the number of vertex classes X_j with $|X_j| \equiv 0 \pmod{d}$ has increased by at least one.

We repeat the deletion step until $|X_j| \equiv 0 \pmod{d}$ for every $j \in [t]$; the previous observation shows that this must occur after at most t steps. Since at each step we deleted $px < td$ copies of K , the K -packing M obtained at termination has size less than dt^2 , as required. The same argument shows that it is possible to choose copies of K as claimed, since at any point the fewer than dt^2 previously-deleted copies of K can intersect fewer than bdt^2 members of a family of pairwise vertex-disjoint copies of K . \square

If $\gcd(K) = 1$, then Lemma 6.5 is in fact sufficient for our purposes (i.e. to delete a K -packing in H so that the number of remaining vertices in each subcluster is divisible by $bk\gcd(K)$). Indeed, in this case we first arbitrarily delete a small number of copies of K so that bk divides the total number of remaining vertices. We then choose s large enough such that $\mathcal{U}_s(K)$ is defined (see Definition 3.5) and so that k divides s , and apply the lemma with the adjacency graph $\text{Adj}(\mathcal{R}')$, $\mathcal{U}_s(K)$, bk and the subclusters V_j^i in place of \mathcal{S}, K, d and the sets X_j respectively, and with $d' = 1$. The graph $\text{Adj}(\mathcal{R}')$ is connected since \mathcal{R}' is connected, and so has only one connected component, so condition (i) of Lemma 6.5 holds by our initial deletions, and condition (ii) Lemma 6.5 follows from Lemma 5.10 and the fact that $\mathcal{U}_s(K)$ has one vertex class of size $bs - 1$ and one of size $bs + 1$, whilst all other vertex classes have size bs . So we obtain a $\mathcal{U}_s(K)$ -packing M in H whose deletion leaves all subclusters with size divisible by bk ; since $\mathcal{U}_s(K)$ admits a perfect K -packing this gives a K -packing as required (this argument is given in more detail in the proof of Lemma 6.7).

However, if $\gcd(K) \geq 2$ then the situation is somewhat more complicated, and we in fact make two applications of Lemma 6.5; once with $\text{Adj}(\mathcal{R}')$ in place of \mathcal{S} as described above, and another with \mathcal{S} being the graph \mathcal{S}' described in the proof outline in Section 4, whose edges indicate that the corresponding subclusters were taken from clusters which form ends of a Φ -dense and Z -sparse triple. In the latter application, condition (ii) of Lemma 6.5 follows as a consequence of Lemma 5.9. However, it is more problematic to ensure that condition (i) is satisfied, as \mathcal{S}' may have multiple connected components. The key here is that, as outlined in Section 4, \mathcal{S}' must have fewer than p components, where p is the least prime factor of $\gcd(K)$. That is, every prime factor of $\gcd(K)$ is strictly greater than r , the number of components of \mathcal{S}' . The next lemma shows that this fact allows us to choose edges of \mathcal{R}' whose index vectors with respect to the partition of $V(\mathcal{R}') = V(\mathcal{S}')$ into components of \mathcal{S}' sum to any chosen ‘target vector’ \mathbf{v} . In the proof of Lemma 6.7 we use these edges of \mathcal{R}' to chose copies of K for deletion to ensure that condition (i) of Lemma 6.5 is satisfied. Note that Lemma 6.6 would not hold if d had some prime factor p equal to r , as demonstrated by the k -graph constructed in Proposition 2.2 for this value of p . So Lemma 6.6 is the point in the proof of Lemma 3.1 at which the minimum codegree condition $\delta(H) \geq n/p + \alpha n$ is necessary (in the case $\gcd(K) > 1$).

For this lemma we use a slightly different definition of index vector. Let \mathcal{P} be a partition of a set X into parts X_1, \dots, X_r ; then for a given d and any $S \subseteq X$ we now define the index vector $\mathbf{i}_{\mathcal{P}}^d(S)$ of S with respect to \mathcal{P} to be the vector in \mathbb{Z}_d^r whose j -th coordinate is $|S \cap X_j|$ modulo d (whereas our previous definition had r in place of d). Again, we sometimes omit the subscript \mathcal{P} and write simply $\mathbf{i}^d(S)$ if \mathcal{P} is clear from the context. Recall that \mathbf{u}_j denotes the j th unit vector of \mathbb{Z}_d^r , i.e. the vector whose j th coordinate is equal to one with all other coordinates equal to zero.

Lemma 6.6. *Suppose that k, d and r are positive integers such that $k \geq 3$ and every prime factor of d is strictly greater than r . Let H be a k -graph on vertex set X , and let \mathcal{P} partition X into parts X_1, \dots, X_r . Also suppose that for any $j_1, \dots, j_{k-1} \in [r]$ there is an edge $\{u_1, \dots, u_k\} \in H$ with $u_i \in X_{j_i}$ for every $i \in [k-1]$. Then for any $\mathbf{v} = (v_1, \dots, v_r) \in \mathbb{Z}_d^r$ such that d divides $\sum_{i=1}^r v_i$ there exist a set S of at most $(r+1)^2$ edges of H and integers a_e for $e \in S$ such that $0 \leq a_e \leq d-1$ for each $e \in S$, d divides $\sum_{e \in S} a_e$ and, working in \mathbb{Z}_d^r , we have $\sum_{e \in S} a_e \mathbf{i}^d(e) = \mathbf{v}$.*

Note that we do not assume that H is \mathcal{P} -partite. Also, throughout the proof of Lemma 6.6 we work within \mathbb{Z}_d^r for all vector calculations (so all equalities of vectors should be interpreted in this context).

Proof. We fix k and d , and proceed by induction on r ; for this note that the fact that every prime factor of d is strictly greater than r implies that every prime factor of d is strictly greater than r' for any $r' \leq r$. For $r = 1$ the lemma is trivial since we must have $\mathbf{v} = (0)$. So fix $r \geq 2$, and assume that the lemma holds with $r - 1$ in place of r .

We claim that for some distinct $i, j \in [r]$ the vector $\mathbf{u}_i - \mathbf{u}_j$ can be written as an integer combination of at most r members of $\mathcal{D} := \{\mathbf{i}^d(e) - \mathbf{i}^d(e') : e, e' \in H\}$ (that is, there are integers c_1, \dots, c_p and vectors $\mathbf{x}_1, \dots, \mathbf{x}_p \in \mathcal{D}$ such that $p \leq r$ and $\mathbf{u}_i - \mathbf{u}_j = \sum_{i \in [p]} c_i \mathbf{x}_i$; we don't place any other restrictions on the integers c_i). To see that this is true, suppose for a contradiction that the claim is false, and fix any distinct $i, j \in [r]$. Since $k \geq 3$, our assumption on H allows us to choose an edge of $e \in H$ which has at least two vertices in X_i . Similarly we may choose an edge $e' \in H$ such that $\mathbf{i}^d(e') = \mathbf{i}^d(e) - 2\mathbf{u}_i + \mathbf{u}_j + \mathbf{u}_\ell$ for some $\ell \in [r]$. Then $\mathbf{i}^d(e) - \mathbf{i}^d(e') = 2\mathbf{u}_i - \mathbf{u}_j - \mathbf{u}_\ell \in \mathcal{D}$. If $\ell = i$, then this gives $\mathbf{u}_i - \mathbf{u}_j \in \mathcal{D}$, giving a contradiction (since $\mathbf{u}_i - \mathbf{u}_j$ can then be expressed as an integer combination of a single member of \mathcal{D}). Similarly, if $\ell = j$, then we obtain $2\mathbf{u}_i - 2\mathbf{u}_j \in \mathcal{D}$. Since $r \geq 2$ we know that d is odd, so $d' := (d + 1)/2$ is an integer with $d'(2\mathbf{u}_i - 2\mathbf{u}_j) = \mathbf{u}_i - \mathbf{u}_j$, and so $\mathbf{u}_i - \mathbf{u}_j$ is an integer combination of a single member of \mathcal{D} , again giving a contradiction. So we must have $\ell \neq i, j$; since i and j were arbitrary this implies that for any distinct $i, j \in [r]$ there is some $\ell = \ell(i, j)$ which is distinct from i and j such that $\mathbf{x}_{i,j} := 2\mathbf{u}_i - \mathbf{u}_j - \mathbf{u}_\ell \in \mathcal{D}$. Fix any i and write $f(j) := \ell(i, j)$ for each $j \neq i$. Then for any $j \neq i$ we can write

$$\mathbf{x}_{i,j} - \mathbf{x}_{i,f(j)} = (2\mathbf{u}_i - \mathbf{u}_j - \mathbf{u}_{f(j)}) - (2\mathbf{u}_i - \mathbf{u}_{f(j)} - \mathbf{u}_{f(f(j))}) = \mathbf{u}_{f(f(j))} - \mathbf{u}_j.$$

This expresses $\mathbf{u}_j - \mathbf{u}_{f(f(j))}$ as an integer combination of $2 \leq r$ members of \mathcal{D} , giving another contradiction unless $f(f(j)) = j$ for any $j \neq i$. So we may assume that the family $\mathcal{F}_i := \{\{j, f(j)\} : j \in [r] \setminus \{i\}\}$ is a partition of $[r] \setminus \{i\}$ into $(r - 1)/2$ pairs (note in particular this implies that r is odd). Then write

$$\mathbf{y}_i := \sum_{\{j,\ell\} \in \mathcal{F}_i} \mathbf{x}_{i,j} = \sum_{\{j,\ell\} \in \mathcal{F}_i} 2\mathbf{u}_i - \mathbf{u}_j - \mathbf{u}_\ell = (r - 1)\mathbf{u}_i - \sum_{j \in [r] \setminus \{i\}} \mathbf{u}_j.$$

So \mathbf{y}_i can be written as an integer combination of at most $(r - 1)/2$ members of \mathcal{D} . Since r and d are coprime, we may fix integers λ, μ such that $\lambda r + \mu d = 1$, whereupon $\lambda(\mathbf{y}_1 - \mathbf{y}_2) = \lambda r \mathbf{u}_1 - \lambda r \mathbf{u}_2 = \mathbf{u}_1 - \mathbf{u}_2$ can be written as an integer combination of at most $r - 1$ members of \mathcal{D} , giving a final contradiction which completes the proof of the claim.

We may therefore assume without loss of generality that $\mathbf{u}_{r-1} - \mathbf{u}_r$ can be written as an integer combination of at most r elements of \mathcal{D} . That is, we may choose a set S_1 of at most $2r$ edges of H and integers m_e for $e \in S_1$ such that $\sum_{e \in S_1} m_e = 0$ and $\sum_{e \in S_1} m_e \mathbf{i}_P^d(e) = \mathbf{u}_{r-1} - \mathbf{u}_r$. Let $Y_j = X_j$ for each $j \in [r - 2]$, and let $Y_{r-1} = X_{r-1} \cup X_r$. Then H is a k -graph on vertex set $X = Y_1 \cup \dots \cup Y_{r-1}$ such that for any $j_1, \dots, j_{k-1} \in [r - 1]$ there is an edge $\{u_1, \dots, u_k\} \in H$ with $v_i \in Y_{j_i}$ for every $i \in [k - 1]$. Let \mathcal{Q} denote the partition of X into the parts Y_1, \dots, Y_{r-1} ; then by our induction hypothesis we may choose a set S_2 of at most r^2 edges of H and integers n_e for $e \in S$ such that d divides $\sum_{e \in S_2} n_e$ and $\sum_{e \in S_2} n_e \mathbf{i}_Q^d(e) = (v_1, \dots, v_{d-2}, v_{d-1} + v_d)$. The latter equation implies that $\sum_{e \in S_2} n_e \mathbf{i}_P^d(e) = (v_1, \dots, v_{d-2}, y, z)$ for some y and z with $y + z = v_{d-1} + v_d$ modulo d , and so

$$(v_{d-1} - y)(\mathbf{u}_{r-1} - \mathbf{u}_r) + \sum_{e \in S_2} n_e \mathbf{i}_P^d(e) = (v_1, \dots, v_{d-2}, v_{d-1}, v_d) = \mathbf{v}.$$

Let $S := S_1 \cup S_2$ and let integers $0 \leq a_e \leq d - 1$ satisfy $a_e \equiv (v_{d-1} - y)m_e + n_e \pmod{d}$ for each $e \in S$ (we take $m_e = 0$ for any $e \notin S_1$ and $n_e = 0$ for any $e \notin S_2$). Then S is a set of at most $r^2 + 2r \leq (r + 1)^2$ edges of H , and the equation above shows that $\sum_{e \in S} a_e \mathbf{i}_P^d(e) = \mathbf{v}$.

Finally,

$$\sum_{e \in S} a_e \equiv (v_{d-1} - y) \sum_{e \in S} m_e + \sum_{e \in S} n_e \equiv 0 + 0 \equiv 0 \pmod{d},$$

so d divides $\sum_{e \in S} a_e$, as required. \square

Finally, we can now give the full statement and proof of Lemma 6.7, showing that we can delete a K -packing in H so that, following these deletions, all subclusters have size divisible by $bk \gcd(K)$. We achieve this by deleting five vertex-disjoint K -packings in succession. The first deletion is simple and ensures that the total number of vertices is divisible by $b \gcd(K)$, whilst the second uses Lemma 6.6 to ensure that $\gcd(K)$ divides the total number of vertices in subclusters within any component of \mathcal{S} . The third then uses Lemma 6.5 to ensure that $\gcd(K)$ divides the size of each subcluster, and the fourth (again straightforward) maintains this property whilst also ensuring that $bk \gcd(K)$ divides the total number of vertices. Finally, our fifth deletion uses Lemma 6.5 again to ensure that $bk \gcd(K)$ divides the number of vertices within any subcluster.

Lemma 6.7. *Suppose that N, s, t, b and k are integers such that $1/N \ll 1/t, 1/s \ll 1/b, 1/k$. Let K be the complete k -partite k -graph with vertex class sizes b_1, \dots, b_k , where $b_1 + \dots + b_k = b$, and suppose that $\gcd(K)$ is defined, that s is divisible by $k \gcd(K)$ and that b_1 and $\gcd(K)$ are coprime. Next let G be a t -partite k -graph with vertex classes Y_1, \dots, Y_t , and suppose that b divides $|Y|$, where $Y = \bigcup_{i \in [t]} Y_i$. Finally suppose that \mathcal{R} is a connected k -graph on $[t]$, and \mathcal{S} is a graph on $[t]$ with r connected components C_1, \dots, C_r , such that the following properties hold.*

- (i) *For any edge $e \in \mathcal{R}$ there are more than N vertex-disjoint copies of $\mathcal{B}(b(s+1))$ in $G[\bigcup_{j \in e} Y_j]$,*
- (ii) *For any edge $uv \in \mathcal{S}$ there is a set $T \in \binom{[t] \setminus \{u, v\}}{k-1}$ such that $G[\bigcup_{j \in \{u, v\} \cup T} Y_j]$ contains more than N vertex-disjoint copies of $\Phi(b(s+1))$ whose end vertex classes lie in Y_u and Y_v and whose central vertex classes lie in the sets Y_j for $j \in T$.*
- (iii) *If $\gcd(K) > 1$, then r (the number of components of \mathcal{S}) is smaller than the least prime factor of $\gcd(K)$, and for any $i_1, \dots, i_{k-1} \in [r]$ there is some edge $e = \{u_1, \dots, u_k\}$ of \mathcal{R} such that $u_j \in V(C_{i_j})$ for each $j \in [k-1]$.*

Then G contains a K -packing M of size at most $N/2b$ such that $bk \gcd(K)$ divides $|Y_j \setminus V(M)|$ for every $j \in [t]$.

Proof. As in Lemma 6.5, we prove the lemma by repeatedly choosing some vertex-disjoint copies of K in G and deleting their vertices from G ; as there, we continue to write Y_j , Y and G for the sets and graph obtained following these deletions. We shall verify at the end of the proof that the K -packing M formed by all the deleted copies of K has size at most $N/2b$, so M covers at most $N/2$ vertices. With this in mind, we can always assume that (i) and (ii) provide at least $N/2$ copies of $\mathcal{B}(b(s+1))$ and $\Phi(b(s+1))$ of the given forms.

Our first step is to delete at most $\gcd(K)$ pairwise vertex-disjoint copies of K from G so that, following these deletions, we have that $b \gcd(K)$ divides $|Y|$. Since each copy of K has b vertices, and b divides $|Y|$, we can indeed achieve this by deleting at most $\gcd(K)$ pairwise vertex-disjoint copies of K from G ; by (i) these copies can be chosen from $G[\bigcup_{j \in e} Y_j]$ for an arbitrary edge $e \in \mathcal{R}$.

The next step is to delete at most $(r+1)^2 \gcd(K)$ copies of K from G so that $\gcd(K)$ divides $\sum_{u \in V(C_i)} |Y_u|$ for each component C_i of \mathcal{S} . If $\gcd(K) = 1$ then no deletions are necessary, whilst if $\gcd(K) > 1$ then we use Lemma 6.6. For each $i \in [r]$, write $V_i := \bigcup_{u \in V(C_i)} Y_u$, and define $v_i \in \{0, \dots, \gcd(K) - 1\}$ to be such that $v_i b_1 \equiv |V_i| \pmod{\gcd(K)}$

(since $\gcd(K)$ and b_1 are coprime a unique such v_i exists). Then since $\gcd(K)$ divides $|Y|$, we have $\sum_{i \in [r]} v_i b_1 \equiv |Y| \equiv 0$ modulo $\gcd(K)$, so $\sum_{i \in [r]} v_i$ is divisible by $\gcd(K)$. We may therefore apply Lemma 6.6 with $\mathcal{R}, \gcd(K)$ and the sets $V(C_i)$ in place of H, d and the sets X_i respectively to obtain at most $(r+1)^2$ edges $e_1, \dots, e_p \in \mathcal{R}$ and integers $a_1, \dots, a_p \in \{0, 1, \dots, \gcd(K) - 1\}$ so that $\gcd(K)$ divides $\sum_{j \in [p]} a_j$ and (working in $\mathbb{Z}_{\gcd(K)}^r$) we have

$$\sum_{j \in [p]} a_j \mathbf{i}_{\mathcal{Q}}^d(e_j) = (v_1, \dots, v_r),$$

where \mathcal{Q} denotes the partition of $[t]$ into parts $V(C_i)$ for $i \in [r]$. For each $j \in [p]$ by (i) we may choose a_j pairwise vertex-disjoint copies of $\mathcal{B}(b(s+1))$ in $G[\bigcup_{\ell \in e_j} Y_\ell]$, within which we can find a_j pairwise vertex-disjoint copies of K . Delete all of these copies of K from G . By definition of $\gcd(K)$, each vertex class of K has size b_1 modulo $\gcd(K)$. Furthermore, since G is t -partite, each of the deleted copies of K has one vertex class contained in Y_ℓ for each $\ell \in e_j$. So for any $i \in [r]$, the total number of vertices deleted from V_i is

$$b_1 \sum_{j \in [p]} a_j |e_j \cap V(C_i)| \equiv b_1 v_i \equiv |V_i| \pmod{\gcd(K)}.$$

So following these deletions we have that $\gcd(K)$ divides $|V_i| = \sum_{u \in V(C_i)} |Y_u|$ for each component C_i of \mathcal{S} . Furthermore, since in total $\sum_{j \in [p]} a_j$ copies of K were deleted, each with b vertices, our assumption that $\gcd(K)$ divides $\sum_{j \in [p]} a_j$ implies that the total number of vertices deleted is divisible by $b \gcd(K)$. So we still have that $b \gcd(K)$ divides $|Y|$ after these deletions.

We now delete at most $N/3b$ further copies of K from G so that $\gcd(K)$ divides $|Y_i|$ for every $i \in [t]$. For this we use Lemma 6.5 with $\mathcal{B}(b \gcd(K)), \gcd(K)$ and the sets Y_i in place of K, d and the sets X_i respectively, with G, \mathcal{S} and t playing the same role here as there, and with $d' = 1$. Then condition (i) of Lemma 6.5 is satisfied as a consequence of our last round of deletions. Also, by (ii), for any edge $uv \in \mathcal{S}$ we can choose $N/2 \geq (bk \gcd(K)) \gcd(K) t^2$ vertex-disjoint copies of $\Phi(b(s+1))$ in G whose end vertex classes lie in Y_u and Y_v , and whose central vertex classes are each a subset of some Y_j . Within each of these copies of $\Phi(b(s+1))$ we can find a copy of $\mathcal{B}(b \gcd(K))$ with one vertex in Y_v , $b \gcd(K) - 1 \equiv -1 \pmod{\gcd(K)}$ vertices in Y_u , and $b \gcd(K) \equiv 0 \pmod{\gcd(K)}$ vertices in each of the other vertex classes intersected by this copy of $\Phi(b(s+1))$. So condition (ii) of Lemma 6.5 is satisfied also, and so Lemma 6.5 yields a $\mathcal{B}(b \gcd(K))$ -packing M' in G of size at most $\gcd(K) t^2$ such that, deleting all vertices covered by M' from G , we find that $\gcd(K)$ divides $|Y_i|$ for every $i \in [t]$. Then, since $\mathcal{B}(b \gcd(K))$ has $kb \gcd(K)$ vertices, it remains the case that $b \gcd(K)$ divides $|Y|$ following these deletions. Recall that $\mathcal{B}(K) = \mathcal{B}(b)$, so $\mathcal{B}(b \gcd(K))$ admits a perfect $\mathcal{B}(K)$ -packing of size $\gcd(K)$, whilst $\mathcal{B}(K)$ admits a perfect K -packing of size k by Proposition 3.6. So there is a K -packing in G which covers the same vertices as M' , so we did indeed delete a K -packing of size at most $kt^2 \gcd(K)^2 \leq N/5b$ in this step.

Next we delete at most $k \gcd(K)$ further copies of K from G so that, following these deletions, we have that $bk \gcd(K)$ divides $|Y|$, as well as preserving the property that $\gcd(K)$ divides $|Y_i|$ for every $i \in [t]$. Since $b \gcd(K)$ divides $|Y|$, we can achieve the latter property by deleting $z \gcd(K)$ copies of K for some integer $0 \leq z \leq k-1$. So choose an arbitrary edge $e \in \mathcal{R}$, and use (i) to choose $z \gcd(K)$ vertex-disjoint copies of K in $G[\bigcup_{i \in e} Y_i]$; then for any $j \in [t]$ the number of vertices deleted from Y_j is equal to $z \gcd(K) b_1 \equiv 0$ modulo $\gcd(K)$, so $|Y_j|$ is still divisible by $\gcd(K)$ following these deletions.

Finally, we apply Lemma 6.5 again to delete a final set of at most $N/5b$ copies of K from G so that, following these deletions, $bk \gcd(K)$ divides $|Y_i|$ for every $i \in [t]$, giving

the desired K -packing M . We shall use the adjacency graph $\text{Adj}(\mathcal{R})$ in place of \mathcal{S} ; since \mathcal{R} is connected, $\text{Adj}(\mathcal{R})$ is connected also, and so condition (i) of Lemma 6.5 holds (with $bk \gcd(K)$ in place of d) as a consequence of our last round of deletions. Furthermore, for any edge uv of $\text{Adj}(\mathcal{R})$ there is an edge $e \in \mathcal{R}$ containing u and v . So by (i) there are at least $N/2 > (bsk)(bk \gcd(K))t^2$ vertex-disjoint copies of $\mathcal{B}(b(s+1))$ in $G[\bigcup_{\ell \in e} Y_\ell]$, each of which contains a copy of $\mathcal{U}_s(K)$ with $bs - \gcd(K)$ vertices in Y_u , $bs + \gcd(K)$ vertices in Y_v and bs vertices in Y_w for any $w \in e \setminus \{u, v\}$ (we can assume that $\mathcal{U}_s(K)$ is defined since we assumed that $1/s \ll 1/b, 1/k$). Since $bs \equiv 0$ modulo $bk \gcd(K)$, we may apply Lemma 6.5 with $\mathcal{U}_s(K)$, $\text{Adj}(\mathcal{R})$, $\gcd(K)$, $bk \gcd(K)$ and the sets Y_j in place of K , \mathcal{S} , d' , d and the sets X_j respectively, whereupon the requirement that $\gcd(K)$ divides $|Y_j|$ for every $j \in [t]$ is satisfied by our previous deletions. This gives a $\mathcal{U}_s(K)$ -packing M'' in G of size at most $bk \gcd(K)t^2$ in G such that, deleting all members of M'' from G , we find that $bk \gcd(K)$ divides $|Y_i|$ for every $i \in [t]$. Since $\mathcal{U}_s(K)$ admits a perfect K -packing of size ks by Proposition 3.6, we may treat M'' as being a K -packing in G of size at most $k^2 bs \gcd(K)t^2 \leq N/5b$, as required.

To complete the proof we must show that at most $N/2b$ copies of K were deleted in total. Indeed, we deleted at most $\gcd(K)$ copies in the first step, at most $\sum_{j \in [p]} a_j \leq (r+1)^2 \gcd(K)$ copies in the second step, at most $N/5b$ copies in the third step, at most $k \gcd(K)$ copies in the fourth step, and at most $N/5b$ copies in the final step, that is, fewer than $N/2b$ copies in total. \square

6.4. Ensuring equality of subcluster sizes. In Lemma 3.2, each vertex class of the k -partite k -graph K has equal size b_1 , so it is insufficient to delete a K -packing in H so that every cluster satisfies certain divisibility conditions. Instead, we must ensure that the clusters in each of our robustly universal k -partite k -graphs have equal size. In this section we prove Lemma 6.10, which gives sufficient conditions for this to be possible. The stronger minimum codegree condition on H will ensure that we can satisfy these conditions, and so apply Lemma 6.10 in the proof of Lemma 3.2 similarly as Lemma 6.7 is used in the proof of Lemma 3.1. However, the results of this section are stated in a more general form which we can also use in the proof of Theorem 1.4.

In this section we proceed under the following setup, in which the k -graphs G and \mathcal{R} , the directed graph \mathcal{S}^+ , base graph \mathcal{S} and clusters U_i play the roles described in Section 4. A *directed graph* D consists of a vertex set V and a set of edges E , where each edge is an ordered pair (u, v) with $u, v \in V$ and $u \neq v$. We write $u \rightarrow v$ to mean that $(u, v) \in E$. The *outdegree* $\deg^+(u)$ of a vertex $u \in V$ is the number of vertices $v \in V$ for which $u \rightarrow v$, and the *minimum outdegree* of D is $\delta^+(D) := \min_{u \in V} \deg^+(u)$. Finally, the *base graph* of D is the (undirected) graph G on V in which uv is an edge of G if either $u \rightarrow v$ or $v \rightarrow u$ in D .

Setup 6.8. Let \mathcal{R} be a k -graph with vertex set $[m]$ which admits a perfect matching $M_{\mathcal{R}}$, and for any $i \in [m]$ let $e(i)$ denote the edge of $M_{\mathcal{R}}$ which contains i . Let \mathcal{S}^+ be a directed graph on $[m]$, and \mathcal{S} be the base graph of \mathcal{S}^+ . Also let G be an m -partite k -graph with vertex classes U_1, \dots, U_m each of size n , and let K be the k -partite k -graph whose vertex classes each have size b_1 . Suppose also that for any sets $V_\ell \subseteq U_\ell$ with $|V_\ell| \geq n/2$ for each $\ell \in [m]$ the following statements hold for $V := \bigcup_{\ell \in [m]} V_\ell$.

- (i) For any edge $e \in \mathcal{R}$ there are at least N vertex-disjoint copies of K in $G[\bigcup_{\ell \in e} V_\ell]$.
- (ii) For any edge $i \rightarrow j$ of \mathcal{S}^+ , there are at least N vertex-disjoint copies of $\Phi(b_1)$ in $G[V]$ whose end vertex classes lie in V_i and V_j and whose central vertex classes lie in the sets V_ℓ for $\ell \in e(i) \setminus \{i\}$.

Note that (ii) implies that $e(i) \neq e(j)$ for any edge $ij \in \mathcal{S}$. If $i \rightarrow j$ is an edge of \mathcal{S}^+ , then condition (ii) allows us to choose a copy of K in G with one vertex in V_j , $b_1 - 1$ vertices in V_i , and b_1 vertices in V_ℓ for $\ell \in e(i) \setminus \{i\}$. Deleting this copy reduces the size of V_j by one relative to the sets V_ℓ for $\ell \in e(j) \setminus \{j\}$, and increases the size of V_i by one relative to the sets V_ℓ for $\ell \in e(i) \setminus \{i\}$. The next lemma states that, if \mathcal{R} is irreducible on $M_{\mathcal{R}}$, we can delete copies of K to achieve the same effect if $j \rightarrow i$ is an edge of \mathcal{S}^+ . This allows us to ignore the direction of edges of \mathcal{S}^+ and consider only the base graph \mathcal{S} when proving Lemma 6.10, which significantly simplifies the argument.

Lemma 6.9. *Adopt Setup 6.8, and suppose additionally that \mathcal{R} is (C, L) -irreducible on $M_{\mathcal{R}}$ for some C and L with $kb_1L + b_1C \leq N$, and that we now have fixed sets V_ℓ with $|V_\ell| \geq n/2$ for each $\ell \in [m]$. Then for any $i, j \in [m]$ with $ij \in \mathcal{S}$ there is a K -packing M in $G[V]$ of size at most $C + L$ such that, if we write $M(\ell) := |V(M) \cap V_\ell|$ for $\ell \in [m]$, we have the following properties.*

- (i) $M(\ell)$ is divisible by b_1 for any $\ell \in [m] \setminus \{i, j\}$.
- (ii) $M(i) = M(\ell) - 1$ for any $\ell \in e(i) \setminus \{i\}$.
- (iii) $M(j) = M(\ell) + 1$ for any $\ell \in e(j) \setminus \{j\}$.
- (iv) $M(\ell) = M(\ell')$ for any $e \in M_{\mathcal{R}} \setminus \{e_i, e_j\}$ and any $\ell, \ell' \in e$.

Proof. Fix any $i, j \in [m]$ with $ij \in \mathcal{S}$. Then there is an edge between i and j in \mathcal{S}^+ , directed either $i \rightarrow j$ or $j \rightarrow i$. Suppose first that $i \rightarrow j$; then there are at least $N \geq 1$ vertex-disjoint copies of $\Phi(b_1)$ in $G[V]$ whose end vertex classes lie in V_i and V_j and whose central vertex classes all lie in the sets V_ℓ for $\ell \in e(i) \setminus \{i\}$. Inside any one of these copies we can find a copy of K with $b_1 - 1$ vertices in V_i , with one vertex in V_j , and with b_1 vertices in V_ℓ for each $\ell \in e(i) \setminus \{i\}$. We can then take M to consist of this single copy of K .

So we may assume that $j \rightarrow i$, so there are at least N vertex-disjoint copies of $\Phi(b_1)$ in $G[V]$ whose end vertex classes lie in V_i and V_j and whose central vertex classes all lie in the sets V_ℓ for $\ell \in e(j) \setminus \{j\}$. Also, since \mathcal{R} is (C, L) -irreducible on $M_{\mathcal{R}}$, we may choose $c \leq C$ and multisets T and T' of edges of \mathcal{R} and $M_{\mathcal{R}}$ respectively such that $|T|, |T'| \leq L$ and such that j appears in c more edges of T than of T' , i appears in c more edges of T' than of T , and any $\ell \in [m] \setminus \{i, j\}$ appears equally often in T as in T' . For each edge $e \in T$, with multiplicity, choose a copy of K in $G[\bigcup_{i \in e} V_i]$; since $N \geq kb_1L$ we may do this so that these copies are all vertex-disjoint. This gives a K -packing M' in $G[V]$ of size at most L such that $M'(\ell)$ is divisible by b_1 for any $\ell \in [m]$, $M'(i) = M'(\ell) - cb_1$ for any $\ell \in e(i) \setminus \{i\}$, $M'(j) = M'(\ell) + cb_1$ for any $\ell \in e(j) \setminus \{j\}$, and $M'(\ell) = M'(\ell')$ for any $e \in M_{\mathcal{R}} \setminus \{e_i, e_j\}$ and any $\ell, \ell' \in e$. Now, since M' covers at most Lkb_1 vertices of $G[V]$, and $N - Lkb_1 \geq cb_1 - 1$, we may choose $cb_1 - 1$ vertex-disjoint copies of $\Phi(b_1)$ in $G[V]$ which do not have any vertices in common with M' , whose end vertex classes lie in V_i and V_j and whose central vertex classes all lie in the sets V_ℓ for $\ell \in e(j) \setminus \{j\}$. Each of these copies contains a copy of K with one vertex in V_i , $b_1 - 1$ vertices in V_j , and b_1 vertices in V_ℓ for each $\ell \in e(j) \setminus \{j\}$; adding these copies of K to M' gives the desired K -packing M . \square

We can now prove the main result of this section.

Lemma 6.10. *Adopt Setup 6.8, and assume that $1/n, 1/m, \beta \ll \alpha, 1/C, 1/L, 1/k, 1/b_1$, and also that $N \geq kb_1L + b_1C$ and $2n/3 \leq n' \leq n$. Let subsets $Y_j \subseteq U_j$ satisfy $(1 - \beta)n' \leq |Y_j| \leq n'$ for each $j \in [m]$, and suppose that b_1 divides $|Y|$, where $Y := \bigcup_{j \in [m]} Y_j$. Also let X_1, \dots, X_s be sets which partition $[m]$ such that*

- (i) *For any $T \subseteq [m]$ with $|T| \leq \alpha m$, any $i \in [s]$ and any $x, y \in X_i \setminus T$, there is a path from x to y in $\mathcal{S}[[m] \setminus T]$ of length at most L .*

- (ii) For any submatching $M'_R \subseteq M_R$ of size $|M'_R| \geq (1 - \alpha/2)|M_R|$ the subgraph $\mathcal{R}[V(M'_R)]$ is (C, L) -irreducible on M'_R .
- (iii) The average size $Q_i := \sum_{j \in X_i} |Y_j|/|X_i|$ of sets Y_j corresponding to vertices of X_i is the same for every $i \in [s]$, and this common average size $Q := Q_1 = \dots = Q_s$ is an integer which is divisible by b_1 .

Then there is a K -packing M in $G[Y]$ such that

- (a) for any edge $e \in M_R$ the sets $Y_j \setminus V(M)$ for $j \in e$ have equal size, and furthermore,
- (b) this common size is a multiple of b_1 .

Proof. For each $j \in [m]$ let $n_j := |Y_j| - Q$, that is, the difference between the size of Y_j and the average size of the sets Y_ℓ . So n_j is an integer with $|n_j| \leq \beta n' \leq \beta n$ for any $j \in [m]$, and for any $i \in [s]$ we have $\sum_{j \in X_i} n_j = 0$ by (iii). Construct a multiset of pairs Γ iteratively as follows. Initially take Γ to be empty. If $n_j = 0$ for every $j \in [m]$, then terminate. Otherwise, since $\sum_{j \in X_i} n_j = 0$ for every $i \in [s]$, there must be $i \in [s]$ and $j, j' \in X_i$ such that $n_j < 0$ and $n_{j'} > 0$. Add (j, j') to Γ , increment n_j by one and decrement $n_{j'}$ by one, and repeat. Since $|n_j| \leq \beta n$ for each $j \in [m]$, we must terminate after at most βnm steps. At this point, we have $|\Gamma| \leq \beta nm$, and, writing s_j for the number of times j appears as the first coordinate of a member of Γ , and t_j for the number of times j appears as the second coordinate of a member of Γ , we have

$$(1) \quad |Y_j| = Q - s_j + t_j$$

for every $j \in [m]$. Furthermore, note that, returning to the original values of n_j , we have $s_j, t_j \leq |n_j| \leq \beta n$ for any $j \in [m]$.

Arbitrarily order the pairs of Γ , and take M initially to be empty; we now add at most $L(C + L)$ copies of K to M , and delete the vertices covered from G , for each member of Γ in turn. So suppose that we are considering the z th pair in Γ , say (j, j') . So $z \leq \beta nm$. Prior to this we have added at most $L(C + L)(z - 1)$ copies of K to M , so at most $kb_1 L(C + L)\beta nm$ vertices have been deleted. For each $\ell \in [m]$ let Y'_ℓ consist of the so far undeleted vertices of Y_ℓ ; we assume for now that $|Y'_\ell| \geq (1 - 7\alpha)n' \geq n/2$, and will justify this assumption later. Also, let M'_R be the submatching of M_R consisting of $e(i)$, $e(j)$, and every $e \in M_R$ other than $e(j)$ and $e(j')$ for which at most αn vertices of $\bigcup_{z \in e} Y_z$ have previously been deleted, and let $T = [m] \setminus V(M'_R)$ and $Y' = \bigcup_{\ell \in V(M'_R)} Y'_\ell$. Then $(\alpha n)(|T|/k) \leq kb_1 L(C + L)\beta nm$, so we find that $|T| \leq \alpha m$. So $\mathcal{R}(V(M'_R))$ is (C, L) -irreducible on M_R by (ii), whilst by (i) we may choose a path $j = v_0, \dots, v_p = j'$ from j to j' in $\mathcal{S}[[m] \setminus T]$ of length $p \leq L$ (since $j, j' \in X_i$ for some i by construction of Γ). Now, for each $x \in [p]$ in turn, apply Proposition 6.9 to choose a K -packing M'_x in $G[Y']$ of size at most $C + L$ which satisfies the conclusions of Proposition 6.9 with v_{x-1} and v_x in place of i and j , delete the vertices of M'_x from the sets Y_ℓ , and add the members of M'_x to M . Having done this for every $x \in [p]$, the net effect is that there are integers $n_z(e) \leq p(C + L)$ for $e \in M_R$ with the following property. For any $e \in M_R$, precisely $b_1 n_z(e)$ vertices were deleted from Y'_ℓ for every $\ell \in e$, except for two cases: $b_1 n_z(e(j)) - 1$ vertices were deleted from Y'_j , whilst $b_1 n_z(e(j')) + 1$ vertices were deleted from $Y'_{j'}$. At this point we proceed to consider the $(z + 1)$ th pair in Γ , and continue in this manner.

After we have completed this process for every pair in Γ , we find that for any $j \in [m]$ the total number of vertices that were deleted from the set Y_j is equal to $t_j - s_j + b_1 \sum_{z \leq |\Gamma|} n_z(e(j))$. Combining this with (1) we obtain

$$|Y_j \setminus V(M)| = Q - b_1 \sum_{z \leq |\Gamma|} n_z(e(j)).$$

Properties (a) and (b) follow, since for any $e \in M_{\mathcal{R}}$ and $j \in e$ we have $e(j) = e$, and we assumed in (iii) that Q is divisible by b_1 . So it remains only to justify our assumption that at any point we had $|Y'_\ell| \geq (1 - 7\alpha)n'$ for any $\ell \in [m]$. For this, fix any $\ell \in [m]$, and consider the number of vertices deleted from Y_ℓ over the course of the procedure. If more than αn vertices of Y_ℓ were deleted in total, then for some z the number of deleted vertices of Y_ℓ first exceeded αn when considering the z th pair of Γ . Whilst considering this pair we deleted at most $L(C + L)$ copies of K , and so at the end of this step the number of vertices deleted from Y_ℓ was at most $\alpha n + kb_1L(C + L) \leq 2\alpha n$. For all subsequent steps the edge $e(\ell)$ was excluded from $M'_{\mathcal{R}}$, and so vertices were only deleted from Y_ℓ when considering pairs (j, j') for which $j \in e(\ell)$ or $j' \in e(\ell)$. The number of such pairs is at most $\sum_{j \in e(\ell)} s_j + t_j \leq 2k\beta n$, and so at most a further $(kb_1L(C + L))(2k\beta n) \leq \alpha n$ vertices were deleted from Y_ℓ , giving a total of at most $3\alpha n \leq 6\alpha n'$ vertices deleted from Y_ℓ over the entire course of the procedure. Since initially we had $(1 - \beta)n' \leq |Y_\ell|$, this justifies our earlier assumption that $|Y'_\ell| \geq (1 - 7\alpha)n'$. \square

6.5. Packing complete k -partite k -graphs. The proof of Lemmas 3.1 and 3.2 will conclude by finding a perfect K -packing within the ‘top layer’ J_- of a D -universal k -partite k -complex J . Provided D is sufficiently large, such a packing exists if the complete k -partite k -graph G on the same vertex classes contains a perfect K -packing. In this section we give sufficient conditions to ensure that this is the case.

For Lemma 3.2, K is a complete k -partite k -graph on vertex classes of equal size. In this case it is elementary to determine whether a complete k -partite k -graph G contains a perfect K -packing.

Fact 6.11. *Let K be the complete k -partite k -graph with vertex classes each of size b_1 , and let G be the complete k -partite k -graph with vertex classes V_1, \dots, V_k . Then G contains a perfect K -packing if and only if $|V_1| = \dots = |V_k|$ and b_1 divides $|V_1|$.*

However, for Lemma 3.1 things are more complicated. The next lemma gives sufficient conditions which ensure that a complete k -partite k -graph G contains a perfect K -packing when K is a complete k -partite k -graph as in Lemma 3.1. Recall for this the definitions of $\mathcal{B}(K)$ and $\mathcal{U}_s(K)$ (Definition 3.5).

Lemma 6.12. *Suppose that $1/n \ll \beta \ll 1/b, 1/k$. Let K be the complete k -partite k -graph with vertex classes of size b_1, \dots, b_k , where $b_1 + \dots + b_k = b$ and the b_i are not all equal. Also let G be a complete k -partite k -graph with vertex classes V_1, \dots, V_k , where $V := V_1 \cup \dots \cup V_k$ has size n . Suppose that*

- (i) $bk \gcd(K)$ divides $|V_i|$ for each $i \in [k]$, and
- (ii) $|V_j| \geq n/k - \beta n$ for every $j \in [k]$.

Then G contains a perfect K -packing.

Proof. Introduce an integer s with $\beta \ll 1/s \ll 1/b, 1/k$. Then we may assume that $\mathcal{U}_s(K)$ is defined and so contains a perfect K -packing by Proposition 3.6; the same is true of $\mathcal{B}(K)$. Also note that (i) implies that $bk \gcd(K)$ divides n . For each $i \in [k - 1]$ define

$$(2) \quad d_i := \frac{|V_i| - n/k}{\gcd(K)} + d_{i-1},$$

with d_0 taken to be zero; then each d_i must be an integer by (i). So by (ii) we have $|d_i| \leq k\beta n + |d_{i-1}|$, which implies that $\sum_{i \in [k-1]} |d_i| \leq k^3\beta n$. Now, for each $i \in [k - 1]$, if d_i is positive then delete d_i copies of $\mathcal{U}_s(K)$ from G , each with $bs + \gcd(K)$ vertices in V_i , $bs - \gcd(K)$ vertices in V_{i+1} and bs vertices in each other vertex class. On the other hand, if d_i is negative

then delete d_i copies of $\mathcal{U}_s(K)$ from G each with $bs - \gcd(K)$ vertices in V_i , $bs + \gcd(K)$ vertices in V_{i+1} and bs vertices in each other vertex class. We also insist that all of these copies of $\mathcal{U}_s(K)$ are pairwise vertex-disjoint; this is not a problem since the total number of copies of $\mathcal{U}_s(K)$ deleted is $N := \sum_{i \in [k-1]} |d_i| \leq k^3 \beta n$, and so the total number of vertices deleted is

$$kbsN \leq bsk^4 \beta n \leq n/k - \beta n \leq \min_{i \in [k]} |V_i|.$$

For each $i \in [k]$ let X_i consist of the undeleted vertices of V_i , and let $X := X_1 \cup \dots \cup X_k$. Then by (2) we obtain

$$|X_i| = |V_i| - bsN - (d_i - d_{i-1}) \gcd(K) = n/k - bsN.$$

We conclude that $|X_1| = \dots = |X_k|$ and that b divides $|X_1|$. So $G[X]$ contains a perfect $\mathcal{B}(K)$ -packing by Fact 6.11; combined with the previously deleted copies of $\mathcal{U}_s(K)$ this gives a perfect $\{\mathcal{B}(K), \mathcal{U}_s(K)\}$ -packing of G . Propositions 3.4 and 3.6 then imply that G contains a perfect K -packing. \square

Unfortunately, Lemma 6.12 is not strong enough for our purposes. Indeed, it requires that the vertex classes of G all have approximately equal size, whilst we wish to find a perfect K -packing within a ‘lopsided’ complete k -partite k -graph G . For this we use the following corollary, which shows that we can indeed do this provided that $\sigma(G) > \sigma(K) + o(1)$, that is, if G is ‘less lopsided’ than K (it is not hard to see that if G and K are complete k -partite k -graphs and $\sigma(G) < \sigma(K)$ then there can be no perfect K -packing in G). Recall for this corollary the definition of $\mathcal{L}(K)$ (Definition 3.5); in particular, $\mathcal{L}(K)$ has $(k-1)!b$ vertices in total, with one vertex class of size $(k-1)!b\sigma(K)$ and $k-1$ vertex classes of size $(k-2)!b(1-\sigma(K))$.

Corollary 6.13. *Suppose that $1/n \ll \beta \ll \alpha, 1/b, 1/k$. Let K be the complete k -partite k -graph with vertex classes of size b_1, \dots, b_k , where $b_1 + \dots + b_k = b$ and the b_i are not all equal. Also let G be a complete k -partite k -graph with vertex classes V_1, \dots, V_k , where $|V_1| \leq |V_2|, \dots, |V_k|$ and $V := V_1 \cup \dots \cup V_k$ has size n . Suppose that*

- (i) $\sigma(G) \geq \sigma(K) + \alpha$,
- (ii) $||V_i| - |V_j|| \leq \beta n$ for every $i, j \in \{2, \dots, k\}$, and
- (iii) $bk \gcd(K)$ divides $|V_i|$ for any $i \in [k]$.

Then G contains a perfect K -packing.

Proof. Let $d := \gcd(K)$, $\phi := \sigma(G)$ and $\sigma := \sigma(K)$, so $|V_1| = \phi n$ and by (ii) we have $|V_j| \geq (1 - \phi)n/(k-1) - \beta n$ for any $2 \leq j \leq k$. Without loss of generality we may assume that $b_1 \leq b_2, \dots, b_k$. Define $x := \frac{\phi - \sigma}{1/k - \sigma}$, so $\alpha \leq x \leq 1$ by (i). Then

$$(3) \quad \phi = \frac{x}{k} + \sigma(1 - x).$$

Define $N := \lfloor (1 - x)n/k!bd \rfloor$, and choose and delete a set of kdN pairwise vertex-disjoint copies of $\mathcal{L}(K)$ in G , such that the smallest vertex class of each copy is contained in V_1 . These copies thus cover $k!b\sigma Nd$ vertices of V_1 and $(k-2)!b(1-\sigma)kdN$ vertices of V_i for each $2 \leq i \leq k$; in particular, the number of vertices covered in any vertex class V_j is divisible by kbd . For each $i \in [k]$ let X_i consist of the undeleted vertices of V_i , and let $X := X_1 \cup \dots \cup X_k$.

So $|X_i|$ is divisible by kbd for any $i \in [k]$ by (iii). Also, by choice of N we have

$$\begin{aligned} xn &\leq |X| = n - k!bdN \leq xn + k!bd, \\ |X_1| &= |V_1| - k!b\sigma dN \geq \phi n - (1-x)\sigma n = xn/k, \text{ and} \\ |X_j| &= |V_j| - (k-2)!b(1-\sigma)kdN \geq \frac{(1-\phi)n}{k-1} - \beta n - \frac{(1-x)(1-\sigma)n}{k-1} = \frac{xn}{k} - \beta n, \end{aligned}$$

for any $2 \leq j \leq k$, where the final equality in each of the last two lines holds by (3). Together these inequalities imply that $|X_j| \geq |X|/k - 2\beta n$ for each $j \in [k]$. Furthermore, together with $x \geq \alpha$ the first inequality shows that we may assume that $1/|X| \ll 2\beta \ll 1/b, 1/k$. So $G[X]$ meets the conditions of Lemma 6.12, and so contains a perfect K -packing. Added to the deleted copies of $\mathcal{L}(K)$ this gives a perfect $\{K, \mathcal{L}(K)\}$ -packing of G ; by Propositions 3.4 and 3.6 it follows that G has a perfect K -packing. \square

7. PROOFS

We have now established all of the preliminary results and definitions we need for the proofs of Lemma 3.1 and Lemma 3.2, for which we proceed as outlined in Section 4.

7.1. Proof of Lemma 3.1. Recall the statement of Lemma 3.1.

Lemma 3.1. *Let K be the complete k -partite k -graph whose vertex classes have sizes b_1, \dots, b_k , where these sizes are not all equal, and suppose that $\gcd(K)$ and b_1 are coprime. Then for any $\alpha > 0$ there exists $n_0 = n_0(K, \alpha)$ such that the following statement holds. Let H be a k -graph on $n \geq n_0$ vertices such that*

- (a) $b := b_1 + \dots + b_k$ divides n ,
- (b) $\delta(H) \geq \sigma(K)n + \alpha n$, and
- (c) if $\gcd(K) > 1$, then $\delta(H) \geq n/p^* + \alpha n$, where p^* is the smallest prime factor of $\gcd(K)$.

Then H contains a perfect K -packing.

Proof. If $\gcd(K) = 1$, define $p^* := b$; this is merely for notational convenience in handling the cases $\gcd(K) = 1$ and $\gcd(K) > 1$ simultaneously. Since $\sigma(K) \geq 1/b$ we may then assume that $\delta(H) \geq n/p^* + \alpha n$ in all cases. Furthermore, in any case we must have $p^* \leq b$. Introduce new constants with

$$\begin{aligned} 1/n &\ll 1/N \ll \varepsilon \ll d^* \ll 1/a \ll \xi, 1/r \ll \nu \ll \mu \\ &\ll c, \eta \ll \theta \ll \psi \ll \beta \ll 1/q \ll \alpha \ll 1/s, 1/D \ll 1/b, 1/k, \end{aligned}$$

and such that s is divisible by $k \gcd(K)$, and choose an integer p such that

$$\sigma(K) + \alpha/3 \leq p/q \leq \sigma(K) + \alpha/2.$$

Note that our constant hierarchy assumes that α is sufficiently smaller than $1/s, 1/D, 1/b, 1/k$; this is not a problem since each property involving α in the statement of Lemma 3.1 is monotone. Using this and the fact that the vertex class sizes b_1, \dots, b_k are not all equal, we may assume that $\sigma(K) \leq 1/k - \alpha/2$, so $pk \leq q$. Finally, we may also assume that $a!r$ divides n . Indeed, by Theorem 3.7 we may greedily delete a K -packing in H of size up to $a!r$ so that the number of vertices remaining in H is divisible by $a!r$, following which the subgraph induced by the remaining vertices of H satisfies the conditions of the lemma (with weaker constants). So to prove the lemma it is sufficient to consider only the case where $a!r$ divides n . We assume this, so in fact no vertices were deleted.

Apply the Regular Approximation Lemma: Let $U := V(H)$, and choose arbitrarily a partition \mathcal{Q} of U into r parts T_1, \dots, T_r of equal size. Let H' be the k -graph on U consisting of all \mathcal{Q} -partite edges of H . So H' is a \mathcal{Q} -partite k -graph on U whose order is divisible by $a!r$, and so we may apply the Regular Approximation Lemma (Theorem 5.1), which yields an a -bounded ε -regular vertex-equitable partition $(k-1)$ -complex \mathcal{P} on U , and a \mathcal{Q} -partite k -graph G on U such that G is ξ -close to H' and the partition k -complex $G[\hat{\mathcal{P}}]$ is ε -regular. Let $Z = G \Delta H'$, so Z is a \mathcal{Q} -partite k -graph on U with $|Z| \leq \xi n^k$, and we have $G \setminus Z \subseteq H' \subseteq G \cup Z$. In particular, any edge of $G \setminus Z$ is also an edge of H . Also let U_1, \dots, U_m be the clusters of \mathcal{P} , and note that the partition $\mathcal{P}^{(1)}$ of U into clusters refines \mathcal{Q} . Since \mathcal{P} is vertex-equitable, every cluster of \mathcal{P} must have the same size, so the number of clusters m is divisible by r . Also, since \mathcal{P} is a -bounded we have $r \leq m \leq ar$. Define $n_1 := |U_i| = n/m$ to be the common cluster size. Then, since $a!r$ divides n and $m \leq ar$ is divisible by r , we deduce that n_1 is divisible by all integers up to $a/2$; in particular, n_1 is certainly divisible by $(q-p)p(k-1)$.

Observe that the vertex set U , the partition \mathcal{Q} of U into parts T_1, \dots, T_r , the partition $(k-1)$ -complex \mathcal{P} with clusters U_1, \dots, U_m , and the \mathcal{Q} -partite k -graphs G and Z therefore satisfy the conditions of Setup 5.7, with U and the clusters U_1, \dots, U_m in place of X and X_1, \dots, X_m , and with constants $1/n \ll \varepsilon \ll d^* \ll 1/a \ll \xi, 1/r \ll \nu \ll \mu \ll c, \eta \ll \theta \ll 1/D \ll 1/k$ playing identical roles there as here. Under this setup, let \mathcal{R} be the reduced k -graph of G and Z as defined in Definition 5.8. So \mathcal{R} has vertex set $[m]$, where vertex i corresponds to the cluster U_i . Since H' contains all \mathcal{Q} -partite edges of H , and $H' \subseteq G \cup Z$, for any \mathcal{Q} -partite $(k-1)$ -tuple $e \in \binom{U}{k-1}$ we have $\deg_{G \cup Z}(e) \geq \deg_{H'}(e) \geq \delta(H) - (k-1)n/r \geq \sigma(K)n + 2\alpha n/3$. So by Lemma 5.11, applied with $\sigma(K) + 2\alpha/3$ in place of γ ,

- (i) all but at most θm^{k-1} many $(k-1)$ -tuples $S \in \binom{[m]}{k-1}$ have

$$\deg_{\mathcal{R}}(S) \geq \sigma(K)m + 2\alpha m/3 - \theta m \geq pm/q + \theta m.$$

Refine the regularity partition into ‘lopsided’ groups: By (i) we may apply Lemma 6.1 to obtain an $\mathcal{A}_{p,q}^k$ -packing \mathcal{F} in \mathcal{R} so that

- (ii) \mathcal{F} covers at least $(1 - \psi)m$ vertices of \mathcal{R} , and
- (iii) $\mathcal{R}[V(\mathcal{F})]$ is connected,

where $V(\mathcal{F})$ denotes the set of vertices of \mathcal{R} covered by \mathcal{F} . Define a graph (*i.e.* 2-graph) \mathcal{S} with vertex set $V(\mathcal{F})$, where ij is an edge of \mathcal{S} if there exists a $(k-1)$ -tuple $S \in \binom{V(\mathcal{F})}{k-1}$ for which the triple (i, S, j) is Φ -dense and Z -sparse (as defined in Definition 5.8). Then every $i \in V(\mathcal{S})$ must lie in some edge S of \mathcal{R} (since i is covered by \mathcal{F}). Since for any \mathcal{Q} -partite $(k-1)$ -tuple $e \in \binom{U}{k-1}$ we have $\deg_{G \cup Z}(e) \geq \deg_{H'}(e) \geq \delta(H) - kn/r \geq n/p^* + 2\alpha n/3$, Lemma 5.11 (applied with $1/p^* + 2\alpha/3$ in place of γ) then implies that there are at least $m/p^* + \alpha m/2$ choices of $j \in [m] \setminus S$ such that the triple $(i, S \setminus \{i\}, j)$ is Φ -dense and Z -sparse. At most ψm of these choices of j do not lie in $V(\mathcal{S})$, so we have $\deg_{\mathcal{S}}(i) \geq m/p^* + \alpha m/2 - \psi m > m/p^*$. So

$$\delta(\mathcal{S}) > m/p^* \geq |V(\mathcal{S})|/p^*,$$

from which we conclude that each of the connected components C_1, \dots, C_{s^*} of \mathcal{S} contains more than m/p^* vertices. In particular, for any $x_1, \dots, x_{k-1} \in [s^*]$, there are at least $\binom{m/p^*}{k-1} > \theta m^{k-1}$ many $(k-1)$ -tuples $\{u_1, \dots, u_{k-1}\}$ with $u_j \in V(C_{x_j})$ for each $j \in [k-1]$, so by (i) at least one of these $(k-1)$ -tuples must have degree at least $pm/q > \psi m$ in \mathcal{R} . This proves that

- (iv) \mathcal{S} has fewer than p^* connected components C_1, \dots, C_{s^*} , and for any $x_1, \dots, x_{k-1} \in [s^*]$ there is some edge $\{u_1, \dots, u_k\} \in \mathcal{R}[V(\mathcal{S})]$ such that $u_j \in V(C_{x_j})$ for each $j \in [k-1]$.

Now, fix any $F \in \mathcal{F}$, and let $U_F := \bigcup_{j \in V(F)} U_j$. Then since each cluster U_j has size n_1 , which is divisible by $(q-p)p(k-1)$, by Lemma 6.2 we may partition U_F into disjoint sets V_j^i with $j \in [k]$ and $i \in [(q-p)p(k-1)]$ such that

- (v) $|V_1^i| = \frac{p}{q} \sum_{j \in [k]} |V_j^i|$ for each i ,
- (vi) $\frac{n_1}{q} \leq \frac{n_1}{q-p} = |V_1^i| \leq |V_2^i| = |V_3^i| = \dots = |V_k^i|$ for each i , and
- (vii) for each i and j there exists $f(i, j) \in [m]$ for which $V_j^i \subseteq U_{f(i, j)}$ and such that for any fixed i the set $\{f(i, j) : j \in [k]\}$ is an edge of \mathcal{R} .

Partition U_F in this manner for every $F \in \mathcal{F}$ to obtain sets V_j^i for $j \in [k]$ and $i \in [t]$, where $t := (q-p)p(k-1)|\mathcal{F}|$. We will refer to the sets V_j^i as *subclusters*.

We naturally obtain from \mathcal{R} a k -graph \mathcal{R}' corresponding to our refined partition into subclusters. Indeed, this has vertex set $[t] \times [k]$, where the vertex (i, j) corresponds to the subcluster V_j^i , and a set $\{(i_1, j_1), \dots, (i_k, j_k)\}$ is an edge of \mathcal{R}' if and only if $\{f(i_1, j_1), \dots, f(i_k, j_k)\}$ is an edge of \mathcal{R} . That is, edges of \mathcal{R}' correspond to k -tuples of subclusters which were taken from clusters of the same edge of the reduced k -graph. In the same way we define a graph \mathcal{S}' on the vertex set $[t] \times [k]$, where $\{(i_1, j_1), (i_2, j_2)\}$ is an edge of \mathcal{S}' if and only if $\{f(i_1, j_1), f(i_2, j_2)\}$ was an edge of \mathcal{S} . It follows from this definition that the components of \mathcal{S}' correspond to the components of \mathcal{S} . That is, \mathcal{S}' has components C'_1, \dots, C'_{s^*} , where for any $\ell \in [s^*]$ we have $(i, j) \in V(C'_\ell)$ if and only if $f(i, j) \in V(C_\ell)$. It then follows from (iv) that

- (viii) \mathcal{S}' has fewer than p^* connected components C'_1, \dots, C'_{s^*} , and for any $x_1, \dots, x_{k-1} \in [s^*]$ there is some edge $e = \{(i_1, j_1), \dots, (i_k, j_k)\} \in \mathcal{R}'$ such that $(i_\ell, j_\ell) \in V(C'_{x_\ell})$ for each $\ell \in [k-1]$. Also,
- (ix) \mathcal{R}' is connected. Indeed, if $f(i, j) = \ell$ and $f(i', j') = \ell'$, and ℓ and ℓ' were contained in a common edge of \mathcal{R} , then (i, j) and (i', j') are contained in a common edge of \mathcal{R}' by definition of \mathcal{R}' . Since $\mathcal{R}[V(\mathcal{F})]$ is connected by (iii), this implies that \mathcal{R}' is connected.

Obtain robustly universal complexes: Fix any $i \in [t]$. Then $e := \{f(i, j) : j \in [k]\}$ is an edge of \mathcal{R} by (vii), and $|V_j^i| \geq n_1/q \geq \eta n_1$ for each $j \in [k]$ by (vi). So we may apply Lemma 5.10 with the clusters $U_{f(i, j)}$ and subclusters V_j^i in place of the clusters X_i and subsets Y_i respectively. This allows us to delete at most $\mu|V_j^i|$ vertices from each subcluster V_j^i to obtain subsets $W_j^i \subseteq V_j^i$ and a k -partite k -complex J^i with vertex classes W_1^i, \dots, W_k^i such that

- (x) $d(J_{[k]}^i) > d^*$ and $|J_{\leq}^i(v)| > d^*|J_{\leq}^i|/|W_j^i|$ for every $v \in W_j^i$,
- (xi) $J_{\leq}^i \subseteq G \setminus Z$, and J^i is η -robustly D -universal.

Let W_0 be the set of all vertices of H which do not lie in any set W_j^i . By (ii) there are at most ψn vertices which lie in clusters U_ℓ for $\ell \in [m] \setminus V(\mathcal{F})$; the set W_0 contains all these, and also the at most μn vertices deleted whilst forming the sets W_j^i . So $|W_0| \leq \psi n + \mu n \leq 2\psi n$. Next, for each i and j choose an integer s_j^i such that $bk \gcd(K)$ divides s_j^i and

$$\beta n_1 \leq s_j^i \leq 2\beta n_1,$$

and choose a subset $X_j^i \subseteq W_j^i$ of size precisely s_j^i uniformly at random and independently of each other choice. Also let $Y_j^i = W_j^i \setminus X_j^i$ for every i and j , and for each $i \in [t]$ write $X^i := \bigcup_{j \in [k]} X_j^i$, $Y^i := \bigcup_{j \in [k]} Y_j^i$, $X = \bigcup_{i \in [t]} X^i$ and $Y = \bigcup_{i \in [t]} Y^i$. So the sets X, Y and W_0 partition $V(H)$. Then we may fix an outcome of these random selections so that

- (xii) for any $i \in [t]$ and any subset $Y'''^i \subseteq Y^i$, the subcomplex $J^i[X^i \cup Y'''^i]$ is D -universal, and
- (xiii) $\delta(H[W_0 \cup Y]) \geq \delta(H) - |X| \geq \alpha n - \alpha n/2 = \alpha n/2$.

Indeed, (xiii) follows from the fact that $|X| \leq tk(2\beta n_1) \leq \alpha n/2$, whilst for any specific $i \in [t]$ (xii) holds with probability $1 - o(1)$ by Proposition 5.5. Since the selections for distinct $i \in [t]$ are independent, with positive probability (xii) holds for every $i \in [t]$.

Delete a K -packing covering ‘bad’ vertices: We will next greedily form a K -packing M in $H[W_0 \cup Y]$ of size $|W_0|$ which covers every vertex of W_0 and which covers at most $\sqrt{\psi}n_1$ vertices from any set Y^i . To do this, suppose that we have already chosen fewer than $|W_0|$ members of M , and that $v \in W_0$ is not yet covered by M . Then M covers fewer than $b|W_0| \leq 2b\psi n$ vertices of H , so there are at most $2b\psi n/(\sqrt{\psi}n_1/2) \leq 4b\sqrt{\psi}m$ sets Y^i for which more than $\sqrt{\psi}n_1/2$ vertices of Y^i are covered by M . Let $Y' \subseteq Y$ consist of all vertices not yet covered by M which lie in sets Y^i in which at most $\sqrt{\psi}n_1/2$ vertices of Y^i are covered by M . Then

$$|Y'| \geq |Y| - (4b\sqrt{\psi}m)kn_1 - 2b\psi n \geq |Y| - \frac{\alpha n}{4},$$

so by (xiii) we have

$$\delta(H[\{v\} \cup Y']) \geq \frac{\alpha n}{2} - |(Y \cup W_0) \setminus Y'| \geq \frac{\alpha n}{2} - \frac{\alpha n}{4} - 2\psi n \geq \frac{\alpha n}{5}.$$

So by Lemma 6.4 there is a copy of K in $H[\{v\} \cup Y']$ which contains v ; choose such a copy and add it to M . Proceeding greedily in this manner, after $|W_0|$ steps we obtain a K -packing M in $H[W_0 \cup Y]$ which covers every vertex of $|W_0|$, and which covers at most $\sqrt{\psi}n_1/2 + b \leq \sqrt{\psi}n_1$ vertices in any set Y^i . Write $Y_j'' := Y_j^i \setminus V(M)$ for each $i \in [t], j \in [k]$ and let $Y' := \bigcup_{i,j} Y_j''$.

Delete a K -packing to ensure divisibility of cluster sizes: We now delete a further K -packing in $H[Y']$, so that after these deletions the number of vertices remaining in any set Y_j^i is divisible by $bk \gcd(K)$. To do this we apply Lemma 6.7 with $\mathcal{R}', \mathcal{S}', H'[Y']$ and kt in place of $\mathcal{R}, \mathcal{S}, G$ and t respectively, and the sets Y_j'' for $i \in [k]$ and $j \in [t]$ in place of the sets Y_j of Lemma 6.7. (We use the subgraph $H'[Y']$ rather than simply $H[Y']$ in place of G due to the requirement in Lemma 6.7 that G should be t -partite.) To see that b divides $|Y'|$, note that $|X|$ is divisible by b by choice of the integers s_j^i , that $|V(M)|$ is divisible by b since M is a K -packing, and $|V(H)|$ is divisible by b by assumption; since X and $V(M)$ are disjoint and $Y' = V(H) \setminus (V(M) \cup X)$ we indeed have that b divides $|Y'|$. Furthermore, \mathcal{R}' is connected by (ix), and condition (iii) of Lemma 6.7 holds by (viii) (since if $\gcd(K) > 1$ then p^* is the smallest prime factor of $\gcd(K)$). So it remains to verify that conditions (i) and (ii) of Lemma 6.7 are satisfied.

For condition (i), consider any edge $e \in \mathcal{R}'$, and let (i_x, j_x) for $x \in [k]$ be the vertices of e . Also let $v_x := f(i_x, j_x)$ for each $x \in [k]$; then $\{v_1, \dots, v_k\}$ is an edge of $\mathcal{R}[V_{\mathcal{F}}]$ by definition of \mathcal{R}' . We apply Lemma 5.10 with clusters U_{v_x} and subclusters $Y_{j_x}''^{i_x}$ in place of the sets X_ℓ and Y_ℓ respectively, which is possible since $|Y_{j_x}''^{i_x}| \geq n_1/q - \sqrt{\psi}n_1 - 2\beta n_1 \geq \eta n_1$ for each pair (i_x, j_x) . Lemma 5.10 then yields a $(k+1)$ -partite k -complex J which covers at least $(1 - \mu)|Y_{j_x}''^{i_x}| \geq n_1/2q$ vertices of each vertex class $Y_{j_x}''^{i_x}$ such that $J_- \subseteq G \setminus Z \subseteq H'$ and such that J is D -universal. By a very similar argument to the proof of Proposition 5.6, the latter fact implies that J_- contains at least $\lfloor n_1/2qb(s+1) \rfloor > N$ vertex-disjoint copies of $\mathcal{B}(b(s+1))$, and so $H'[\bigcup_{(i_x, j_x) \in e} Y_{j_x}''^{i_x}]$ does so also.

Now let $(i_1, j_1)(i_{k+1}, j_{k+1})$ be an edge of \mathcal{S}' , and write $v_1 = f(i_1, j_1)$ and $v_{k+1} = f(i_{k+1}, j_{k+1})$. Then $v_1 v_{k+1}$ is an edge of \mathcal{S} by definition of \mathcal{S}' , and so by definition of \mathcal{S} there exists a $(k-1)$ -tuple $S \in \binom{V(\mathcal{S}) \setminus \{v_1, v_{k+1}\}}{k-1}$ such that (u, S, v) is a Φ -dense and Z -sparse triple. Let v_2, \dots, v_k be the vertices of S , and for each $2 \leq x \leq k$ choose some i_x and j_x such that $f(i_x, j_x) = v_x$. We apply Lemma 5.9 with $\{v_1, \dots, v_k\}$ and $\{v_2, \dots, v_{k+1}\}$ in place of A and B respectively, with the clusters U_{v_x} and subclusters $Y_{j_x}''^{i_x}$ for $x \in [k+1]$ in place of the sets X_ℓ and Y_ℓ

respectively. Similarly as before, this is possible since $|Y_{j_x}^{i_x}| \geq \eta n_1$ for each pair (i_x, j_x) , and then Lemma 5.9 then yields a $(k+1)$ -partite k -complex J which covers at least $n_1/2q$ vertices of each vertex class $Y_{j_x}^{i_x}$ such that $J_- \subseteq G \setminus Z \subseteq H'$ and such that J is D -universal on Φ (where we consider vertex x of Φ to correspond to the pair (i_x, j_x)). Then by Proposition 5.6, J_- contains $\lfloor (n_1/2q)/kbs \rfloor > N$ vertex-disjoint copies of $\Phi(b(s+1))$ whose end vertex classes lie in $Y_{j_x}^{i_x}$ for $x = 1$ and $x = k+1$ and whose central vertex classes lie in $Y_{j_x}^{i_x}$ for $2 \leq x \leq k$. Since $J_- \subseteq H'$, this establishes condition (ii) of Lemma 6.7.

So we may indeed apply Lemma 6.7 as claimed. This yields a K -packing M' in $H'[Y']$ (and therefore also in $H[Y']$) of size at most N/b so that, taking $Y''^i_j := Y'^i_j \setminus V(M')$ for each i and j , we have that $bk \gcd(K)$ divides $|Y''^i_j|$ for every $i \in [t]$ and $j \in [k]$.

Blow-up a perfect K -packing in the remaining k -graph: To finish the proof, let $L^i_j := X^i_j \cup Y''^i_j$ for every $i \in [t]$ and $j \in [k]$, and for each $i \in [t]$ let $L^i := \bigcup_{j \in [k]} L^i_j$. So the sets L^i partition $V(H) \setminus V(M \cup M')$. Fix any $i \in [t]$, and observe that the k -complex $J^i[L^i]$ is D -universal by (xii). Since $J_-^i \subseteq G \setminus Z \subseteq H$, this implies that $H[L^i]$ contains a perfect K -packing if $\mathcal{K}[L^i]$, the complete k -partite k -graph on vertex classes L^i_1, \dots, L^i_k , does also. For any $j \in [k]$, by choice of the sets X^i_j and the K -packing M' both $|X^i_j|$ and $|Y''^i_j|$ are divisible by $bk \gcd(K)$, so $bk \gcd(K)$ divides $|L^i_j|$ also. Furthermore, L^i_j was formed from V^i_j by first deleting at most $\mu|V^i_j|$ vertices to form W^i_j , and then deleting the at most $\sqrt{\psi}n_1 + N$ vertices covered by $M \cup M'$. Since $|V^i_j| \geq n_1/q$ by (vi), in total at most $\beta|V^i_j|$ vertices were deleted in forming L^i_j from V^i_j , and in particular we have $|L^i_j| \geq |V^i_j|/2$. Writing $V^i := \bigcup_{j \in [k]} V^i_j$, by (v), (vi) and our choice of p , it follows that for any $j \in [k]$ we have

$$|L^i_j| \geq (p/q)|V^i| - \beta|V^i_j| \geq (p/q)|L^i| - 2\beta|L^i_j| \geq (\sigma(K) + \alpha/4)|L^i|,$$

so $\sigma(\mathcal{K}[L^i]) \geq \sigma(K) + \alpha/4$, and also that $||L^i_j| - |L^i_{j'}|| \leq \beta|V^i| \leq 2\beta|L^i|$ for any $2 \leq j, j' \leq k$. So by Corollary 6.13 $\mathcal{K}[L^i]$ admits a perfect K -packing, and so $H[L^i]$ contains a perfect K -packing M^i . Having chosen such a K -packing M^i for every $i \in [t]$, the union $M \cup M' \cup \bigcup_{i \in [t]} M^i$ is a perfect K -packing in H . \square

7.2. Proof of Lemma 3.2. Recall the statement of Lemma 3.2.

Lemma 3.2. *Let K be the complete k -partite k -graph whose vertex classes each have size b_1 . Then for any $\alpha > 0$ there exists $n_0 = n_0(K, \alpha)$ such that if $n \geq n_0$ is divisible by $b_1 k$ and H is a k -graph on n vertices with $\delta(H) \geq n/2 + \alpha n$ then H contains a perfect K -packing.*

In several places the proof of this lemma is similar to the proof of Lemma 3.1, in which case we refer to that proof.

Proof. Let $b := kb_1$, so $|V(K)| = b$, and introduce new constants with

$$1/n \ll \varepsilon \ll d^* \ll 1/a \ll \xi, 1/r \ll \nu \ll \mu \ll c, \eta \ll \theta \ll \psi \ll 1/C, 1/L \ll \alpha, 1/D \ll 1/b, 1/k,$$

As in the proof of Lemma 3.1, we have assumed without loss of generality that α is sufficiently smaller than $1/b$ and $1/k$. Furthermore, by the same argument as in the proof of Lemma 3.1 we may assume that n is divisible by $a!r$.

We begin by following the exact same steps as in the section ‘Apply the Regular Approximation Lemma’ of the previous proof, to obtain a partition \mathcal{Q} of $U := V(H)$ into parts T_1, \dots, T_r , a subgraph $H' \subseteq H$ consisting of all \mathcal{Q} -partite edges of H , a partition $(k-1)$ -complex \mathcal{P} on U with clusters U_1, \dots, U_m , and \mathcal{Q} -partite k -graphs G and $Z = G \triangle H'$ which

satisfy the conditions of Setup 5.7 (with variables taking the same values there as here). Likewise, as before we let \mathcal{R} be the reduced k -graph of G and Z as defined in Definition 5.8, so \mathcal{R} has vertex set $[m]$. Similarly as before we find that any \mathcal{Q} -partite $(k-1)$ -tuple $e \in \binom{U}{k-1}$ has

$$(4) \quad \deg_{G \cup Z}(e) \geq \deg_H(e) - (k-1)n/r \geq n/2 + 2\alpha n/3,$$

and we apply Lemma 5.11 to find that (with plenty of room to spare) all but at most θm^{k-1} many $(k-1)$ -tuples $S \in \binom{[m]}{k-1}$ have $\deg_{\mathcal{R}}(S) \geq m/k + \alpha m/2$. So we can apply Corollary 6.3 (with $\alpha/2$ in place of α) to find a matching $M_{\mathcal{R}}$ in \mathcal{R} which covers $m' \geq (1-\psi)m$ vertices of \mathcal{R} such that $\mathcal{R}[V(M'_{\mathcal{R}})]$ is (C, L) -irreducible on $M'_{\mathcal{R}}$ for any $M'_{\mathcal{R}} \subseteq M_{\mathcal{R}}$ with $|M'_{\mathcal{R}}| \geq (1-\alpha/4)|M_{\mathcal{R}}|$. Without loss of generality we assume that $V(M_{\mathcal{R}}) = [m']$.

Having chosen $M_{\mathcal{R}}$, we now define the graph \mathcal{S} on $V(M_{\mathcal{R}}) = [m']$; this definition is different to that used in the proof of Lemma 3.1. Indeed, we first define a directed graph \mathcal{S}^+ on $[m']$, where $i \rightarrow j$ is an edge of \mathcal{S} if the triple $(i, e_i \setminus \{i\}, j)$ is Φ -dense and Z -sparse, where e_i is the edge of $M_{\mathcal{R}}$ which contains i . We then define \mathcal{S} to be the base graph of \mathcal{S}^+ . By (4) and Lemma 5.11, applied with $1/2 + 2\alpha/3$ in place of γ , we find that for any $i \in [m']$ there are at least $m/2 + 2\alpha m/3 - \theta m$ choices of $j \in [m] \setminus e_i$ such that the triple $(i, e_i \setminus \{i\}, j)$ is Φ -dense and Z -sparse. At most ψm of these choices of j are not members of $[m']$, so we have $\delta(\mathcal{S}) \geq \delta^+(\mathcal{S}^+) > m/2 + \alpha m/2 \geq (1/2 + \alpha/2)m'$.

Obtain robustly universal complexes: We now obtain robustly universal complexes covering almost all of the vertices in clusters corresponding to edges of \mathcal{R} , similarly as in the proof of Lemma 3.1 (but here we do not divide our clusters into subclusters, so there is no need to define \mathcal{R}' and \mathcal{S}'). Fix any $e \in M_{\mathcal{R}}$. Then by Lemma 5.10 we can delete at most μn_1 vertices from each set U_j with $j \in e$ to obtain subsets W_j and a k -partite k -complex J^e with vertex classes W_1, \dots, W_k such that J^e is η -robustly D -universal, $J^e \subseteq G \setminus Z$, $d(J^e) > d^*$ and $|J^e(v)| > d^*|J^e|/|W_j|$ for every $v \in W_j$.

Let W_0 be the set of all vertices of H which do not lie in any set W_j . So W_0 contains the at most ψn vertices in clusters U_ℓ for $\ell \in [m] \setminus [m']$, and the at most μn vertices deleted from clusters U_j in forming the sets W_j . So we have $|W_0| \leq \psi n + \mu n \leq 2\psi n$. Next, fix an integer n_X with $\alpha n_1/3 \leq n_X \leq \alpha n_1/2$ such that n_X is divisible by b . For each $j \in [m']$ choose a subset $X_j \subseteq W_j$ of size precisely n_X uniformly at random, and take $Y_j := W_j \setminus X_j$. Then just as in the proof of Lemma 3.1, we find that, as there, we can fix an outcome of these random selections so that

(i) $\delta(H[W_0 \cup Y]) \geq \alpha n/2$, and

(ii) for any $e \in M_{\mathcal{R}}$ and any subset $Y_e'' \subseteq Y_e$, the subcomplex $J^e[X_e \cup Y_e'']$ is D -universal,

where we write $Y_e := \bigcup_{j \in e} Y_j$, $X_e := \bigcup_{j \in e} X_j$, and $Y := \bigcup_{e \in M_{\mathcal{R}}} Y_e$. Also let $X := \bigcup_{e \in M_{\mathcal{R}}} X_e$, and observe that the sets W , X and Y partition $V(H)$. So our assumption that b divides $|V(H)|$, together with our choice of n_X , implies that b divides both $|X|$ and $|W_0 \cup Y|$.

Delete a K -packing covering ‘bad’ vertices: Exactly as in the corresponding part of the proof of Lemma 3.1, (i) allows us to repeatedly apply Lemma 6.4 to greedily form a K -packing M^1 in $H[W_0 \cup Y]$ of size $|W_0|$ which covers every vertex of W_0 and which covers at most $\sqrt{\psi}n_1$ vertices from Y_e for any $e \in M_{\mathcal{R}}$. Following this, we apply Theorem 3.7 up to m' times to greedily choose a K -packing M^2 in $H[Y \setminus V(M^1)]$ of size at most m' , so that bm' divides $|Y \setminus V(M^1 \cup M^2)|$ (this is possible since $Y \setminus V(M^1)$ is identical to $(Y \cup W_0) \setminus V(M^1)$, so has size divisible by b). Write $Y'_j := Y_j \setminus V(M^1 \cup M^2)$ for every $j \in [m']$, and let $Y' := Y \setminus V(M^1 \cup M^2)$. Note that Y'_j was formed from U_j by deleting at most μn_1 vertices to form W_j , then exactly n_X vertices to form Y_j , and then at most $\sqrt{\psi}n_1 + bm'$ vertices which were covered by $M^1 \cup M^2$.

So, writing $n'_1 := n_1 - n_X$, for any $j \in [m]$ we have

$$(1 - 3\sqrt{\psi})n'_1 \leq n'_1 - 2\sqrt{\psi}n_1 \leq |Y'_j| \leq n'_1.$$

Delete a K -packing to ensure equality of cluster sizes: We now use Lemma 6.10 to delete a further K -packing M^3 in $H[Y']$ so that, following these deletions, the clusters Y'_ℓ for $\ell \in e$ have equal size for any edge $e \in M_{\mathcal{R}}$, and this common size is divisible by b_1 . For this, recall that $\mathcal{R}' := \mathcal{R}[V(M_{\mathcal{R}})]$ is a k -graph with vertex set $[m']$, and that $M_{\mathcal{R}}$ is a perfect matching in \mathcal{R}' , \mathcal{S}^+ is a directed graph on $[m']$, and \mathcal{S} is the base graph of \mathcal{S}^+ . We will show that $K, \mathcal{R}', M_{\mathcal{R}}, \mathcal{S}, \mathcal{S}^+$ and the clusters U_ℓ for $\ell \in [m']$ satisfy the conditions of Setup 6.8 with H', m', n_1 and $kb_1L + b_1C$ in place of G, m, n and N respectively. So fix any sets $V_\ell \subseteq U_\ell$ with $|V_\ell| \geq n_1/2$ for each $\ell \in [m']$, and let $V := \bigcup_{\ell \in [m']} V_\ell$.

Let $e \in \mathcal{R}'$. We apply Lemma 5.10 with clusters U_ℓ and subclusters V_ℓ in place of the sets X_ℓ and Y_ℓ respectively. Lemma 5.10 then yields a $(k+1)$ -partite k -complex J which covers at least $(1 - \mu)|V_\ell| \geq n_1/3$ vertices of each vertex class V_ℓ such that $J_- \subseteq G \setminus Z \subseteq H'$ and such that J is D -universal. So J_- contains at least $\lfloor n_1/3b_1 \rfloor > N$ vertex-disjoint copies of K , and so $H'[\bigcup_{\ell \in e} V_\ell]$ does so also. This demonstrates that condition (i) of Setup 6.8 is satisfied.

Now let $i \rightarrow j$ be an edge of \mathcal{S}^+ . By definition of \mathcal{S}^+ it follows that $(i, e_i \setminus \{i\}, j)$ is a Φ -dense and Z -sparse triple. Define $A := e_i$ and $B := \{j\} \cup e_i \setminus \{i\}$, and apply Lemma 5.9 with the sets V_ℓ and clusters U_ℓ for $\ell \in A \cup B$ in place of the sets Y_ℓ and X_ℓ respectively. Then Lemma 5.9 yields a $(k+1)$ -partite k -complex J with $J_- \subseteq G \setminus Z \subseteq H'$ which covers at least $(1 - \mu)|V_\ell| \geq n_1/3$ vertices of each V_ℓ and is D -universal on Φ . So by Proposition 5.6 J_- contains at least $\lfloor n_1/3b_1 \rfloor \geq N$ vertex-disjoint copies of $\Phi(b_1)$ in $J_- \subseteq H'[V]$ whose end vertex classes lie in V_i and V_j and whose central vertex classes lie in V_ℓ for $\ell \in e_i \setminus \{i\}$, so condition (ii) of Setup 6.8 is satisfied.

We will apply Lemma 6.10 with the trivial partition of $[m']$ into one set $X_1 = [m']$ (so $s = 1$). So condition (iii) of Lemma 6.10 requires simply that $|Y'|/m'$ is an integer which is divisible by b_1 ; this holds by our choice of M^2 and the fact that b_1 divides b . Furthermore, condition (ii) of Lemma 6.10 holds by our choice of $M_{\mathcal{R}}$. Finally, condition (i) of Lemma 6.10 holds since $\delta(\mathcal{S}) > (1/2 + \alpha/2)m'$, so for any $x, y \in [m']$ there are more than $\alpha m'$ paths of length two from x to y , and so these paths cannot all be removed by the deletion of at most $\alpha m'$ vertices of \mathcal{S} not including x or y . We can therefore apply Lemma 6.10, with $3\sqrt{\psi}$ and n'_1 in place of β and n' , and with b_1, α, k, C and L playing the same role there as here. From this we obtain a K -packing M^3 in $H'[Y']$ such that, writing $Y''_j := Y'_j \setminus V(M^3)$ for each $j \in [m']$, for each $e \in M_{\mathcal{R}}$ there is an integer n_e such that b_1 divides n_e and $|Y''_j| = n_e$ for any $j \in e$.

Blow-up a perfect K -packing in the remaining k -graph: To finish the proof, let $L_j := X_j \cup Y''_j$ for every $j \in [m']$, and for each $e \in M_{\mathcal{R}}$ let $L_e := \bigcup_{j \in e} L_j$. So the sets L_e for $e \in M_{\mathcal{R}}$ partition $V(H) \setminus V(M^1 \cup M^2 \cup M^3)$. Now fix any $e \in M_{\mathcal{R}}$. By choice of M^3 we have $|L_j| = |Y''_j| + |X_j| = n_e + n_X$ for any $j \in e$, and so $|L_j|$ is divisible by b_1 since both n_e and n_X were. So the complete k -partite k -graph $\mathcal{K}[L_e]$ with vertex classes L_j for $j \in e$ admits a perfect K -packing by Fact 6.11. Since the k -complex $J^e[L_e]$ is D -universal by (ii) it follows that $J^e_-[L_e]$ admits a perfect K -packing also. Finally, since $J^e_- \subseteq G \setminus Z \subseteq H$, this implies that $H[L_e]$ contains a perfect K -packing M^e . Choose M^e in this way for each $e \in M_{\mathcal{R}}$; then $M^1 \cup M^2 \cup M^3 \cup \bigcup_{e \in M_{\mathcal{R}}} M^e$ is a perfect K -packing in H . \square

8. PACKING LOOSE CYCLES

Recall that we write C_s^k to denote the loose cycle k -graph on $s(k-1)$ vertices, which was defined for any $s > 1$ to have $s(k-1)$ vertices $\{1, \dots, s(k-1)\}$ and s edges $\{\{j(k-1) + 1, \dots, j(k-1) + k\} \text{ for } 0 \leq j < s\}$, with addition taken modulo $s(k-1)$. Also recall that $\tau(K)$ denotes the proportion of vertices of K in a smallest vertex cover of K , whilst $\sigma(K)$ denotes the proportion of vertices of K in a smallest vertex class of a k -partite realisation of K . In this section we prove Theorem 1.4, giving the asymptotic value of $\delta(C_s^k, n)$ for any k and s . To begin, we establish the values of $\gcd(C_s^k)$, $\tau(C_s^k)$ and $\sigma(C_s^k)$.

Proposition 8.1. *For any $k \geq 3$ and $s \geq 2$ we have*

$$\tau(C_s^k) = \sigma(C_s^k) = \frac{\lceil s/2 \rceil}{s(k-1)}.$$

Furthermore, we also have $\gcd(C_s^k) = 1$ except in the case $s = k = 3$, whilst $\gcd(C_3^3)$ is undefined.

Proof. Note that the vertices $j(k-1) + 1$ for $0 \leq j < s$ are the vertices of C_s^k which lie in two edges of C_s^k . Let C be the (graph) cycle on these s vertices (in order). Then any proper k -colouring of C (as a graph) can be extended to a k -partite realisation of C_s^k by colouring the $k-2$ uncoloured vertices of each edge of C_s^k with the $k-2$ colours not used to colour the two coloured vertices. Furthermore, the size of each vertex class V_i is then simply s minus the number of vertices of C with colour i . Now, if $s \geq 4$ then we can 3-colour C with $\lfloor s/3 \rfloor$ red vertices, $\lfloor s/3 \rfloor + 1$ green vertices and all remaining vertices blue. Extending this colouring to a k -partite realisation of C_s^k as described above, we find that the red and green vertex classes differ in size by one, from which we conclude that $\gcd(C_s^k) = 1$. If instead $s = 3$ and $k \geq 4$, then we 3-colour C with one red, one blue and one green vertex; extending this 3-colouring of C to a k -partite realisation of C_s^k we find that the blue, green and red vertex classes each have one fewer vertex than each other vertex class, so again we have $\gcd(C_s^k) = 1$. Finally, if $s = 2$, then whilst C is no longer a simple graph, if we colour its two vertices red and blue and then extend this colouring to a k -partite realisation of C_s^k , we find that the red and blue vertex classes each have one fewer vertex than each other vertex class, again giving $\gcd(C_s^k) = 1$. So $\gcd(C_s^k) = 1$ in any case except for $k = s = 3$.

Next observe that any vertex of C_s^k lies in at most two edges of C_s^k , so any vertex cover of C_s^k has size at least $\lceil s/2 \rceil$; taking vertices $j(k-1) + 1$ for even $0 \leq j < s$ gives a vertex cover of this size. So $\tau(C_s^k) = \lceil s/2 \rceil / s(k-1)$ as claimed. We must have $\sigma(C_s^k) \geq \tau(C_s^k)$ since any vertex class of any k -partite realisation of C_s^k is a vertex cover of C_s^k . So it remains only to show that $\sigma(C_s^k) \leq \lceil s/2 \rceil / s(k-1)$, that is, that C_s^k has a k -partite realisation in which some vertex class has size $\lceil s/2 \rceil$. For this, observe that we can 3-colour C with $\lfloor s/2 \rfloor$ blue vertices, $\lfloor s/2 \rfloor$ red vertices, and either one or zero green vertices. Extending this colouring to a k -partite realisation of C_s^k we find that the blue vertex class has size $s - \lfloor s/2 \rfloor = \lceil s/2 \rceil$, as required.

Finally, observe that C_3^3 has only one 3-partite realisation up to permutation of the vertex classes $\{1, 4\}$, $\{2, 5\}$, $\{3, 6\}$. Since each vertex class has equal size, $\gcd(C_3^3)$ is undefined. \square

Except in the case $k = s = 3$, Proposition 8.1 shows that Theorem 1.2 and Proposition 2.3 give asymptotically matching upper and lower bounds on $\delta(C_s^k, n)$. However, since any 3-partite realisation of C_3^3 has two vertices in each vertex class, C_3^3 has type 0, and so our main theorems provide only the bound $\delta(C_3^3, n) \leq n/2 + o(n)$ which applies to all k -partite k -graphs. However, by modifying the proof of Lemma 3.2 we can actually prove that $\delta(C_3^3, n) \leq n/3 + o(n)$, giving the correct asymptotic threshold in this case also. For this we need the

following proposition, which allows us to find copies of C_3^3 with an odd number of vertices on each side of a partition of $V(H)$. Note that the k -graph constructed in Proposition 2.1 demonstrates that this proposition does not hold if we replace C_3^3 by the 3-partite 3-graph K with two vertices in each vertex class; this is the point at which the proof of Theorem 1.4 fails to hold for K .

Proposition 8.2. *Suppose that $1/n \ll \alpha$. Let H be a 3-graph on n vertices with $\delta(H) \geq n/3 + \alpha n$, and suppose that sets A and B partition $V(H)$ and satisfy $|A|, |B| \geq n/3 + \alpha n$. Then there is a copy of C_3^3 in H with an odd number of vertices in each of A and B .*

Proof. Suppose for a contradiction that no such copy of C_3^3 exists. There are $|A||B| \geq n^2/9$ pairs (x, y) with $x \in A$ and $y \in B$. Each of these pairs lies in at least $\delta(H) \geq n/3$ edges of H , so there are at least $n^3/81$ edges $e \in H$ which have at least one vertex in each of A and B . So without loss of generality we may assume that there are at least $n^3/200$ edges $e \in H$ with precisely two vertices in A . We now ‘colour’ the edges of the complete graph $K[A]$ on A with colours red and blue. Indeed, we colour xy red if there are at least 3 vertices $w \in B$ with $\{x, y, w\} \in H$, and we colour xy blue if there are at least 6 vertices $w \in A$ such that $\{x, y, w\} \in H$. So every edge xy receives at least one colour; if both conditions are satisfied then we give xy both colours, meaning that we can treat it as being either colour. Since any pair xy lies in at most n edges, we find that there are at least $(n^3/200 - 2n^2)/n \geq n^2/300$ red edges of $K[A]$.

Observe that there can be no triangle in $K[A]$ with three red edges. Indeed, if xyz is such a triangle then we may choose distinct $w_1, w_2, w_3 \in B$ such that $(x, y, w_1), (x, z, w_2)$ and (y, z, w_3) are each edges of H , thus forming a copy of C_3^3 with three vertices in A and three in B . Similarly, there can be no triangle in $K[A]$ with two blue edges and one red edge, as then we can form a copy of C_3^3 with one vertex in B and five in A . Now, choose any vertex $x \in A$ which lies in a red edge, and define $A_1 = \{y \in A \setminus \{x\} : xy \text{ is red}\}$ and $A_2 := A \setminus A_1$. So A_1 and A_2 partition A , and by our previous observations no edge of $K[A_1]$ or $K[A_2]$ is red. So all edges of $K[A_1]$ and $K[A_2]$ are blue and not red; it follows that every edge yz with $y \in A_1$ and $z \in A_2$ is red and not blue (so in fact every edge of $K[A]$ has only one colour). So every edge of $K[A]$ must have received a single colour. Moreover the red edges of $K[A]$ form a complete bipartite subgraph of $K[A]$ with vertex classes A_1 and A_2 . Since the number of red edges of $K[A]$ is at least $n^2/300$ it follows that $|A_1|, |A_2| \geq n/300$. Without loss of generality we may assume that $|A_1| \leq |A_2|$, so $|A_1| \leq (n - |B|)/2 \leq n/3 - \alpha n/2$.

Now let $y, z \in A_1$. Then there are at least $\delta(H) \geq n/3 + \alpha n$ vertices w such that $\{w, y, z\} \in H$. At most $n/3 - \alpha n/2$ of these vertices w lie in A_1 , and since yz is not red at most 2 of these vertices w lie in B . So there are at least αn vertices $w \in A_2$ such that $\{w, y, z\} \in H$; summing over all pairs $y, z \in A_1$ we find that there are at least $\binom{|A_1|}{2} \alpha n \geq 10^{-6} \alpha n^3$ edges of H with two vertices in A_1 and one vertex in A_2 . Since there are $|A_1||A_2| \leq n^2$ pairs yz with $y \in A_1$ and $z \in A_2$, we deduce that some such pair yz lies in at least $10^{-6} \alpha n \geq 6$ such edges of H . But then yz is blue, a contradiction. \square

We can now give the proof of Theorem 1.4. Note that, as mentioned above, the proof follows immediately from Proposition 8.1 except in the case $k = s = 3$.

Theorem 1.4. *For integers $k \geq 3$ and $s \geq 2$ we have*

$$\delta(C_s^k, n) = \begin{cases} \frac{n}{2(k-1)} + o(n) & \text{if } s \text{ is even, and} \\ \frac{s+1}{2s(k-1)}n + o(n) & \text{otherwise.} \end{cases}$$

Proof. Suppose first that we do not have $k = s = 3$. Then Proposition 8.1 shows that $\gcd(C_s^k) = 1$ and $\sigma(C_s^k) = \lceil s/2 \rceil / s(k-1)$, so Theorem 1.2 implies that $\delta(C_s^k, n) \leq \lceil s/2 \rceil n / s(k-1) + o(n)$. On the other hand, Proposition 8.1 also shows that $\tau(C_s^k) = \lceil s/2 \rceil / s(k-1)$, so Proposition 2.3 implies that $\delta(C_s^k, n) \geq \lceil s/2 \rceil n / s(k-1)$. This completes the proof except for the case $k = s = 3$.

Now suppose that $k = s = 3$. By Proposition 8.1 we have $\tau(C_3^3) = 1/3$, and it follows by Proposition 2.3 that $\delta(C_3^3, n) \geq n/3$. So it suffices to prove that $\delta(C_3^3, n) \leq n/3 + o(n)$. That is, we must show that for any $\alpha > 0$ there exists n_0 such that for any $n \geq n_0$ which is divisible by 6, any k -graph H on n vertices with $\delta(H) \geq n/3 + \alpha n$ contains a perfect C_3^3 -packing. To do this, let K denote the complete 3-partite 3-graph with vertex classes each of size two, and note that any copy of K contains a copy of C_3^3 on the same vertex set, so we can treat copies of K in H as being copies of C_3^3 for the purpose of finding a C_3^3 -packing in H . We mimic the proof of Lemma 3.2 as it would apply to K . Indeed, we use the same hierarchy of constants as in the proof of Lemma 3.2, except that we now have the fixed values $b_1 = 2, b = 6$ and $k = 3$, and we introduce two additional constants β and β' with $\psi \ll \beta \ll \beta' \ll 1/C, 1/L$. We proceed exactly as in the proof of Lemma 3.2 for most steps of the proof. In the first section of the proof, the calculation of (4) now only gives us $\deg_{G \cup Z}(e) \geq n/3 + 2\alpha n/3$, so Lemma 5.11 now yields the weaker result that at most θm^2 pairs $S \in \binom{[m]}{2}$ have $\deg_{\mathcal{R}}(S) \geq m/3 + \alpha m/2$, but this is still sufficient to apply Corollary 6.3 as in the proof of Lemma 3.2. However, at the end of the first section of that proof our weaker minimum codegree condition now only implies that $\delta(\mathcal{S}) \geq \delta^+(\mathcal{S}^+) > (1/3 + \alpha/2)m'$. This condition is only used in the step ‘Delete a K -packing to ensure equality of cluster sizes’, and indeed the rest of the proof proceeds exactly as before until that step (in particular we obtain $n_X \leq \alpha n_1/2$, $n'_1 = n_1 - n_X$, $m' \geq (1 - \psi)m$, K -packings M^1 and M^2 , and sets X_j and Y'_j as there).

In this step we wish to find a C_3^3 -packing M^3 in $H[Y']$ so that for any $e \in M_{\mathcal{R}}$ the sets Y'_j for $j \in e$ have equal size, and this common size is even. We demonstrate below how this can be done, but first note that once we have found such a C_3^3 -packing M^3 in $H[Y']$, the final step of the proof proceeds exactly as before to give a perfect C_3^3 -packing in H .

Suppose first that for any set $T \subseteq V(\mathcal{S})$ with $|T| \leq \alpha m'/6$ the graph $\mathcal{S} \setminus T$ formed by deleting the vertices of T from \mathcal{S} is connected (as was the case in the proof of Lemma 3.2 as a consequence of our stronger bound on $\delta(\mathcal{S})$). Then $\delta(\mathcal{S} \setminus T) \geq \delta(\mathcal{S}) - \alpha m'/6 > m'/3$, and for any $x, y \in V(\mathcal{S}) \setminus T$ there is a path from x to y in $\mathcal{S} \setminus T$. It follows that for any $x, y \in V(\mathcal{S}) \setminus T$ the shortest path in $\mathcal{S} \setminus T$ from x to y has length at most five, as in a path of length six or more two of the first, fourth and seventh vertices must share a common neighbour in $\mathcal{S} \setminus T$, allowing us to construct a shorter path. We can therefore apply Lemma 6.10 exactly as in the proof of Lemma 3.2, with the trivial partition of $V(\mathcal{S})$ into one set $X_1 = V(\mathcal{S})$, except that now we have $\alpha/6$ in place of α (condition (i) of Lemma 6.10 then follows by our remarks above, and all other conditions hold exactly as before).

We may therefore assume that there exists a set $T \subseteq V(\mathcal{S})$ with $|T| \leq \alpha m'/6$ for which the graph $\mathcal{S}' := \mathcal{S} \setminus T$ is disconnected. Observe that $\delta(\mathcal{S}') \geq \delta(\mathcal{S}) - \alpha m'/6 \geq m'/3 + \alpha m'/3$, so \mathcal{S}' has precisely two connected components C_1 and C_2 , each with at least $m'/3 + \alpha m'/3$ vertices and so at most $2m'/3 - \alpha m'/3$ vertices. In particular, any vertices $x, y \in V(C_1)$ have at least $\alpha m'$ neighbours in $V(C_1)$, and the analogous condition holds for V_2 . Furthermore, every vertex $v \in T$ has at least $m'/3 + \alpha m'/2$ neighbours in \mathcal{S} , and so has either at least $m'/6$ neighbours in $V(C_1)$ or at least $m'/6$ neighbours in $V(C_2)$. Form a set V_1 by adding to $V(C_1)$ all those vertices of T with at least $m'/6$ neighbours in $V(C_1)$, and let $V_2 = V(\mathcal{S}) \setminus V_1$, so $V(C_2) \subseteq V_2$ and every vertex of $T \cap V_2$ has at least $m'/6$ neighbours in $V(C_2)$. Then V_1 and V_2 partition $[m']$, and $m'/3 + \alpha m'/3 \leq |V_1|, |V_2| \leq 2m'/3 - \alpha m'/3$. Furthermore, writing

$\mathcal{S}_1 := \mathcal{S}[V_1]$, for any vertices $x, y \in V_1$ there is a path from x to y of length at most four in \mathcal{S}_1 from x to y , even after the deletion of at most $\alpha m'/6$ vertices other than x and y . Indeed, even following this deletion, x must have a neighbour w in $V(C_1)$, and y must have a neighbour z in $V(C_1)$, and then z and w have a common neighbour in $V(C_1)$ by our observation above that z and w have at least $\alpha m'$ common neighbours in \mathcal{S}' . A similar argument shows that if we delete at most $\alpha m'/6$ vertices of $\mathcal{S}_2 := \mathcal{S}[V_2]$ then there is a path of length at most four between any two vertices in the remaining subgraph.

Now recall from the proof of Lemma 3.2 that every set Y'_j has size $(1 - 3\sqrt{\psi})n'_1 \leq |Y'_j| \leq n'_1$. In particular, it follows that

$$(5) \quad n - 2\alpha n/3 \leq (1 - 3\sqrt{\psi})(1 - \alpha/2)n_1(1 - \psi)m \leq (1 - 3\sqrt{\psi})n'_1m' \leq |Y'| \leq n'_1m' \leq n.$$

Fix an integer Q which is divisible by 6 such that $(1 - \beta)n'_1 \leq Q \leq (1 - \beta + \psi)n'_1$.

Claim 8.3. *There exists a C_3^3 -packing M^* in $H[Y']$ of size*

$$N := \frac{|Y'| - m'Q}{6}$$

so that $|V(M^*) \cap Y'_j| \leq \beta'n'_1$ for any $j \in [m']$ and $|V(M^*) \cap \bigcup_{j \in V_1} Y'_j| = \sum_{j \in V_1} |Y'_j| - |V_1|Q$.

To prove the claim, we first observe that N is an integer, since Y' consists of all vertices of H except those in X (which was chosen in the proof of Lemma 3.2 to have size divisible by $b = 6$) and those covered by the K -packing $M^1 \cup M^2$, so $|Y'|$ is divisible by 6, and we chose Q to be divisible by 6. Also, we have $(\beta - \beta^2)m'n'_1 \leq 6N \leq \beta m'n'_1$ by (5) and our choice of Q . Write $A := \bigcup_{j \in V_1} Y'_j$ and $B := \bigcup_{j \in V_2} Y'_j$. Since $|Y'| \geq n - 2\alpha n/3$ by (5), we have $\delta(H[Y']) \geq (1/3 + \alpha/3)|Y'|$, so, by three successive applications of Proposition 8.2, we may obtain a set E_0 of three vertex-disjoint copies of C_3^3 in $H[Y']$, each of which has an odd number of vertices in each of A and B .

Now recall that all but at most θm^2 pairs $S \in \binom{[m]}{2}$ had $\deg_{\mathcal{R}}(S) \geq m/3 + \alpha m/2 \geq m'/3 + \alpha m'/2$. Since $m' \geq (1 - \psi)m$ and $|V_1|, |V_2| \geq m'/3$ we may greedily form a set of $m'/10$ disjoint pairs (x, y) with $x \in V_1, y \in V_2$ and $\deg_{\mathcal{R}}(\{x, y\}) \geq m'/3$. So certainly either there are at least $m'/10$ edges of \mathcal{R} with precisely two vertices in V_1 or there are at least $m'/10$ edges of \mathcal{R} with precisely two vertices in V_2 . Without loss of generality we assume the former, that there is a set F_1 of $m'/10$ vertex-disjoint edges of \mathcal{R} each with precisely two vertices in V_1 . The same argument for pairs (x, y) with $x, y \in V_2$ shows that there must be a set F_2 of $m'/10$ vertex-disjoint edges of \mathcal{R} each with at least two vertices in V_2 . Using Lemma 5.10 exactly as Lemma 5.9 was used in the proof of Lemma 3.2, we can obtain at least $\beta'n'_1/5$ vertex-disjoint copies of K in $H'[\bigcup_{j \in e} Y'_j]$ for each edge $e \in F_1 \cup F_2$. So we can greedily form K -packings E_1 and E_2 in $H[Y']$ each of size $(\beta'n'_1/5)(m'/40) = \beta'n'_1m'/200 \geq N$ such that $V(E_1) \cap V(E_2) \cap V(E_3) = \emptyset$, every copy of K in E_1 has precisely four vertices in A , every copy of K in E_2 has either four or six vertices in B , and collectively E_1, E_2 and E_3 cover at most $4\beta'n'_1/5 + 24 \leq \beta'n'_1$ vertices in any set Y'_j .

Recall that we want M^* to cover $|A| - |V_1|Q$ vertices of A . Initially take M^* to consist precisely N edges taken from E_0 and E_1 . By choosing an appropriate subset of the edges of E_0 to include, we can ensure that $|V(M^*) \cap A| - (|A| - |V_1|Q)$ is divisible by four. However, this initial selection of M^* has

$$|V(M^*) \cap A| \geq 4(N - 3) \geq \frac{2(\beta - \beta^2)m'n'_1}{3} - 12 > \left(\frac{2}{3} - \frac{\alpha}{3}\right)\beta m'n'_1 \geq |A| - |V_1|Q,$$

so too many vertices are taken from A (the final inequality holds since $|A| \leq n'_1|V_1|$ and $|V_1| \leq (2/3 - \alpha/3)m'$). On the other hand, if we were to replace all edges of E_1 in M^* by

edges of E_2 , then a similar calculation would show that $|V(M^*) \cap A| < |A| - |V_1|Q$, that is, that too few vertices are taken from A . Starting from our initial M^* , we repeatedly replace an edge $e \in E_1$ in M^* by an edge $e' \in E_2$, beginning with those edges $e' \in E_2$ with $|e' \cap B| = 6$, and then using edges $e' \in E_2$ with $|e' \cap B| = 4$ if these run out. Then each replacement by an edge e' with $|e' \cap B| = 6$ decreases $|V(M^*) \cap A|$ by four, and each replacement by an edge e' with $|e' \cap B| = 4$ decreases $|V(M^*) \cap A|$ by two. Since our initial M^* was chosen so that $|V(M^*) \cap A| - (|A| - |V_1|Q)$ was divisible by four, at some point in this process we must have $|V(M^*) \cap A| = |A| - |V_1|Q$. By our choice of E_1 and E_2 , at this point we also have $|V(M^*) \cap Y'_j| \leq \beta' n'_1$ for any $j \in [m']$, so this M^* is the desired C_3^3 -packing. This completes the proof of Claim 8.3.

Retuning to the proof of Theorem 1.4, fix some M^* as in Claim 8.3, and delete the vertices covered by M^* from H . Having done so, the sets $Y_j^* := Y'_j \setminus V(M^*)$ for each $j \in [m']$ of undeleted vertices satisfy

$$(6) \quad (1 - 2\beta')n'_1 \leq |Y'_j| - \beta'n'_1 \leq |Y_j^*| \leq |Y'_j| \leq n'_1$$

for any $j \in [m']$, and

$$(7) \quad Q = \frac{|\bigcup_{i \in V_1} Y_i^*|}{|V_1|} = \frac{|\bigcup_{i \in V_2} Y_i^*|}{|V_2|}.$$

where for the second equality we used the fact that M^* covered $6N = |Y'| - m'Q$ vertices in total, so, since V_1 and V_2 partition $[m']$, we have

$$|V(M^*) \cap \bigcup_{j \in V_2} Y'_j| = |Y'| - m'Q - \sum_{j \in V_1} |Y'_j| + |V_1|Q = \sum_{j \in V_2} |Y'_j| - |V_2|Q.$$

We now apply Lemma 6.10 similarly as in Lemma 3.2, but now with V_1 and V_2 in place of X_1 and X_2 , giving a partition of $[m']$ into two parts. Again we have $n_1 - n_X$ and m' in place of n and m respectively and $M_{\mathcal{R}}$ and \mathcal{S} play the same role there as here, but we now have the sets $Y_j^*, H'[Y^*]$, $\alpha/6$ and $2\beta'$ in place of Y_j , G , α and β respectively, and we set $b_1 = 2$ and $k = 3$. The sets Y_j^* satisfy the size condition of Lemma 6.10 by (6), and condition (iii) of Lemma 6.10 then holds by (7) (with Q playing the same role here as there). The conditions of Setup 6.8 are satisfied exactly as in the proof of Lemma 3.2, since these do not depend on the sets Y_j^* . Likewise condition (ii) of Lemma 6.10 holds by our choice of $M_{\mathcal{R}}$ exactly as in the proof of Lemma 3.2. Finally, condition (i) of Lemma 6.10 holds by our comments on \mathcal{S}_1 and \mathcal{S}_2 prior to the statement of Claim 8.3.

So we may indeed apply Lemma 6.10 as claimed, to obtain a K -packing M^{**} in $H'[Y^*]$ such that for any $e \in M_{\mathcal{R}}$ the sets $Y_j'' := Y_j^* \setminus V(M^{**}) = Y'_j \setminus V(M^* \cup M^{**})$ for $j \in e$ obtained by these deletions have equal size n_e , where n_e is divisible by $b_1 = 2$. So we may take $M^3 = M^* \cup M^{**}$, following which, as stated earlier, the final step of the proof proceeds exactly as in the proof of Lemma 3.2. \square

9. CONCLUDING REMARKS

9.1. Non-complete k -partite k -graphs. In Section 2 we saw that Theorems 1.1, 1.2 and 1.3 are asymptotically best possible for all complete k -partite k -graphs; we now consider those incomplete k -partite k -graphs for which the same is true. For brevity, throughout the following discussion the words ‘best possible’ should be interpreted as meaning best possible up to an $o(n)$ error term.

Recall that Proposition 2.3 showed that Theorem 1.2 is best possible for all k -partite k -graphs K of type 1 with $\tau(K) = \sigma(K)$, and also for all k -partite k -graphs K of type $d \geq 2$

with $\tau(K) = \sigma(K) \geq 1/p$, where p is the smallest prime factor of d . In particular, any k -partite k -graph on b vertices which contains a matching of size $\sigma(K)b$ (that is, a matching which covers an entire vertex class of some k -partite realisation) has $\sigma(K) = \tau(K)$.

However, there are k -partite k -graphs for which $\tau(K) \neq \sigma(K)$. For an example, choose disjoint sets U_1, \dots, U_k each of size greater than k , and for each $j \in [k]$ choose a marked vertex $u_j \in U_j$. Let K^* be the k -partite k -graph with vertex classes U_1, \dots, U_k whose edges are all k -tuples of vertices including at least one of the marked vertices u_j . Then it is not hard to see that K^* has only one k -partite realisation up to permutations of the vertex classes U_1, \dots, U_k , so $\sigma(K^*) = \min_{j \in [k]} |U_j|/b > k/b$, where $b := |V(K^*)|$. However, $\{u_j : j \in [k]\}$ is a vertex cover of K^* , so we have $\tau(K^*) \leq k/b < \sigma(K^*)$. Having seen the construction of Proposition 2.3, it is natural to ask whether the best possible versions of Theorems 1.2 and 1.3 would have $\tau(K)$ in place of $\sigma(K)$, providing an upper bound to match the lower bound of Proposition 2.3. However, we can show this is not the case by considering the k -graph K^* defined above and the following variation of the construction of Proposition 2.3. Take disjoint empty vertex sets A and B with sizes $|A| = \lceil (k+1)n/b \rceil - 1$ and $|B| = n - |A|$, and let H be the 3-graph on vertex set $A \cup B$ whose edges are all 3-tuples $e \in \binom{A \cup B}{3}$ with $1 \leq |e \cap A| \leq k-1$. Then $\delta(H) = |A| = \lceil (k+1)n/b \rceil - 1$, but H does not have a perfect K^* -packing. This is because for any copy of K^* in H , $A \cap V(K^*)$ is a vertex cover of K^* . However, we cannot have $u_j \in A$ for every $j \in [k]$ because $\{u_1, \dots, u_k\}$ is an edge of K^* . Any other vertex cover of K^* has size at least $k+1$, so any copy of K^* in H must have at least $k+1$ vertices in A . It follows that any K^* -packing in H has size at most $|A|/(k+1) < n/b$, so is not perfect. It is therefore not obvious what should be the general rule for the behaviour of $\delta(K, n)$ for k -partite k -graphs K of type 1 which have $\tau(K) \neq \sigma(K)$.

We now consider the divisibility-based extremal constructions, for which we make the following definitions. Let K be a k -partite k -graph; then we say that two vertices u, v of K are *tightly-linked* if there is a set S of $k-1$ vertices of K such that $u \cup \{S\}$ and $v \cup \{S\}$ are both edges of K . Observe that u and v must then lie in the same vertex class of any k -partite realisation of K . Now let U_1, \dots, U_k be the vertex classes of a k -partite realisation of K , and form a graph on $V(K)$ by joining any tightly-linked pair of vertices with an edge. So each connected component of this graph is a subset of a vertex class of K ; we say that K is *tightly k -partite* if the connected components of this graph are precisely the vertex classes of K . So, for example, any complete k -partite k -graph is tightly k -partite. Note that if K is tightly k -partite then it has only one k -partite realisation up to permutation of the vertex classes (but the converse does not hold: for example, C_3^3 has only one k -partite realisation but is not tightly k -partite).

Recall that Proposition 2.1 showed that Theorem 1.1 is best possible for any k -partite k -graph K which satisfies property (P1) for some p , and that Theorem 1.3 is best possible for any k -partite k -graph H which satisfies property (P2) for some $p \geq 2$ with $1/p \geq \sigma(H)$. In Section 2 we showed that property (P1) holds for any complete k -partite k -graph of type 0; the same argument shows that property (P1) holds for any tightly k -partite k -graph K of type 0. Indeed, for any $i \in [k]$ and any $x, y \in U_i$, the fact that H is tightly k -partite implies that there is a sequence x, z_1, \dots, z_ℓ, y such that each consecutive pair is tightly-linked in K , and so, given a set $A \subseteq V(K)$ such that $|e \cap A|$ is divisible by p for any $e \in K$, the vertices x, z_1, \dots, z_ℓ, y are either all in A or all not in A . Property (P1) then follows exactly as before. Similarly, in Section 2 we showed that property (P2) holds for any complete k -partite k -graph of type $d \geq 2$ and any p which divides d . Exactly as before we can adapt this argument to show that the same statement holds with ‘tightly k -partite’ in place of ‘complete k -partite’. In conclusion, we find that Theorem 1.1 is best possible for any tightly k -partite k -graph of

type 0, and that Theorem 1.3 is best possible for any tightly k -partite k -graph K of type $d \geq 2$ such that the smallest prime factor p of d satisfies $1/p \geq \sigma(K)$. Again, it is not clear what the general rule should be for the behaviour of $\delta(K, n)$ for k -partite k -graphs K of type 0 or $d \geq 2$ which are not tightly k -partite. One example of the latter category is the cycle C_3^3 , which has type 0 and for which Theorem 1.4 showed that $\delta(C_3^3, n) = n/3 + o(n) = \sigma(C_3^3)n + o(n)$. The principal difference between the proofs of Theorem 1.4 and Lemma 3.2 was the use of Proposition 8.2 to find copies of C_3^3 with an odd number of vertices on each side of a partition of $V(H)$; it seems likely that the value of $\delta(K, n)$ for k -partite k -graphs K of type 0 or type $d \geq 2$ which are not tightly k -partite depends on the minimum codegree required to give an analogue of Proposition 8.2, in a manner reminiscent of the results of [14] for perfect matchings.

9.2. Almost-perfect packings. For many applications, it suffices to find an almost-perfect K -packing in a k -graph H , that is, a K -packing covering all but a small number of vertices. It is natural to consider the minimum codegree needed to guarantee such a packing, and this can be obtained by small changes to our methods. First, observe that if in Proposition 2.3 we instead take the set A to have size $\lceil \tau(n - C) \rceil - 1$, where $\tau = \tau(K)$, then we obtain a k -graph G on n vertices with $\delta(G) = |A| = \lceil \tau(n - C) \rceil - 1$ in which any K -packing has size at most $|A|/\tau b < (n - C)/b$ (where b is the number of vertices of K). So any K -packing in G leaves more than C vertices uncovered, proving the following proposition.

Proposition 9.1. *Let K be a k -partite k -graph. Then for any $C > 0$ and any n there exists a k -graph G on n vertices with $\delta(G) \geq \lceil \tau(K)(n - C) \rceil - 1$ such that no K -packing in G covers all but at most C vertices of G .*

In the other direction, we now prove Theorem 1.5, which gives an upper bound similar to that of Theorem 1.2 but which holds for all k -partite k -graphs K (whereas Theorem 1.2 applied only when $\gcd(K) = 1$). Combined with Proposition 9.1 this gives the asymptotically correct threshold for the almost-packing problem for any k -partite k -graph K with $\tau(K) = \sigma(K)$; as we have seen, this includes all complete k -partite k -graphs.

Theorem 1.5. *Let K be a k -partite k -graph. Then there exists a constant $C = C(K)$ such that for any $\alpha > 0$ there exists $n_0 = n_0(K, \alpha)$ such that any k -graph H on $n \geq n_0$ vertices with $\delta(H) \geq \sigma(K)n + \alpha n$ admits a K -packing covering all but at most C vertices of H .*

Proof. Suppose first that $\sigma(K) < 1/k$, and fix U_1, \dots, U_k to be the vertex classes of a k -partite realisation of K in which $|U_1| = \sigma(K)b$, where $b := |V(K)|$. Note that since $\sigma(K) < 1/k$, the vertex classes U_1, \dots, U_k cannot all be the same size. Let K^* be the complete k -partite k -graph on these vertex classes; then it suffices to find a perfect K^* -packing in H . To do this, we follow the proof of Lemma 3.1 exactly as in the case when $\gcd(K) = 1$ until the step ‘Delete a K -packing to ensure divisibility of cluster sizes’, in which we applied Lemma 6.7 to delete a K^* -packing in H such that following this deletion all subclusters had size divisible by $bk \gcd(K^*)$. We cannot now apply Lemma 6.7 as we may have $\gcd(K^*) \neq 1$; however, it is straightforward to modify (and hugely simplify) the argument of Lemma 6.7 to show that we can delete a K^* -packing in H such that, following this deletion, at most one subcluster in any connected component of \mathcal{S}' does not have size divisible by $bk \gcd(K^*)$. Our condition $\delta(H) \geq \sigma(K)n + \alpha n \geq n/b + \alpha n$ implies that \mathcal{S}' has fewer than b connected components (exactly as $\delta(H) \geq n/p + \alpha n$ implied that \mathcal{S}' had fewer than p connected components). So at most b subclusters do not have size divisible by $bk \gcd(K^*)$; by deleting up to $bk \gcd(K^*)$ vertices from each of these we can ensure that every subcluster has size divisible by $bk \gcd(K^*)$. Following the remainder of the proof exactly as before, we obtain a K^* -packing in H which

covers all vertices except for the at most $b^2k \gcd(K^*) \leq b^3k$ deleted vertices, so we may take $C = b^3k$.

The remaining possibility is that $\sigma(K) = 1/k$, in which case we have $\delta(H) \geq n/k + \alpha n$. Since $\sigma(K) = 1/k$ every vertex class of any k -partite realisation of K must have equal size, so we may assume without loss of generality that K is the complete k -partite k -graph with vertex classes each of size b_1 , for some positive integer b_1 . In this case we mimic the proof of Lemma 3.2, and as in the proof of Theorem 1.4, all steps proceed as before except for the choice of the K -packing M^3 . Using the notation of the proof of Lemma 3.2, we proceed as in the proof of Theorem 1.4 to find M^3 , first obtaining a partition of $[m']$ into parts V_1, \dots, V_s for some $s < k$ such that for any $j \in [s]$ we have $|V_j| \geq m'/k + \alpha m'/2k$ and the property that, even after the deletion of up to $\alpha m'/2k$ vertices of $\mathcal{S}[V_j]$, the remaining subgraph contains a short path between any two vertices. We cannot then obtain M^* as in Claim 8.3, since we now have no analogue of Proposition 8.2, but instead we can simply delete up to $2kb$ vertices of $\bigcup_{i \in V_j} Y'_i$ for each $j \in [s]$ before proceeding similarly as the proof of Claim 8.3 to obtain a K -packing M^* such that the sets Y_i^* of vertices of Y'_i which were not deleted or covered by M^* have the desired property, that is, that the average size of sets Y_i^* with $i \in V_j$ is the same for each $j \in [s]$, and divisible by b_1 . From this we obtain M^3 exactly as in the proof of Theorem 1.4, and the final step of the proof then proceeds exactly as for Lemma 3.2, giving a K -packing in H covering all vertices except for the at most $2ksb < k^2b$ deleted vertices, so we may take $C = k^2b$. \square

9.3. Non k -partite k -graphs. All of the results of this paper pertain to k -partite k -graphs, but we can also consider the value of $\delta(H, n)$ for k -graphs H which are not k -partite. However, for most such k -graphs H we do not even know the asymptotic value of the Turán density, that is, the number of edges needed in a large k -graph G to guarantee even a single copy of H in G . By contrast, in this paper we used several times the fact that the Turán density of a k -partite k -graph is zero. A lack of knowledge of the Turán density of H is not an essential obstacle to finding $\delta(H, n)$; indeed, Keevash and Mycroft determined the exact value of $\delta(K_4^3, n)$ for large n even though finding the Turán density of K_4^3 remains a significant open problem. It would be interesting to know whether $\delta(H, n)$ is determined by the parameters $\gcd(H)$ and $\sigma(H)$ (whose definitions extend naturally to the non k -partite case), in the same way as for graphs. However, we note that it is not sufficient to consider only the smallest r for which a k -graph H admits an r -partite realisation. Indeed $K_4^3 - e$ and K_4^3 both admit a 4-partite realisation with one vertex in each vertex class, and no 3-partite realisation, but as we saw in Section 1, results of Lo and Markström [26, 27] and of Keevash and Mycroft [16] show that $\delta(K_4^3 - e, n)$ and $\delta(K_4^3, n)$ differ by $n/4 + o(n)$.

A natural starting point to consider would be the complete 4-partite 3-graph K with vertex classes of size b_1, b_2, b_3 and b_4 (so the edges are any triple whose vertices lie in three different vertex classes). The fact that $\delta(K_4^3, n) \leq 3n/4$ strongly suggests that we have $\delta(K, n) \leq 3n/4 + o(n)$ for any values of b_1, b_2, b_3 and b_4 . On the other hand, if $\gcd(\{b_1, b_2, b_3, b_4\}) > 1$, then we can modify a construction of Pikhurko [29] to show that $\delta(K, n) \geq 3n/4 - 2$, and in fact the same construction gives the same threshold if $\gcd(K) = 2$, where $\gcd(K) := \gcd(\{b_i - b_j : i, j \in [4]\}) > 1$. So we expect that $\delta(K, n) = 3n/4 + o(1)$ in these cases. If $\delta(K, n)$ exhibits behaviour analogous to that of k -partite k -graph case, then we would also expect $\delta(K, n)$ to be lower in the case $\gcd(K) = 1$. Assume without loss of generality that $b_1 \leq b_2, b_3, b_4$ and define $\sigma = \sigma(K) := \frac{b_1}{b_1 + b_2 + b_3 + b_4}$. Given disjoint sets A and B with $|A| = \lceil \sigma n \rceil - 1$ and $|B| = n - |A|$, a well-known random construction due to Czygrinow and Nagle [5] gives a k -graph G' on $|B|$ with $\delta(G') \geq |B|/2 - o(n)$ which does not contain any copy of K_4^3 . Form

a k -graph G on $A \cup B$ whose edges are the edges of G' together with any triple of vertices which intersects A . Then $\delta(G) = \delta(G') + |A| = n/2 + |A|/2 - o(n) = (1/2 + \sigma)n - o(n)$, and G cannot contain a perfect K -packing, as any copy of K in G must have at least b_1 vertices in A . So we have $\delta(K, n) \geq (1/2 + \sigma)n - o(n)$; it would be interesting to know whether this is the asymptotically correct threshold for $\gcd(K) = 1$. However, demonstrating that this is the correct threshold would imply that the minimum codegree which guarantees the existence of a copy of K_4^3 in a k -graph on n vertices is asymptotically $n/2$, so would require the solution of a well-known open problem (see [5] for further details).

10. ACKNOWLEDGEMENTS

I would like to thank Katherine Staden for her helpful comments on an earlier version of this manuscript. I also thank an anonymous referee for many helpful remarks.

REFERENCES

- [1] R. Aharoni, A. Georgakopoulos and P. Sprüssel, Perfect matchings in r -partite r -graphs, *European Journal of Combinatorics* **30** (2009), 39–42.
- [2] N. Alon, P. Frankl, H. Huang, V. Rödl, A. Ruciński and B. Sudakov, Large matchings in uniform hypergraphs and the conjectures of Erdős and Samuels, *Journal of Combinatorial Theory, Series A* **119** (2012), 1200–1215.
- [3] N. Alon and R. Yuster, H -factors in dense graphs, *Journal of Combinatorial Theory, Series B* **66** (1996), 269–282.
- [4] A. Czygrinow, L. DeBiasio and B. Nagle, Tiling 3-uniform hypergraphs with $K_4^3 - 2e$, *Journal of Graph Theory* **75** (2014), 124–136.
- [5] A. Czygrinow and B. Nagle, A note on codegree problems for hypergraphs, *Bulletin of the ICA* **32** (2001), 63–69.
- [6] A. Czygrinow, V. Kamat, Tight co-degree condition for perfect matchings in 4-graphs, *Electronic Journal of Combinatorics* **19** (2012), P20.
- [7] D. E. Daykin and R. Häggkvist, Degrees giving independent edges in a hypergraph, *Bulletin of the Australian Mathematical Society* **23** (1981), 103–109.
- [8] P. Erdős, On extremal problems of graphs and generalized graphs, *Israel Journal of Mathematics* **2** (1964), 183–190.
- [9] W.T. Gowers, Hypergraph Regularity and the multidimensional Szemerédi Theorem, *Annals of Mathematics* **166** (2007), 897–946.
- [10] A. Hajnal and E. Szemerédi, Proof of a conjecture of Erdős, *Combinatorial Theory and its Applications, Vol.2* (Ed. P. Erdős, A. Rényi and V.T. Sós), Amsterdam, Netherlands: North-Holland, pp. 601–623, 1970.
- [11] H. Hàn, Y. Person, M. Schacht, On perfect matchings in uniform hypergraphs with large minimum vertex degree, *SIAM Journal on Discrete Mathematics* **23** (2009), 732–748.
- [12] J. Han, Y. Zhao, Minimum vertex degree threshold for C_4^3 -tiling, *Journal of Graph Theory* **79** (2015), 300–317.
- [13] P. Keevash, A hypergraph blowup lemma, *Random Structures & Algorithms* **39** (2011), 275–376.
- [14] P. Keevash, F. Knox, R. Mycroft, Polynomial-time perfect matchings in dense hypergraphs, *Advances in Mathematics* **269** (2014), 265–334.
- [15] P. Keevash, D. Kühn, R. Mycroft and D. Osthus, Loose Hamilton cycles in hypergraphs, *Discrete Mathematics* **311** (2011), 544–559.
- [16] P. Keevash and R. Mycroft, A geometric theory of hypergraph matching, *Memoirs of the American Mathematical Society* **233** (2014), monograph 1098.
- [17] I. Khan, Perfect matching in 3-uniform hypergraphs with large vertex degree, *SIAM Journal of Discrete Mathematics* **27** (2013), 1021–1039.
- [18] I. Khan, Perfect Matchings in 4-uniform hypergraphs, preprint, arXiv:1101.5675.
- [19] J. Komlós, Tiling Turán theorems, *Combinatorica* **20** (2000), 203–218.
- [20] J. Komlós, G.N. Sárközy and E. Szemerédi, Proof of the Alon-Yuster conjecture, *Discrete Mathematics* **235** (2001), 255–269.
- [21] D. Kühn and D. Osthus, Loose Hamilton cycles in 3-uniform hypergraphs of high minimum degree, *Journal of Combinatorial Theory, Series B* **96** (2006), 767–821.

- [22] D. Kühn and D. Osthus, The minimum degree threshold for perfect graph packings, *Combinatorica* **29** (2009), 65–107.
- [23] D. Kühn, D. Osthus and A. Treglown, Matchings in 3-uniform hypergraphs, *Journal of Combinatorial Theory, Series B* **103** (2013), 291–305.
- [24] J. Lenz, D. Mubayi, Perfect packings in quasirandom hypergraphs, *Journal of Combinatorial Theory, Series B*, to appear.
- [25] A. Lo, K. Markström, Minimum codegree threshold for $(K_4^3 - e)$ -factors, *Journal of Combinatorial Theory, Series A* **120** (2013), 708–721.
- [26] A. Lo and K. Markström, Perfect matchings in 3-partite 3-uniform hypergraphs, *Journal of Combinatorial Theory, Series A* **127** (2014), 22–57.
- [27] A. Lo and K. Markström, F -factors in hypergraphs via absorption, *Graphs and Combinatorics* **31** (2015), 679–712.
- [28] K. Markström and A. Ruciński, Perfect matchings (and Hamilton cycles) in hypergraphs with large degrees, *European Journal of Combinatorics* **32** (2011), 677–687.
- [29] O. Pikhurko, Perfect matchings and K_4^3 -tilings in hypergraphs of large codegree, *Graphs and Combinatorics* **24** (2008), 391–404.
- [30] V. Rödl and A. Ruciński, Dirac-type questions for hypergraphs – a survey (or more problems for Endre to solve), An Irregular Mind (Szemerédi is 70), *Bolyai Society Mathematical Studies* **21** (2010), 561–590.
- [31] V. Rödl, A. Ruciński and E. Szemerédi, Perfect matchings in large uniform hypergraphs with large minimum collective degree, *Journal of Combinatorial Theory, Series A* **116** (2009), 616–636.
- [32] V. Rödl and M. Schacht, Regular partitions of hypergraphs: regularity lemmas, *Combinatorics, Probability and Computing* **16** (2007), 833–885.
- [33] V. Rödl and M. Schacht, Regular partitions of hypergraphs: counting lemmas, *Combinatorics, Probability and Computing* **16** (2007), 887–901.
- [34] V. Rödl and J. Skokan, Regularity lemma for uniform hypergraphs, *Random Structures & Algorithms* **25** (2004), 1–42.
- [35] A. Treglown and Y. Zhao, Exact minimum degree thresholds for perfect matchings in uniform hypergraphs, *Journal of Combinatorial Theory, Series A* **119** (2012), 1500–1522.
- [36] A. Treglown and Y. Zhao, Exact minimum degree thresholds for perfect matchings in uniform hypergraphs II, *Journal of Combinatorial Theory, Series A* **120** (2013), 1463–1482.

Richard Mycroft,
 School of Mathematics,
 University of Birmingham,
 Birmingham B15 2TT,
 United Kingdom.
 r.mycroft@bham.ac.uk