

Impact of habituated dietary protein intake on fasting and postprandial whole-body protein turnover and splanchnic amino acid metabolism in elderly men

Højfeldt, Grith; Bülow, Jacob; Agergaard, Jakob; Asmar, Ali; Schjerling, Peter; Simonsen, Lene; Bülow, Jens; Van hall, Gerrit; Holm, Lars

DOI:
[10.1093/ajcn/nqaa201](https://doi.org/10.1093/ajcn/nqaa201)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Højfeldt, G, Bülow, J, Agergaard, J, Asmar, A, Schjerling, P, Simonsen, L, Bülow, J, Van hall, G & Holm, L 2020, 'Impact of habituated dietary protein intake on fasting and postprandial whole-body protein turnover and splanchnic amino acid metabolism in elderly men: a randomized, controlled, crossover trial', *The American journal of clinical nutrition*, vol. 112, no. 6, pp. 1468-1484. <https://doi.org/10.1093/ajcn/nqaa201>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is a pre-copyedited, author-produced version of an article accepted for publication in *The American Journal of Clinical Nutrition* following peer review. The version of record [Grith Højfeldt, Jacob Bülow, Jakob Agergaard, Ali Asmar, Peter Schjerling, Lene Simonsen, Jens Bülow, Gerrit van Hall, Lars Holm, Impact of habituated dietary protein intake on fasting and postprandial whole-body protein turnover and splanchnic amino acid metabolism in elderly men: a randomized, controlled, crossover trial, *The American Journal of Clinical Nutrition*, Volume 112, Issue 6, December 2020, Pages 1468–1484] is available online at: <https://doi.org/10.1093/ajcn/nqaa201>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Impact of habituated dietary protein intake on fasting and postprandial whole body protein turnover and splanchnic amino acid metabolism in elderly men: a randomized controlled, crossover trial.

Grith Højfeldt¹, Jacob Bülow¹, Jakob Agergaard¹, Ali Asmar^{2,3}, Peter Schjerling^{1,4}, Lene Simonsen², Jens Bülow^{2,6}, Gerrit van Hall^{5,6}, Lars Holm^{1,6,7}

¹Institute of Sports Medicine, Department of Orthopedic Surgery M81, Bispebjerg Hospital, Copenhagen, Denmark (GH, JaB, JA, PS, LH)

²Department of Clinical Physiology and Nuclear Medicine, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark (AA, LS, JB)

³Department of Clinical Physiology, Nuclear Medicine and PET, Rigshospitalet, Copenhagen, Denmark (AA)

⁴Center for Healthy Aging, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark (PS)

⁵Clinical Metabolomics Core Facility, Clinical Biochemistry, Rigshospitalet, Copenhagen, Denmark (GvH)

⁶Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark (GvH, LH)

⁷School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK (LH)

Corresponding Author: Grith Højfeldt, address: Bispebjerg Hospital, Indgang 8, 1. sal, Nielsine Nielsens Vej 11, 2400 København NV, phone: +45 38635042 e-mail: grithwh@gmail.com

Sources of supports: The Danish Dairy Research Foundation; Arla Foods Ingredients Group P/S; The Excellence Programme 2016 University of Copenhagen – project CALM. We disclose that

the funding entities have neither been taking part in the study design, trial conduction, sample analyses nor data interpretation in the present study.

Running head: Habituated protein intake & whole body turnover.

List of abbreviations: Body weight (BW), fractional synthesis rate (FSR), high protein (HP), lean body mass (LBM), net balance (NB), plasma flow (PF), partial volume of distribution (pV), rate of appearance (R_a), rate of disappearance (R_d), recommended dietary allowance (RDA), recommended protein (RP).

Clinical Trial Registry: Journal number NCT02587156, [Clinicaltrials.org](https://clinicaltrials.org).

Data described in the manuscript, code book, and analytic code will be made available upon request pending application and approval.

1 **Abstract**

2 *Background:* Efficacy of protein absorption and subsequent amino acid utilization may be
3 reduced in elderly. Higher protein intakes have been suggested to counteract for this.

4 *Objective:* To elucidate how habituated level of protein intake affects the fasted state and the
5 stimulatory effect of a protein-rich meal on protein absorption, whole body protein turnover
6 and splanchnic amino acid metabolism.

7 *Design:* 12 males (65-70 years) were included in a double-blinded crossover intervention
8 study, consisting of a 20-day habituation period to a protein intake at recommended dietary
9 allowance (RDA) or high-level (1.1 g/kg lean body mass (LBM)/day or >2.1 g/kg LBM/day),
10 each followed by an experimental trial with a primed, constant infusion of D₈-phenylalanine
11 and D₂-tyrosine. Arterial and hepatic venous blood samples were obtained after an overnight
12 fast and repeatedly four hours after a standardized meal including intrinsically labeled whey
13 protein concentrate and calcium-caseinate proteins. Blood was analyzed for amino acid
14 concentrations and phenylalanine and tyrosine tracer enrichments wherefrom whole-body and
15 splanchnic amino acid and protein kinetics were calculated.

16 *Results:* High compared to the recommended level of protein intake resulted in a higher fasting
17 whole body protein turnover with a resultant 0.03 (± 0.01 SEM) $\mu\text{mol/kg LBM/min}$ lower net
18 balance ($P < 0.05$), which was not rescued by the intake of a protein dense meal. The plasma
19 protein fractional synthesis rate was 0.13 (± 0.06 SEM) %/h lower ($P < 0.05$) following
20 habituation to high protein. Further a higher fasting and postprandial amino acid removal was
21 observed following habituation to high protein, yielding higher urea excretion and increased
22 phenylalanine oxidation rates ($P < 0.01$).

23 *Conclusion:* Three weeks of habituation to high protein intake (>2.1 g protein/kg LBM/day)
24 led to a significantly higher net protein loss in the fasted state. This was not compensated for in
25 the 4-hour -post-prandial period after intake of a meal high in protein.

26 *Keywords:* Habitual protein intake, recommended protein intake, protein turnover, protein
27 breakdown, whole body protein turnover, intrinsically labelled proteins, stable-isotope tracers,
28 whey protein, caseinate protein

29

30

31

32

33

34

35

36

37

38

39

40

41

42 **Introduction**

43 The recommended daily requirement of 0.83 g dietary protein/kg body weight/day (1–3) for
44 supporting lean mass maintenance in all adults have repeatedly been challenged and suggested
45 to be elevated for individuals above 65 years of age (4). The basis for this awareness arises
46 from follow-up and observational cross-sectional studies (5–7). A limitation of these findings
47 is that differences most often are found between moderately high and lower than
48 recommended dietary allowance (RDA) protein intakes (5,6,8). Suggestions made to elevate
49 the protein intake above the current recommendations for the elderly assume that an additive
50 effect of protein intake will also be seen between recommended and higher protein intakes.
51 This assumption has not been tested.

52 One cause for an enhanced protein requirement for elderly individuals is anabolic resistance
53 (9–11). The mechanism for which is unknown but may be found in the route from oral intake
54 of protein to peripheral efficacy to enhance protein synthesis and or reduce protein breakdown.
55 The first pass splanchnic tissues extraction has been shown to increase with age (12,13), thus
56 following a protein containing meal, less amino acids is presented to the circulation and made
57 available to promote protein synthesis. The amount of amino acids in the circulation is affected
58 by the absorption and digestion rates of proteins as well as the ability of the peripheral tissue to
59 take up amino acids from the circulation and incorporate them into body proteins and vice
60 versa to release amino acids into the circulation.

61 Metabolic homeostasis can occur within rather wide ranges of protein exposures. This is
62 reflected by e.g. increased capacity to degrade and oxidize amino acids and hence scavenge
63 nitrogen by increasing urea production and excretion when protein intake is increased (14–16).
64 Fasting whole body protein kinetics have also been shown to be enhanced after habituation to
65 divergent dietary protein levels in both younger (17,18) and older (19) individuals. In contrary,

66 Gorissen and colleagues habituated older individuals for two weeks to high (1.5 g/kg body
67 weight (BW)/d) vs. low (0.7 g/kg BW/d) protein intake and observed neither an effect on
68 fasting whole body protein synthesis and breakdown rates nor a response to plain protein
69 feeding: However, a significantly increased amino acid oxidation was observed following
70 habituation to high protein (20).

71 With reference to the suggestions to elevate the dietary protein for older adults, it remains to
72 be elucidated whether habituation to higher protein intake have an impact different from the
73 currently recommended on the whole body protein turnover in a fasted state and how it
74 affects the post-prandial protein handling .The primary outcome was the difference in plasma
75 protein enrichment from intrinsically labeled whey protein 4 hours after ingestions. This was
76 measured to assess the post-prandial response to a mixed meal following habituation for 20
77 days to a normal (RDA) and high dietary protein intake in older individuals. Secondary
78 outcomes included whole body and plasma specific protein metabolism both in a fasted and
79 post-prandial state. We hypothesized that a high protein diet would increase the amino acid
80 oxidation and the nitrogen removal apparatus but simultaneously improve digestion/absorption
81 driving a more positive postprandial net protein synthesis response.

82

83

84

85

86

87

88 **Subjects, materials and methods**

89 *Participants*

90 Twelve men (65-70 years) completed this double-blinded randomized crossover trial (**Figure**
91 **1**). Before inclusion, all participants were screened by a physician and deemed overall healthy
92 based on blood samples, medical history and an interview. The exclusion criteria were
93 diabetes; pain in the skeletal muscle, connective tissue or back; known arthrosclerosis; alcohol
94 intake above 21 units per week; and regular intake of drugs or dietary supplements anticipated
95 to affect body protein turnover. Figure 1 shows a flowchart of the enrollment. 69 males were
96 assessed for eligibility, 12 were included and went through both habituation periods. There
97 were no dropouts.

98 Study design, purpose, risks and discomforts involved were explained (written and orally) to
99 each participant in accordance with the Helsinki declaration, after which the subjects gave
100 written consent of participation. The study took place at Bispebjerg and Frederiksberg
101 Hospital, Copenhagen. The local ethical committee of the Capital Region of Denmark
102 approved the study, protocol number H-15005598 and the study was registered at
103 Clinicaltrials.org under journal number NCT02587156.

104 Pre-screening involved dual-energy X-ray absorptiometry (DXA) scanning (Lunar iDXA; GE
105 Medical Systems, Pewaukee, WI, USA, with enCORE v.16 software) to determine the body
106 composition of the participants, with emphasis on the lean body mass (LBM) used to calculate
107 the desired dietary protein intake and the tracer amount needed. Participant characteristics can
108 be seen in **Table 1**.

109

110 *Experimental design*

111 The experimental design is shown in **Figure 2**.

112 The crossover intervention consisted of 20 days habituation to the recommended level of
113 protein in one period (recommended protein, RP) and habituation to a high level of protein
114 intake in another period (high protein, HP). The order of the habituation periods was randomly
115 assigned by a draw in a double blinded fashion.

116 In the 20-day habituation period the participants received four daily intakes of identically
117 packed supplements containing 20 g of whey hydrolysate and 10 g of sucrose in the HP period
118 and an isocaloric identically packed supplement consisting of 20 g maltodextrin and 10 g
119 sucrose in the RP period. The supplements came in powder form and the participants were
120 instructed to dissolve them a cold beverage of their choosing. The four daily supplements were
121 spread out during the day based on the participants customary protein distribution (e.g. two
122 supplements with breakfast, one with lunch and one with dinner if the participant ingested least
123 protein at breakfast in his daily diet. The pattern of supplement intake was decided by the
124 investigator) aiming at an even protein intake at all 3 main meals in the HP period. In the RP
125 period, participants were instructed in the exact same manner, however as their supplement
126 contained carbohydrate, their protein distribution throughout the day was not even.

127 Each habituation period started with a start-up period of seven days, where the only dietary
128 alterations were the four daily supplements. Subjects were guided towards a basic diet of 1.1 g
129 protein/kg LBM/day based on a 3-day dietary registration during the seven start up days. The
130 remaining 13 days of the 20-day habituation period were controlled by regular online dietary
131 registrations. Protein intake was adjusted to lean body mass instead of total body weight to
132 account for varying body compositions assuming that the lean body mass (LBM) is mainly
133 responsible for amino acid and protein metabolism. The 1.1. g protein/kg LBM/day
134 corresponds to 0.8 g protein/kg BW/day assuming a whole body fat content of 27%.

135

136 From day 19, participants refrained from any sort of strenuous physical activity. At 8 a.m. on
137 day 20, a 24-hour urine collection was started. At the last meal of day on day 20, all
138 participants had a standardized meal. The experimental trial on day 21 was identical for all
139 participants irrespective of the preceding habituation period. Following day 21, the participants
140 had a washout period of minimum 45 days, before crossing over to the alternate protein intake
141 intervention in a double blinded fashion. An overview of the 21-day period is illustrated in
142 figure 2.

143

144 *Production of intrinsically labeled proteins (given on day 21)*

145 Five Danish Holstein Friesian cows were infused with stable isotopes, four with L-[ring-D₅]-
146 phenylalanine and 1 cow with L-[¹⁵N]-phenylalanine, the protocol is described in detail
147 elsewhere (21). In brief, milk was collected from 11 milkings, pasteurized and caseinate and
148 whey proteins were purified, yielding L-[ring-D₅]-phenylalanine labeled whey with an
149 enrichment of 15.44 ± 0.24 mole percent excess (MPE) (\pm SEM) and L-[¹⁵N]-phenylalanine
150 labeled caseinate with an enrichment of 20.81 ± 0.02 MPE (\pm SEM).

151

152 *Day 21 – experimental trial day*

153 Participants fasted for 12 h overnight and arrived at 8 a.m. by taxi to the hospital ward. Upon
154 arrival they finished their first 24-hour urine sample period and a second 24-hour urine sample
155 period was started. Hereafter, the participants changed into hospital attire and were placed in a
156 bed in the supine position and a catheter was inserted in an antecubital vein and a background
157 blood sample was taken (-100 min, end of meal intake is set to 0 min). Hereafter, primed
158 continuous infusions were started of D₈-phenylalanine (prime: 4 μ mol/kg LBM, continuous: 4
159 μ mol/kg LBM/h in a fasted state, 4.8 μ mol/kg LBM/h in postprandial state), D₂-tyrosine
160 (prime: 2.2 μ mol /kg BW, continuous: 2.3 μ mol/kg BW/h in a fasted state, 2.75 μ mol/kg

161 LBM/h in a postprandial state), $^{15}\text{N}_2$ -urea (prime 84 $\mu\text{mol/kg}$ BW, continuous 9 $\mu\text{mol/kg}$
162 BW/h) and indocyanine green (ICG, prime: 1 mg, continuous infusion: 7-14 mg/h). At -40
163 minutes a catheter was placed in a radial artery under local anesthesia. Fluoroscopy was used
164 to place a catheter in a hepatic vein (through vena femoralis). The radial artery catheter as well
165 as the liver catheter were used for blood collection which were taken simultaneously. The
166 catheters were kept clear by frequent flushing with saline.

167 No sooner than 90 minutes after the start of the stable isotope tracer infusion, two fasting blood
168 samples were taken from each of the sampling sites (hepatic vein and artery). The consecutive
169 blood samples were taken at -12 and -10 minutes, analysis of these results were pooled and
170 collectively referred to as -10. Following blood sampling, a protein dense mixed breakfast
171 meal was served. The meal had to be ingested within 10 minutes and contained 0.61 g
172 protein/kg LBM, of which 0.23 g/kg LBM came from ^{15}N -phenylalanine intrinsically labeled
173 caseinate protein, and 0.31 g/kg LBM came from D_5 -phenylalanine intrinsically labeled whey
174 protein. The D_5 -labeled whey protein was dissolved in water and served as a protein drink. The
175 ^{15}N -labeled caseinate was mixed with raspberry jam, which was spread with butter on a bun
176 and served with the whey drink. The protein content in the butter and the bun constituted the
177 remaining 0.07 g/kg LBM of the total 0.61 g protein /kg LBM. **Table 2** shows the
178 macronutrient content in the breakfast.

179

180 After meal ingestions, blood samples were drawn simultaneous from the hepatic vein and
181 radial arteria at time 30, 60, 90, 120, 150, 180, and 240 min. After the last sample was
182 obtained, all catheters were removed, participants received a standardized lunch containing
183 0.80 g protein/kg LBM and a urine sample was collected, and the participants got a taxi home.
184 At home they ingested a standardized dinner ensuring an equal protein content of 0.61 g /kg
185 LBM. Thus, all participants ingested 2.02 g protein/kg LBM across day 21 independent of

186 their habituated protein intake. Next morning at 8 a.m. and prior to food intake, the second 24-
187 hour urine collection ended.

188

189 *Blood analysis*

190 Arterial plasma samples were converted to serum by use of 1 unit thrombin pr. 200 μ L EDTA
191 plasma. The serum samples were analyzed for insulin, using ALPCO Insulin ELISA assay
192 (Catalog number: 80-INSHU-E01.1, E10.1, ALPCO).

193 Free amino acids were extracted from arterial plasma for measures of amino acid
194 concentrations and amino acid tracer abundances. An internal standard in a 500 μ L 50% acetic
195 acid solution was added to 100 μ L plasma, following which plasma was poured over cation
196 exchange columns with resin (AG 50W-X8 resin, Bio-Rad laboratories, Hercules, Ca, USA),
197 which had been prepped by adding 3 x 2 ml 1M HCl creating an acidic environment. The
198 columns were washed 5 times with 3 mL of deionized water before the amino acids were
199 eluted by adding 2 x 2 mL 4M NH_4OH collected in vials. Solvent was evaporated under a
200 stream of N_2 flow at 70°C and derivatized using PITC derivatization agent, converting the
201 samples into their phenylthiocabamyl (PITC) derivative. 10 μ L of the 100 μ L derivatized
202 samples were loaded and analyzed on an ultra performance liquid chromatography system
203 coupled to a triple stage quadrupole mass spectrometer (LC-MS/MS) (Thermo Fischer
204 Scientific, San Jose, CA, USA) as described by Bornø et al. (22). Plasma protein bound tracer
205 enrichments, were analyzed from plasma protein precipitated with 500 μ L ice cold acetone pr
206 50 μ L plasma, hydrolyzed by 1 ml 1 M HCl and 1 ml resin slurry, and left overnight at 110°C.
207 Following hydrolysis, the samples were purified over cation exchange resin, as described
208 above. For measures of deuterium enrichments, samples were PITC derivatized and run on LC-
209 MS/MS system as described above. For determining the ^{15}N abundances, amino acids were
210 converted to the N-acetyl-propyl (NAP) derivatives and analyzed on a gas chromatography-

211 combustion-isotope ratio mass spectrometer (GC-C-IRMS) system as previously described by
 212 Bornø et al. (23).

213 Plasma and urine samples for analysis of urea concentration as well as urea tracer enrichments
 214 were prepared with internal standards and run over resin columns just like the samples for
 215 phenylalanine and tyrosine enrichments. Following cation exchange on the resin columns and
 216 overnight drying under N₂ stream the samples were derivatized by adding 40 µl acetonitrile
 217 and 40 µl N-methyl-N-(tert-butyldimethylsilyl) trifluoroacetamide (MtBSTFA) + 1% tert-
 218 butyl-dimethylchlorosilane (tBDMCS), followed by vortex mixing and kept at 70°C for 30
 219 minutes. The derivatized samples were separated on a GC column and isotope ratios were
 220 analyzed on a triple-stage quadrupole mass spectrometer.

221

222 *Splanchnic flow*

223 The splanchnic blood flow was measured via Fick's principle, using a primed continuous
 224 infusion of indocyanine green (ICG) as indicator as previously described (24). The plasma ICG
 225 concentration was determined by spectrophotometry at wavelengths 805 and 900 nm. The light
 226 absorption at 900 nm is a measure of turbidity and is thus regarded as the background "noise".
 227 The calculation of blood flow is based on the difference in ICG concentration found in the
 228 artery and hepatic venous blood. Thus, the splanchnic plasma flow is calculated using Fick's
 229 principle (25).

$$\text{Splanchnic plasma flow} = \frac{\text{ICG influx}}{C_{\text{ICG-a}} - C_{\text{ICG-hv}}}$$

230 Where C_{ICG-a}=ICG concentrations in the artery, and C_{ICG-hv}=ICG concentrations in the hepatic
 231 vein.

232

233 *Tracers*

234 Concentrations of the labeled phenylalanine and tyrosine are calculated as their tracer to tracee
 235 ratio (TTR) multiplied by the concentrations of unlabeled compounds. Enrichments are given
 236 as mole percent excess ($MPE = TTR / (1 + TTR)$). For D₈-phenylalanine enrichments the
 237 transamination product (D₇-phenylalanine), and oxidation products (D₇- and D₆-tyrosine) were
 238 also measured and the sum D₈- and D₇-phenylalanine as well as D₇- and D₆-tyrosine were used
 239 for all calculations.

240 In the postprandial period, the tracers are in non-steady state and all amino acid kinetics in this
 241 period are based on both ingestion of D₅-phenylalanine labeled whey, ¹⁵N-phenylalanine
 242 labeled caseinate, continuously infused D₈-phenylalanine and continuously infused D₂-
 243 tyrosine. The calculations are done using Steele equation (26) with modification introduced by
 244 Proietto et al. (27). An approach which is generally accepted for calculating whole body amino
 245 acid and protein kinetics (20,28–30). The Steele equation incorporates the change in pool size
 246 defined as: $Pool\ size = pV * \frac{dC}{dt}$, where pV is the partial volume of distribution set at 0.125,
 247 dC is the difference in concentrations [μ mol/l] across the time interval, dt. This means that for
 248 all calculations which include non-steady state conditions, the values are calculated as delta
 249 values between timepoints, thus the time resolution for these will be -5, 15, 45, 75, 105, 135,
 250 165 and 210 minutes.

251

252 Plasma protein FSR: The direct incorporation of amino acids into plasma protein was given by:

$$FSR = \left(\frac{\Delta E\ protein\ [MPE]}{E\ Precursor\ pool\ [MPE] * \Delta time\ [h]} \right) * 100\%$$

253 Where ΔE protein is the change in enrichment in plasma protein, and the precursor is the
 254 enrichment in the plasma taken from the hepatic vein.

255

256 Ingested amino acid appearance rate: The exogenous rate of appearance (R_a), is the appearance
 257 rate into the systemic circulation of phenylalanine derived from the ingested proteins, thus
 258 phenylalanine from whey and caseinate, into the hepatic vein. In order to calculate this, the
 259 total R_a is needed. In the fasting, steady state, the total and exogenous R_a is given by:

$$Total R_a = \frac{IR}{E_{(t)}}$$

$$Exogenous R_a = \frac{Total R_a \cdot dE_{OT(t)}}{E_{protein}}$$

260 In the postprandial, non-steady state

$$Total R_a = \frac{IR}{\hat{E}_{(t)}} - \frac{pV \cdot \left[\frac{\hat{C}_{(t)}}{1 + \hat{E}_{(t)}} \right] \cdot \left[\frac{dE_{(t)}}{dt} \right]}{\hat{E}_{(t)}}$$

$$Exogenous R_a = \frac{Total R_a \cdot dE_{OT(t)} + pV \cdot dC_{(t)} \cdot \left[\frac{dE_{(t)}}{dt} \right]}{E_{protein}}$$

261 Where IR= tracer infusion rate [$\mu\text{mol/kg LBM/min}$], $E_{(t)}$ = enrichment at time t, $dE_{OT(t)}$ = delta
 262 enrichment from the oral tracer (D_5 -phenylalanine or ^{15}N -phenylalanine) at time t, $\hat{C}_{(t)}$ =
 263 average concentration between values surrounding time t, $\hat{E}_{(t)}$ =average enrichment between
 264 values surrounding time t, $dE_{(t)}$ =delta enrichment, dt =delta time and $E_{protein}$ =the enrichment in
 265 the ingested protein. All concentrations and enrichments are measured in the hepatic vein.

266

267 Net amino acid balance across the splanchnic tissues: The net balance (NB) across the
 268 splanchnic tissues is given by (31):

$$Splanchnic NB = (C_a - C_{hv}) \cdot PF$$

269 Where C_a and C_{hv} are phenylalanine concentrations measured in the artery and hepatic vein,
 270 respectively. PF is the plasma flow in the hepatic vein.

271

272 Whole body protein turnover: All measures in this section are based on concentrations and
 273 enrichments in the arterial blood. The endogenous R_a represents phenylalanine appearing from
 274 within the body, used as a measure of protein breakdown. It is given by:

$$Endogenous R_a = Total R_a - Exo R_a - IR$$

275 The formula is the same for the fasted, steady state and the postprandial, non-steady state.
 276 However, the appearance rates used are the ones calculated in the steady and non-steady state,
 277 respectively.

278 The rate limiting step in phenylalanine oxidation is the irreversible hydroxylation of
 279 phenylalanine to tyrosine (32). This conversion is given by:

280 In the fasted, steady state

$$Phe \rightarrow tyr = Total R_a \cdot \frac{E_{tyr}}{E_{phe}}$$

281 In the postprandial, non-steady state

$$Phe \rightarrow tyr = Total R_a \cdot \frac{\hat{E}_{tyr}}{\hat{E}_{phe}}$$

282 Where \hat{E}_{tyr} and \hat{E}_{phe} are the mean enrichments of either D₇-tyrosine or D₈-phenylalanine
 283 between two surrounding timepoints.

284 Subtracting the ‘phenylalanine-to-tyrosine conversion rate’ from the total Rate of
 285 disappearance (R_d), gives a measure of the phenylalanine removed by other processes than
 286 oxidations, hence a measure of synthesis. The R_d is given by:

287 In the fasted, steady state

$$Total R_d = Total R_a$$

288 In the postprandial, non-steady state

$$Total R_d = pV \cdot \frac{dC}{dt} + Total R_a$$

289 As for the endogenous R_a , the formula for the endogenous R_d is the same in the fasted steady
290 state and the postprandial non-steady state. However, the disappearance rate and the
291 conversion of phenylalanine to tyrosine used are the ones calculated in the steady and non-
292 steady state respectively. Thus, the endogenous R_d is given by:

$$\text{Endogenous } R_d = \text{Total } R_d - (\text{Phe} \rightarrow \text{tyr})$$

293

294 *Deviations*

295 All 12 subjects completed all trial days. However, on the experimental day 21, three subjects
296 did not receive infusion of D_2 -tyrosine and are not included in calculations of phenylalanine
297 conversion to tyrosine and whole-body protein synthesis measurements. One participant was
298 excluded from all postprandial measurements due to mixing error with the labeled proteins.
299 D_7 -phenylalanine and D_6 -tyrosine (deriving from transamination of D_8 -phenylalanine and
300 conversion from D_7 -phenylalanine to D_6 -tyrosine) was only measured for 5 participants. Based
301 on the known D_8 -phenylalanine/ D_7 -phenylalanine and D_7 -tyrosine/ D_6 -tyrosine ratios for these
302 five participants, the D_7 -phenylalanine and D_6 -tyrosine have been calculated for all
303 participants.

304 There were no dropouts.

305

306

307

308

309

310

311 Statistics

312 Intervention groups' responses to feeding over time were compared using two-way ANOVA
313 with repeated measures. Interaction effects were tested using SIDAK post hoc test and time
314 effects were tested using Dunnet's post hoc test. Difference between the two habituation
315 periods in the fasted state was compared using a paired t-test. The insulin data is presented as
316 the numerical values, however as data was not normally distributed the statistical analysis were
317 performed on log transformed data. The primary outcome was the difference in plasma protein
318 enrichment of D₅-phenylalanine from the intrinsically labeled whey protein 4 hours after
319 ingestions. This was studied in 12 individuals, allowing us to detect a difference of 0.01 MPE
320 with the expected standard deviation of 0.01, a significance level of 5% and a power of 80%.
321 GraphPad Prism 7.0 was used for all statistical tests. Data is presented as means ±SEM unless
322 otherwise stated. Significant level was set to $p < 0.05$. The intervention effect was assessed by
323 two-tail Student's *t*-test on within-subject dissimilarity as a response to high- and low-protein
324 intake.

325

326

327

328

329

330

331

332

333 **Results**

334 The participants were on average 66.6 year of age and were overall healthy based on normal
335 blood pressure, lipid blood profile and hemoglobin A1c. Their average protein intake (1.5 g/kg
336 LBM/day) was higher than the current recommendation. All participant characteristics are
337 shown in Table 1.

338 With an intake of 1.22 ± 0.04 g/kg LBM/day (0.82 ± 0.03 g/kg BW/day), the full 20-day
339 dietary composition for the RP period was slightly higher than the aim (1.1 g/kg LBM/day).
340 However, the last 13 days of the habituation period, were right on point. For the HP period the
341 aim was >2.1 g/kg LBM/day. The actual intake covering the full 20-day period was 24%
342 higher, with an intake of 2.61 ± 0.04 g/kg LBM/day (1.76 ± 0.04 g/kg BW/day). The protein
343 intakes in the two habituation periods are seen in **Table 3**.

344

345 *Nitrogen and amino acid losses*

346 The urine urea quantity during the last 24 hours of the habituation was significantly higher
347 following habituation to HP (**Figure 3A**, $P<0.01$), emphasizing a high compliance to the
348 dietary protocol among the participants. During the 24 hours which included the trial day all
349 participants had the exact same relative protein intake irrespective of the habituation period.
350 Urine collected during the 8 h of the trial had a similar urea quantity independent of
351 habituation period (**Figure 3B**). In contrast, the total 24 hour urine of the entire trial day 21
352 had a higher urine urea after habituation to HP (**Figure 3C**, $P<0.05$) in spite of the same
353 protein intake through the entire day.

354 In line with these findings, the urea R_a is significantly higher following habituation to HP in
355 the fasted state (**Figure 4A**, $P<0.01$). The urea R_a remained elevated following habituation to

356 HP as compared to RP throughout the 4-hour postprandial period (**Figure 4B**). However, there
357 was no difference in the absolute change from fasted to postprandial urea R_a between the two
358 habituation periods (baseline corrected; data not shown). In accordance with this, the
359 phenylalanine hydroxylation rate to tyrosine (**Figure 4C and D**) was significantly higher
360 following habituation to HP both in the fasted (Figure C, $P<0.01$) and in the postprandial four
361 hour period (Figure 4D, time: $P<0.0001$, intervention $P<0.001$).

362 *Plasma protein fractional synthesis rate and nutrient protein-derived tracer abundances*

363 The fractional synthesis rates (FSR) of plasma proteins are shown in **Figure 5A**. There was a
364 significant higher plasma FSR in the 4-hour postprandial period following the RP period
365 compared to the HP period ($P=0.046$).

366 The enrichment from the intrinsically labeled whey (D_5 -phenylalanine labeled) and caseinate
367 (^{15}N -phenylalanine labeled) proteins were assessed in the plasma protein 1 and 4 hours
368 postprandial. Values are presented as mole percent excess (MPE) for D_5 -phenylalanine and
369 atom percent excess (APE) for ^{15}N -phenylalanine enrichments. The incorporation of
370 phenylalanine deriving from both whey and caseinate was greater at 4 hours compared to 1
371 hour postprandial (**Figure 5B and C**, $P<0.0001$). For the incorporation of phenylalanine
372 deriving from the whey protein into plasma proteins there was a significantly higher
373 incorporation following RP compared to HP intake 4 hours postprandial (Figure 5C,
374 $P=0.0007$).

375

376 *Insulin concentrations*

377 Serum insulin increased with the meal intake and remained elevated from baseline in the first 3
378 hours postprandial. It was unaffected by the preceding protein habituation period both in the
379 fasted and the postprandial state (**Figure 6**).

380

381 *Phenylalanine and protein kinetics*

382 The R_a of phenylalanine from the whey and caseinate was measured in blood samples both
383 from the hepatic vein and the radial artery. While there is a higher appearance rate measured in
384 the hepatic vein, the R_a pattern is the same from the two sampling sites. There is a significant
385 time effect of the R_a of phenylalanine from the whey and the caseinate. However, no
386 differences existed between habituation periods (**Figure 7**).

387 The whole-body protein breakdown rate was significantly higher following habituation to HP
388 compared with RP ($P<0.05$) in the fasted state (**Figure 8A**). In the postprandial period, the
389 whole-body protein breakdown was significantly decreased (**Figure 8B**, time: $P<0.0001$) with
390 no difference between habituation periods. The whole-body protein synthesis was higher in the
391 fasted state following habituation to HP as compared to RP (Figure 8A, $P<0.01$). Post-
392 prandially, the whole-body protein synthesis increased (**Figure 8C**, time: $P<0.0001$) but more
393 so following habituation to HP as compared to RP ($P<0.05$). As an overall result, the whole-
394 body protein turnover rate was higher when habituated to HP compared to RP. However, and
395 most important for lean body mass maintenance, the net protein turnover balance in the fasted
396 state was less negative following habituation to RP (**Figure 8A**, $P<0.05$). The net protein
397 balance was increased in the postprandial period ($P<0.0001$), albeit no differences between
398 habituation periods (Figure 8D).

399 There were no differences between interventions in the change from fasting to the postprandial
400 state for all protein kinetic parameters (the endogenous rate of appearance, phenylalanine to
401 tyrosine conversion, endogenous rate of disappearance, and the resultant net balance, data not
402 shown).

403

404 *Amino acid concentrations and net splanchnic tissues amino acid balance*

405 The arterial amino acid concentrations and splanchnic net balances are presented in **Table 4**.

406 The amino acid concentrations in the fasted state were similar for RP and HP except for

407 glycine. After meal and protein intake the concentrations of glycine (Gly), asparagine (Asn),

408 alanine (Ala), tyrosine (Tyr) and isoleucine (Ile) were significantly higher following habituation

409 to RP at several timepoints compared with subjects following HP diet ($P < 0.05$). In addition,

410 tendencies for higher concentrations were also observed for lysine (Lys), tryptophan (Trp),

411 methionine (Met) and threonine (Thr) ($P < 0.07$) in RP versus HP habituated participants. The

412 measured net balances across the splanchnic bed revealed no intervention (RP versus HP)

413 effect. It should be acknowledged that the splanchnic balance techniques bear relatively higher

414 variability as for example the single arterial amino acid concentration measurement. In the fed

415 state, the net balances for all amino acids but glutamine (Gln) became more negative indicating

416 a net release of amino acids from the splanchnic tissues to the circulation. However, no

417 changes were observed for glutamine (Gln) in the fasted compared to fed state.

418

419

420

421

422

423

424

425 **Discussion**

426 The main findings of the present study are: 1. Twenty days of habituation to HP intake
427 enhances the whole body protein turnover rate; 2. HP habituation leads to a more negative
428 overnight fasted net protein balance; 3. The higher overnight fasted net protein loss with HP
429 was not compensated for in the 4 h postprandial period after a protein-dense meal containing
430 the same amount of protein for HP and RP; 4. Postprandial plasma protein FSR and
431 consequently incorporation of meal derived phenylalanine into plasma proteins were
432 significantly higher following habituation to RP.

433

434 Similar results as our finding, that whole body protein turnover is increased in response to
435 habituation to HP, were reported by Pannemans and colleagues in young (17) and older (19)
436 participants, who were habituated to either 12 or 21 E% protein. Whereas after habituation to
437 0.70 g protein/kg BW/day or 1.53 g protein/kg BW/day for 14 days, Gorissen and colleagues
438 (20) show no difference in whole body protein synthesis nor breakdown, in response to a
439 protein intake, thus no change in protein turnover. The major difference being that Gorissen
440 et al. (20) studied the postprandial response to only ~0.42 g whey protein/kg LBM without
441 concomitant energy intake. Such amount of protein may have been insufficient to stimulate
442 protein turnover kinetics optimally in older men. Moore and colleagues showed that for plateau
443 stimulation of muscle protein synthesis rates requires an intake of 0.61 g protein/kg LBM in
444 older individuals (33), which we provided in the mixed meal.

445 An enhanced protein turnover rate may be advantageous as an improved capacity to repair and
446 remodel tissues and maintain protein function or adaptation to a physiological challenge.
447 However, in the overnight fasted state HP habituation led to a more negative whole-body net
448 protein balance, which has also been shown by Hursel et al. comparing long term habituation

449 to rather high (2.4 g protein/kg/d) with rather low protein (0.4 g protein/kg/d)(18). In the
450 present study, the more negative whole-body net balance was not counterbalanced by a more
451 positive net balance in the postprandial state. In a conservative scenario, assuming a fasting
452 state condition of 8 hours a day, the negative net balance means a net protein loss of ~3 g of
453 whole body proteins/day. This is equivalent to ~1 kg body protein/year assuming that the body
454 proteins consists of 4.5% phenylalanine (34). It should be noted that this estimated loss is
455 solely based on the fasting condition and does not represent a real-life situation. However, the
456 fact that the adaptations to higher protein intakes may affect the utilization of amino acids in a
457 negative direction should be considered. Combined, these results imply that care should be
458 taken before providing large amounts of proteins. However, it should also be kept in mind that
459 too low protein intake should at all times be prevented, as it has shown to lead to lean body
460 mass erosion (5,35).

461

462 The rate limiting step in phenylalanine degradation hydroxylation of phenylalanine to tyrosine
463 (32) was elevated following habituation to HP both in the fasted and the 4-hour postprandial
464 state. Furthermore, the higher amount of urea excreted in the urine and the higher urea rate of
465 appearance in the blood under fasting conditions at day 21 emphasize that high protein intake
466 results in higher nitrogen excretion. Of importance, even on day 22, where the participants
467 ingested the exact same protein amount at all meals, the urine urea content remained 23%
468 elevated in the HP condition (Figure 3C), emphasizing a reduced ability to retain amino acids
469 and nitrogen when habituated to high protein intake. Such effect can be described as an
470 impaired utilization of amino acids when habituated to HP diet.

471 All circulating amino acids concentrations with the exception of glycine were similar for RP
472 and HP in the fasted state whereas postprandially, the glycine, asparagine, alanine, tyrosine and

473 isoleucine concentrations were higher in RP. A higher postprandial amino acid concentration
474 can originate from a difference in digestion/absorption rate, splanchnic first by-pass extraction
475 and/or reduced peripheral clearance and increased production rates. The exogenous whey and
476 caseinate phenylalanine appearance rates (Figure 7) reveal similar protein digestion and
477 absorption rates after RP and HP habituation and the postprandial whole body protein
478 synthesis, degradation and net balance were also similar indicating similar peripheral clearance
479 and release rates. Therefore, the higher arterial concentrations of some amino acids after RP
480 habituation most likely originate from a lower net splanchnic first by-pass extraction of these
481 amino acids. To quantify arterial-hepatic venous differences, blood samples were taken from
482 the radial artery and the hepatic vein, reflecting the net balance across the splanchnic tissues
483 (36) despite the mixing of the arterial blood to the liver with portal venous content. In the
484 postprandial period, there was a net release for all amino acids, except glutamine, into the
485 circulation from the splanchnic tissue reflecting the expected net uptake. For glutamine the
486 demand in the splanchnic tissue extract most content in the meal, which is in agreement with
487 previous findings from Stoll et al. estimating that in piglets the gut tissue utilizes more than
488 95% of dietary glutamine (37). Together with alanine, glutamine is a key nitrogen carrier and
489 intermediate in amino acid transamination processes. Moreover, glutamine is the preferred
490 energy source of rapidly dividing cells such as those present in the intestine (36).

491 For all of the measured amino acids the net balance across the splanchnic tissues was similar
492 following the two habituation periods. Despite the even net balance across the splanchnic
493 tissues phenylalanine oxidation occurring primarily in the liver was significantly increased
494 following habituation to HP intake (Figure 4). Thus, another process must utilize more amino
495 acids in the RP condition, which appeared to be for plasma protein synthesis, which primarily
496 occurs in the hepatocytes (38). In agreement, the whey-derived amino acid tracer (D₅-
497 phenylalanine) was more abundant in the plasma proteins four hours after meal intake after RP

498 habituation. Overall, these results indicate that during habituation to high protein diets, the
499 amino acid metabolism is higher, leaving less amino acids for the translational apparatus even
500 in the postprandial period.

501

502 A limitation of the present study is that we cannot extrapolate our findings to the entire 24-
503 hour protein net balance. Irrespectively, it was clear that the four-hour net protein balance
504 response to the same meal did not compensate for the net protein loss identified during the end
505 of an overnight fasting period, despite the fact that it was a protein rich meal. The postprandial
506 net protein balance might have differed between the two habituations had the breakfast meal
507 not been identical but contained a protein level comparable to the habituated period as seen by
508 Kim et al. (40). In this study, following habituation to a high protein intake we measured the
509 response to a protein intake similar to the habituated level, and following habituation to the
510 recommended protein level we measured the response to an intake of protein which was twice
511 as high as the habituated level. This means that following the RP period, the meal stimuli is out
512 of the ordinary. Hence their response could potentially be exacerbated.

513

514 In conclusion, a protein intake higher than recommended level resulted in a significantly more
515 negative fasting whole-body net protein balance. The ingestion of a mixed protein dense meal
516 did not compensate for the lowered fasting net protein balance in the first four postprandial
517 hours. Further, the findings of a lower plasma protein FSR and less abundance of dietary whey
518 derived phenylalanine in plasma proteins point towards a higher first pass splanchnic
519 extraction of amino acids from the meal when habituated to high protein content in the diet.
520 This is most likely caused by a higher amino acid catabolism in the liver yielding the observed
521 higher urea production and excretion.

522 Taking the limitations into consideration, these results indicate potential drawbacks of
523 ingesting a 'higher than currently recommended protein level to improve the body's protein
524 utilization. Rather it increases the body's catabolism of amino acids, measured as increased
525 amino acids oxidation and nitrogen excretion.

526

527

528

529

530

531

532 **Acknowledgements**

533 The authors' responsibilities were as follow: GH, JB, GvH and LH designed the study, GH,
534 JaB, AA, JB, LS, GvH and LH Conduced the research, GH, JB, LS, JA, PS, GvH and LH
535 analyzed the data, GH drafted the manuscript and had primary responsibility for final content,
536 all authors read and approved the final manuscript. None of the authors report a conflict of
537 interest related to the study.

538

References:

1. Trumbo P, Schlicker S, Yates AA, Poos M. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids. *Journal of the American Dietetic Association*. 2002. p. 1621–30.
2. EFSA Nda (Panel on Dietetic Products, Nutrition AA. Scientific Opinion on Dietary Reference Values for protein. *EFSA J*. 2012;10:2557–623.
3. World Health Organization, Food and Agriculture Organization of the United Nations UNU. Protein and Amino Acid Requirements in Human Nutrition. World Health Organization technical report series. 2007.
4. Nordic Council of Ministers NC of M. Nordic Nutrition Recommendations 2012. Nordic Nutrition Recommendations 2012. 2012 Aug.
5. Houston DK, Nicklas BJ, Ding J, Harris TB, Tylavsky FA, Newman AB, Lee JS, Sahyoun NR, Visser M, Kritchevsky SB, et al. Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *Am J Clin Nutr*. 2008;87:150–5.
6. Meng X, Zhu K, Devine A, Kerr DA, Binns CW, Prince RL. A 5-year cohort study of the effects of high protein intake on lean mass and BMC in elderly postmenopausal women. *J Bone Miner Res*. 2009;24:1827–34.
7. Loenneke JP, Loprinzi PD, Murphy CH, Phillips SM. Per meal dose and frequency of protein consumption is associated with lean mass and muscle performance. *Clin Nutr*. Elsevier Ltd; 2016;35:1506–11.
8. Hughes VA, Frontera WR, Roubenoff R, Evans WJ, Fiatarone Singh MA. Longitudinal

- changes in body composition in older men and women: Role of body weight change and physical activity. *Am J Clin Nutr.* 2002;76:473–81.
9. Guillet C, Prod'homme M, Balage M, Gachon P, Giraudet C, Morin L, Grizard J, Boirie Y. Impaired anabolic response of muscle protein synthesis is associated with S6K1 dysregulation in elderly humans. *FASEB J.* 2004;18:1586–7.
 10. Volpi E, Mittendorfer B, Rasmussen BB, Wolfe RR. The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. *J Clin Endocrinol Metab.* 2000;85:4481–90.
 11. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A, Wolfe RR. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. *Am J Physiol Endocrinol Metab.* 2006;291:E381-7.
 12. Volpi E, Mittendorfer B, Wolf SE, Wolfe RR. Oral amino acids stimulate muscle protein anabolism in the elderly despite higher first-pass splanchnic extraction. *Am J Physiol.* 1999;277:E513–20.
 13. Boirie Y, Gachon P, Beaufrere B. Splanchnic and whole-body elderly men¹³ leucine kinetics in young and elderly men. *Am J Clin Nutr.* 1997;65:489–95.
 14. Steffee P, Goldsmith RS, Young R, Scrimshaw NS, Steffee WP, Goldsmith RS, Pencharz PB, Scrimshaw NS, Young VR. Dietary protein intake and dynamic aspects of whole body nitrogen metabolism in adult humans. *Metabolism.* 1976;25:281–97.
 15. Forslund AH, Hambræus L, Olsson RM, El-Khoury AE, Yu YM, Young VR. The 24-h whole body leucine and urea kinetics at normal and high protein intakes with exercise in

- healthy adults. *Am J Physiol.* 1998;275:E310-20.
16. Young VR, El-Khoury AE, Raguso CA, Forslund AH, Hambraeus L. Rates of urea production and hydrolysis and leucine oxidation change linearly over widely varying protein intakes in healthy adults. *J Nutr.* 2000;130:761–6.
 17. Pannemans DL, Halliday D, Westerterp KR, Kester AD. Effect of variable protein intake on whole-body protein turnover in young men and women. *Am J Clin Nutr.* 1995;61:69–74.
 18. Hursel R, Martens EAP, Gonnissen HKJ, Hamer HM, Senden JMG, Van Loon LJC, Westerterp-Plantenga MS. Prolonged adaptation to a low or high protein diet does not modulate basal muscle protein synthesis rates - A substudy. *PLoS One.* 2015;10:1–15.
 19. Pannemans DL, Halliday D, Westerterp KR. Whole-body protein turnover in elderly men and women: responses to two protein intakes. *Am J Clin Nutr.* 1995;61:33–8.
 20. Gorissen SH, Horstman AM, Franssen R, Kouw IW, Wall BT, Burd NA, de Groot LC, van Loon LJ. Habituation to low or high protein intake does not modulate basal or postprandial muscle protein synthesis rates: a randomized trial. *Am J Clin Nutr.* 2016;
 21. Reitelseder S, Tranberg B, Agergaard J, Dideriksen K, Højfeldt G, Merry ME, Storm AC, Poulsen KR, Hansen ET, van Hall G, et al. Phenylalanine stable isotope tracer labeling of cow milk and meat and human experimental applications to study dietary protein-derived amino acid availability. *Clin Nutr.* 2020;44:1–50.
 22. Bornø A, Van Hall G. Quantitative amino acid profiling and stable isotopically labeled amino acid tracer enrichment used for in vivo human systemic and tissue kinetics measurements. *J Chromatogr B Anal Technol Biomed Life Sci.* Elsevier B.V.;

- 2014;951–952:69–77.
23. Bornø A, Hulston CJ, Van Hall G. Determination of human muscle protein fractional synthesis rate: An evaluation of different mass spectrometry techniques and considerations for tracer choice. *J Mass Spectrom.* 2014;49:674–80.
 24. Asmar A, Asmar M, Simonsen L, Madsbad S, Holst JJ, Hartmann B, Sorensen CM, Bülow J. Glucagon-like peptide-1 elicits vasodilation in adipose tissue and skeletal muscle in healthy men. *Physiol Rep.* 2017;5:1–12.
 25. Henriksen JH, Winkler K. Hepatic blood flow determination. A comparison of ^{99m}Tc-diethyl-IDA and indocyanine green as hepatic blood flow indicators in man. *J Hepatol.* 1987;4:66–70.
 26. Steele R. Influences of glucose loading and of injected insulin on hepatic glucose output. *Ann N Y Acad Sci.* 1959;82:420–30.
 27. Proietto J, Rohner-Jeanrenaud F, Ionescu E, Terrettaz J, Sauter JF, Jeanrenaud B. Non-steady-state measurement of glucose turnover in rats by using a one-compartment model. *Am J Physiol.* 1987;252:E77-84.
 28. Boirie Y, Gachon P, Corny S, Fauquant J, Maubois JL, Beaufrère B. Acute postprandial changes in leucine metabolism as assessed with an intrinsically labeled milk protein. *Am J Physiol.* 1996;271:E1083-91.
 29. Groen BBL, Res PT, Pennings B, Hertle E, Senden JMG, Saris WHM, van Loon LJC. Intra-gastric protein administration stimulates overnight muscle protein synthesis in elderly men. *AJP Endocrinol Metab.* 2012;302:E52–60.
 30. Dangin M, Guillet C, Garcia-Rodenas C, Gachon P, Bouteloup-Demange C, Reiffers-

- Magnani K, Fauquant J, Ballèvre O, Beaufrère B. The rate of protein digestion affects protein gain differently during aging in humans. *J Physiol.* 2003;549:635–44.
31. Smith GI, Patterson BW, Klein SJ, Mittendorfer B. Effect of hyperinsulinaemia-hyperaminoacidaemia on leg muscle protein synthesis and breakdown: reassessment of the two-pool arterio-venous balance model. *J Physiol.* 2015;593:4245–57.
 32. Thompson GN, Pacy PJ, Merritt H, Ford GC, Read M a, Cheng KN, Halliday D. Rapid measurement of whole body and forearm protein turnover using a [2H5]phenylalanine model. *Am J Physiol.* 1989;256:E631–9.
 33. Moore DR, Churchward-Venne TA, Witard O, Breen L, Burd NA, Tipton KD, Phillips SM. Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *J Gerontol A Biol Sci Med Sci.* 2015;70:57–62.
 34. Beach EF, Munks B, Robinson A. The Amino Acid Composition of Animal Tissue Protein. *J Biol Chem.* 1943;148:431–9.
 35. Mangano KM, Sahni S, Kiel DP, Tucker KL, Dufour AB, Hannan MT. Dietary protein is associated with musculoskeletal health independently of dietary pattern: the Framingham Third Generation Study. *Am J Clin Nutr.* 2017;105:714–22.
 36. Neis EPJG, Sabrkhany S, Hundscheid I, Schellekens D, Lenaerts K, Olde Damink SW, Blaak EE, Dejong CHC, Rensen SS. Human splanchnic amino-acid metabolism. *Amino Acids.* Springer Vienna; 2017;49:161–72.
 37. Stoll B, Henry J, Reeds PJ, Yu H, Jahoor F, Burrin DG. Catabolism dominates the first-pass intestinal metabolism of dietary essential amino acids in milk protein-fed piglets. *J*

- Nutr. 1998;128:606–14.
38. Gersovitz M, Munro HN, Udall J, Young VR. Albumin synthesis in young and elderly subjects using a new stable isotope methodology: response to level of protein intake. *Metabolism*. 1980;29:1075–86.
 39. Jaleel A, Nehra V, Persson X-MT, Boirie Y, Bigelow M, Nair KS. In vivo measurement of synthesis rate of multiple plasma proteins in humans. *Am J Physiol Endocrinol Metab*. 2006;291:E190-7.
 40. Kim I, Shin Y, Schutzler SE, Azhar G, Wolfe RR, Ferrando AA. Quality of meal protein determines anabolic response in older adults. *Clin Nutr*. Elsevier Ltd; 2018;37:2076–83.

Table 1: Subject characteristics at inclusion

Participant characteristics N=12	
Age [years]	66.6 ± 1.6
Height [m]	1.79 ± 0.04
Body weight [kg]	84.3 ± 10.9
BMI [kg/m ²]	26.3 ± 3.3
LBM [kg]	56.4 ± 4.4
Fat %	29.9 ± 5.7
Systolic blood pressure [mmHg]	137 ± 25
Diastolic blood pressure [mmHg]	83 ± 11
Normal dietary protein intake [g/kg LBM/d]	1.5 ± 0.3
Daily energy intake [kcal/LBM]	36.5 ± 7.7
E% protein	18 ± 2.6
Hemoglobin A1c (IFCC) [mmol/mol]	35.2 ± 2.6
Thyrotropin (TSH) [$\times 10^{-3}$ IU/l]	1.5 ± 0.8
Total Cholesterol [mmol/l]	5.5 ± 0.7
HDL cholesterol [mmol/l]	1.7 ± 0.4
LDL Cholesterol [mmol/l]	3.2 ± 0.6
Triglyceride [mmol/l]	1.3 ± 0.8

Values are means ± SD; BMI, body mass index; LBM, lean body mass; E%, energy percent.

Table 2: Macronutrient composition of the breakfast meal.

Trial day breakfast	
Calories [kcal]	441
Protein [g/kg LBM]	0.61
- D ₅ -phe labeled whey [g/kg LBM]	0.31
- ¹⁵ N-phe labeled casein [g/kg LBM]	0.23
Carbohydrates [g/kg LBM]	1.06
Fat [g/kg LBM]	0.21

LBM, lean body mass

Table 3: Total daily calorie and protein intake for the recommended and high protein intake periods including the supplement.

20-day habituation period N=12		
Recommended protein period	Basic diet	Basic diet + supplements
Calorie intake week 2-3 [kcal/day]	2117 ± 84	1624 ± 84
Protein intake day 1-7 [g/kg LBM/day]	1.42 ± 0.09	1.42 ± 0.09
Protein intake day 1-7 [g/kg BW/day]	0.96 ± 0.07	0.96 ± 0.07
Protein intake day 8-20 [g/kg LBM/day]	1.11 ± 0.03	1.11 ± 0.03
Protein intake day 8-20 [g/kg BW/day]	0.75 ± 0.03	0.75 ± 0.03
High protein period	Basic diet	Basic diet +supplements
Calorie intake week 2-3 [kcal/day]	2207 ± 58	1666 ± 58
Protein intake day 1-7 [g/kg LBM/day]	2.78 ± 0.09	1.35 ± 0.07 *
Protein intake day 1-7 [g/kg BW/day]	1.87 ± 0.08	0.91 ± 0.06 *
Protein intake day 8-20 [g/kg LBM/day]	2.53 ± 0.05	1.10 ± 0.03 *
Protein intake day 8-20 [g/kg BW/day]	1.71 ± 0.05	0.74 ± 0.02 *

Data excluding the supplements are noted in the parentheses. LBM, lean body mass; BW, body weight. Values are mean ± SEM. *denotes p<0.001 between interventions.

Table 4: Arterial amino acid concentrations and splanchnic tissues net balances

		Arterial conc.[$\mu\text{mol/L}$] \pm SEM; NB, net balance [$\mu\text{mol/min}$] \pm SEM							
		-10min	30min	60min	90min	120min	150min	180min	240min
Asp	Arterial RP	5 \pm 1	10 \pm 1	12 \pm 3	6 \pm 1	9 \pm 4	6 \pm 1	5 \pm 1	8 \pm 3
	\$ HP	5 \pm 0	10 \pm 1	7 \pm 1	5 \pm 0	5 \pm 1	5 \pm 0	5 \pm 0	5 \pm 1
	NB RP	-4 \pm 1	-23 \pm 5	-5 \pm 3	-5 \pm 1	1 \pm 4	-6 \pm 3	-3 \pm 1	-6 \pm 2
	\$ HP	-3 \pm 0	-14 \pm 3	-5 \pm 2	-3 \pm 1	-2 \pm 1	-2 \pm 1	-2 \pm 0	-2 \pm 0
Glu	Arterial RP	111 \pm 27	127 \pm 14	126 \pm 19	125 \pm 25	110 \pm 20	125 \pm 31	106 \pm 23	107 \pm 23
	\$ HP	93 \pm 10	114 \pm 7	104 \pm 8	96 \pm 6	99 \pm 15	91 \pm 9	89 \pm 8	92 \pm 10
	NB RP	-101 \pm 17	-174 \pm 30	-115 \pm 15	-125 \pm 20	-100 \pm 21	-119 \pm 35	-104 \pm 15	-129 \pm 22
	\$ HP	-81 \pm 11	-131 \pm 22	-88 \pm 24	-103 \pm 19	-67 \pm 21	-81 \pm 15	-85 \pm 11	-83 \pm 14
Ser	Arterial RP	113 \pm 15	167 \pm 15	183 \pm 23	147 \pm 14	144 \pm 27	138 \pm 19	120 \pm 15	130 \pm 20
	\$ HP	102 \pm 5	147 \pm 7	131 \pm 8	114 \pm 7	104 \pm 4	106 \pm 6	104 \pm 7	105 \pm 6
	NB RP	17 \pm 3	-62 \pm 17	1 \pm 15	11 \pm 11	27 \pm 19	-10 \pm 18	-6 \pm 6	-17 \pm 16
	\$ HP	17 \pm 2	-32 \pm 22	-2 \pm 17	4 \pm 3	11 \pm 5	8 \pm 16	6 \pm 6	9 \pm 9
Gly	Arterial RP	*223 \pm 27	*244 \pm 27	*263 \pm 32	*234 \pm 22	*219 \pm 25	*234 \pm 27	209 \pm 21	*229 \pm 25
	\$ HP	184 \pm 11	187 \pm 12	176 \pm 13	169 \pm 12	166 \pm 9	179 \pm 13	180 \pm 16	191 \pm 15
	NB RP	26 \pm 7	-14 \pm 16	20 \pm 8	17 \pm 14	18 \pm 7	-8 \pm 27	-11 \pm 9	-22 \pm 19
	\$ HP	15 \pm 5	-19 \pm 22	10 \pm 19	-2 \pm 16	-2 \pm 9	-7 \pm 24	-4 \pm 12	-1 \pm 18
Asn	Arterial RP	45 \pm 7	86 \pm 11	*91 \pm 11	*79 \pm 7	64 \pm 7	*73 \pm 12	64 \pm 9	60 \pm 8
	\$ HP	44 \pm 3	77 \pm 5	71 \pm 5	62 \pm 6	56 \pm 4	57 \pm 4	56 \pm 5	55 \pm 5
	NB RP	9 \pm 2	-42 \pm 9	-16 \pm 4	0 \pm 6	-3 \pm 4	-8 \pm 5	-10 \pm 3	-8 \pm 4
	\$ HP	7 \pm 2	-27 \pm 13	-9 \pm 9	-7 \pm 7	-4 \pm 3	-4 \pm 8	-3 \pm 4	-1 \pm 5
Gln	Arterial RP	707 \pm 78	857 \pm 100	893 \pm 105	840 \pm 76	721 \pm 68	798 \pm 83	722 \pm 74	741 \pm 76
	\$ HP	648 \pm 29	747 \pm 41	756 \pm 51	718 \pm 50	646 \pm 45	692 \pm 47	687 \pm 49	735 \pm 52
	NB RP	122 \pm 27	39 \pm 53	65 \pm 27	152 \pm 51	78 \pm 32	75 \pm 62	41 \pm 28	13 \pm 51
	\$ HP	97 \pm 14	-8 \pm 77	58 \pm 77	37 \pm 56	47 \pm 38	74 \pm 65	71 \pm 25	61 \pm 58
His	Arterial RP	77 \pm 9	101 \pm 11	110 \pm 11	100 \pm 9	88 \pm 8	97 \pm 11	89 \pm 11	91 \pm 10
	\$ HP	73 \pm 3	93 \pm 5	94 \pm 5	85 \pm 5	79 \pm 3	83 \pm 6	84 \pm 5	88 \pm 6

	NB	RP	8±2	-18±7	-10±3	4±6	0±3	-9±8	-9±4	-14±6
	\$	HP	3±2	-21±11	-7±10	-7±6	-3±4	-5±8	-6±3	-4±7
Thr	Arterial	RP	118±17	217±23	234±25	216±18	182±17	199±25	175±21	166±18
	\$	HP	111±6	197±10	183±9	165±11	156±13	145±8	137±9	134±8
	NB	RP	19±4	-110±25	-45±5	-13±14	-12±9	-24±19	-19±6	-16±11
	\$	HP	16±3	-74±34	-25±24	-19±20	9±16	0±18	2±7	6±10
Ala	Arterial	RP	266±29	*407±50	*493±54	*487±43	402±40	*431±39	396±38	401±42
	\$	HP	225±17	323±29	361±28	362±34	336±24	349±29	356±31	363±30
	NB	RP	152±21	19±37	-3±19	93±41	83±20	77±27	70±19	94±29
	\$	HP	115±10	21±45	31±39	48±36	80±18	93±46	98±19	109±30
Pro	Arterial	RP	167±18	290±25	322±31	332±27	293±25	335±28	306±25	300±24
	\$	HP	155±11	277±18	281±20	278±22	263±15	271±20	272±26	269±19
	NB	RP	9±5	-142±31	-85±7	-68±26	-68±29	-98±45	-87±17	-82±26
	\$	HP	0±6	-127±47	-72±38	-89±38	-56±18	-55±35	-47±14	-32±18
Arg	Arterial	RP	84±11	138±14	139±16	128±12	105±10	119±15	102±11	95±10
	\$	HP	70±5	116±7	105±7	96±6	86±4	88±7	84±7	81±5
	NB	RP	13±3	-44±13	-14±2	-1±8	-10±9	-15±14	-13±5	-13±8
	\$	HP	8±2	-30±21	-10±13	-13±15	-1±5	-3±13	-1±4	-1±7
Tyr	Arterial	RP	62±7	125±12	*132±13	*125±10	105±10	*116±12	*102±10	95±10
	\$	HP	64±3	114±7	110±7	98±5	88±4	87±4	84±6	81±4
	NB	RP	8±3	-45±19	-10±3	2±8	-4±7	-7±12	-3±4	-5±8
	\$	HP	8±2	-30±24	-11±15	-9±13	-1±4	-3±12	2±3	1±6
Val	Arterial	RP	235±28	409±38	429±40	417±30	349±29	388±42	341±37	323±33
	\$	HP	260±11	427±22	419±21	386±20	344±18	345±22	326±21	315±18
	NB	RP	7±6	-154±38	-81±5	-35±24	-49±23	-64±34	-58±11	-62±21
	\$	HP	-4±7	-144±63	-71±49	-85±38	-49±20	-42±37	-34±12	-28±23
Met	Arterial	RP	21±2	51±5	51±5	45±2	35±3	39±5	34±3	28±3
	\$	HP	20±1	47±3	40±3	34±3	30±1	29±2	27±2	25±2
	NB	RP	5±1	-24±5	-6±2	2±3	-1±2	-3±3	-2±1	-1±2
	\$	HP	4±0	-16±10	-3±5	-3±5	0±2	0±4	1±1	2±2
Ile	Arterial	RP	52±8	177±22	175±21	*152±13	112±12	*129±21	111±17	91±11
	\$	HP	55±4	176±14	154±14	121±8	108±10	91±5	82±5	74±4

	NB	RP	0±1	-121±28	-52±5	-24±14	-34±13	-30±15	-28±11	-26±7
	\$	HP	-3±3	-102±44	-52±26	-44±18	-10±14	-22±13	-19±5	-14±7
Leu	Arterial	RP	124±14	376±36	394±41	336±28	258±23	283±38	242±33	201±24
	\$	HP	135±9	400±33	358±30	287±17	242±16	220±17	203±20	180±15
	NB	RP	5±4	-247±58	-98±10	-38±27	-50±19	-64±22	-51±7	-37±10
	\$	HP	-7±4	-224±82	-106±50	-82±34	-28±20	-37±24	-26±8	-10±10
Tryp	Arterial	RP	54±7	97±10	106±12	98±9	79±7	86±9	72±8	65±7
	\$	HP	51±2	87±5	85±4	76±3	67±3	66±4	61±4	58±3
	NB	RP	3±1	-31±10	-10±1	1±6	-4±5	-7±10	-5±3	-7±5
	\$	HP	1±1	-28±14	-6±9	-9±7	-3±3	-2±8	-1±2	-1±5
Phe	Arterial	RP	52±6	91±7	92±8	85±5	73±5	83±8	73±7	71±6
	\$	HP	58±2	90±4	83±4	79±4	74±4	75±4	73±5	72±3
	NB	RP	7±2	-25±7	-5±3	1±5	-4±4	-7±7	-6±2	-7±5
	\$	HP	5±1	-21±13	-4±8	-9±8	-3±5	-4±9	-1±3	0±5
Lys	Arterial	RP	177±19	368±33	374±34	325±21	260±20	295±34	257±26	231±22
	\$	HP	187±9	363±19	318±17	273±15	241±9	242±13	230±15	219±12
	NB	RP	18±5	-186±47	-56±4	-10±22	-22±19	-30±23	-32±9	-28±14
	\$	HP	13±5	-146±70	-32±39	-33±33	-10±13	-14±32	-8±10	-5±19
Ess.	Arterial	RP	411±49	962±93	997±96	904±65	719±61	800±98	694±85	614±66
	\$	HP	450±21	1003±61	931±59	794±40	693±33	655±40	611±42	569±33
BCAA	NB	RP	13±10	-521±123	-232±14	-97±62	-132±52	-158±69	-137±26	-125±37
	\$	HP	-16±14	-470±188	-229±121	-212±90	-86±50	-101±74	-78±23	-51±32
Ess.excl.	Arterial	RP	591±67	1078±99	1118±107	1009±69	835±66	934±105	820±83	764±72
	\$	HP	577±24	1010±46	922±45	818±43	744±23	739±38	709±43	689±33
BCAA	NB	RP	81±17	-462±117	-151±16	-18±66	-53±52	-96±85	-87±30	-84±53
	\$	HP	56±14	-351±180	-95±112	-96±100	-7±43	-28±95	-10±30	1±56
Total	Arterial	RP	2703±313	4337±423	4618±470	4273±326	3605±308	3971±432	3523±365	3430±336
	\$	HP	2541±93	3992±193	3836±203	3501±197	3186±127	3220±181	3137±206	3139±177
	NB	RP	328±76	-1413±400	-532±61	-40±272	-154±182	-360±375	-340±133	-374±245
	\$	HP	206±57	-1188±621	-410±443	-432±375	-89±174	-105±377	-56±127	9±232

NB=Splanchnic tissues net balance, a negative value means a higher amino acid concentration in the hepatic vein, i.e. a net amino acid release from the splanchnic tissue into the main circulation. Ess. BCAA, essential branched-chain amino acids; ess.excl. BCAA, essential amino acids excluding branched-chain amino acids. \$ denotes an overall time effect, with a significant change from fasting, * denotes an interaction effect with a Sidak post hoc test revealing significant difference between RP and HP ($P < 0.05$). N=11, data is mean \pm SEM. Ess. =essential .

Figure legends:

Figure 1: Flow chart of the enrollment process.

Figure 2: Overview of the 20-day trial period, with emphasis on the day 21 trial day.

Figure 3: Urea amount in the urine A) 24 hour urine day 20-21, post '20 day habituation', B) 8 hour urine during the trial, including lunch C) 24 hour urine day 21-22, including the trial period. RP, recommended protein period; HP, high protein period. * denotes significant differences between interventions assessed by a paired t-test. Fig. A+B N=10, fig. C N=12, values are mean \pm SEM.

Figure 4: Nitrogen and amino acid loss as well as phenylalanine conversion in the fasted and postprandial state. Figure A and B show the urea rate of appearance in the hepatic vein in the post habituation, fasted state (A), pre and four-hour postprandial (B). (N=10, values are mean \pm SEM). Figure C and D shows the phenylalanine conversion to tyrosine. RP, recommended protein period; HP, high protein period; R_a , rate of appearance. (N=8, values are mean \pm SEM). \$ denotes significant change from fasting, * denotes significant difference between habituated protein levels. In the fasted state comparison between groups are done by a paired t-test while changes over time and between groups are done by a two-way ANOVA with repeated measures.

Figure 5: 0-4 hours fractional synthesis rate (FSR) for plasma proteins (A) and incorporation of amino acids from ingested caseinate (^{15}N -phenylalanine) (B) and whey (D_5 -phenylalanine) (C) into plasma proteins. The plasma protein FSR in the 4 hours postprandial period (A) is significantly higher following habituation to RP compared HP intake ($P < 0.05$). Phenylalanine from the breakfast meal into plasma protein is significantly increased with time (B and C). RP, recommended protein period; HP, high protein period; FSR, fractional synthesis rate;

APE, atom percent excess; MPE, mole percent excess. \$ denotes significant change from 1 hour ($P < 0.05$). The enrichment deriving from the D_5 -phenylalanine from the whey protein is significantly higher in the plasma proteins 4 hours post meal intake when participants are habituated to RP intake. * denotes significant difference between habituations ($P < 0.001$). Plasma FSR comparison between groups are done by a paired t-test while changes in enrichment over time and between groups are done by a two-way ANOVA with repeated measures. Values mean \pm SEM, $N=11$.

Figure 6: The insulin concentrations in the post habituation, fasted state (A), and immediately pre and four hours postprandial (B). RP, recommended protein period; HP, high protein period. \$ denotes significant change from fasting. In the fasted state comparison between groups are done by a paired t-test while changes over time and between groups are done by a two-way ANOVA with repeated measures. ($N=12$, values are mean \pm SEM).

Figure 7: Exogenous rate of appearance of the ingested protein in the hepatic vein (A) representing the rate at which the intrinsically labeled phenylalanine enters the circulation And in the radial artery (B), representing the rate of appearance into the radial artery. For the appearance rate in both the hepatic vein and the radial artery, there is a significant time effect ($p < 0.0001$) and a difference between R_a of Whey and R_a of casein on time, intervention and interaction ($p < 0.0001$). RP, recommended protein period; HP, high protein period; Exo, exogenous: R_a , rate of appearance. Changes over time and between groups are done by a two-way ANOVA with repeated measures. ($N=11$, values are mean \pm SEM)

Figure 8: Whole body protein turnover in the fasted (A) and immediately pre and postprandial state (B, C, D). The endogenous release of phenylalanine represents the whole body protein breakdown in fasted (A) and fed state (B). The endogenous rate of disappearance

represents the protein synthesis in the fasted (A) and fed state (C). The net balance is shown in the fasted (A) and fed state (D). RP, recommended protein period; HP, high protein period; endo, endogenous; R_a , rate of appearance; R_d , rate of disappearance. \$ denotes significant change from fasting, * denotes significant difference between habituated protein levels. In the fasted state comparison between groups are done by a paired t-test while changes over time and between groups are done by a two-way ANOVA with repeated measures. (N=8, values are mean \pm SEM).