

Natural selection on crosstalk between gene regulatory networks facilitates bacterial adaptation to novel environments

Taylor, Tiffany B.; Shepherd, Matthew; Jackson, Robert; Silby, Mark W.

DOI:
[10.1016/j.mib.2022.02.002](https://doi.org/10.1016/j.mib.2022.02.002)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Taylor, TB, Shepherd, M, Jackson, R & Silby, MW 2022, 'Natural selection on crosstalk between gene regulatory networks facilitates bacterial adaptation to novel environments', *Current Opinion in Microbiology*, vol. 67, 102140. <https://doi.org/10.1016/j.mib.2022.02.002>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Natural selection on crosstalk between gene regulatory networks facilitates bacterial adaptation to novel environments

Tiffany B Taylor¹, Matthew J Shepherd¹,
Robert W Jackson² and Mark W Silby³



At the level of the gene, mutation is the raw material for natural selection. However, at the level of the gene regulatory network (GRN), variation is revealed to selection via promiscuous regulator activity ('crosstalk'), which creates opportunities for genetic innovation that can facilitate adaptation. Many genetic and environmental features can contribute to increasing potential for crosstalk by facilitating non-cognate interactions between regulatory elements. If a novel interaction provides a fitness benefit, rewired GRNs with strengthened affinity for newly forged connections can be selected. Here, we identify factors that facilitate opportunities for crosstalk and rewiring between GRNs, consider whether features of some GRNs make them more 'rewireable' than others and if these features might constrain evolution towards convergent outcomes. We explore patterns from laboratory and natural microbial populations that show changes within GRNs during adaptation. Finally, we discuss the prospects and open questions in the field.

Addresses

¹ Milner Centre for Evolution, Department of Biology & Biochemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

² School of Biosciences and Birmingham Institute of Forest Research (BIFoR), University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

³ Department of Biology, University of Massachusetts Dartmouth, 285 Old Westport Road, North Dartmouth, MA 02747, USA

Corresponding author: Taylor, Tiffany B (t.b.taylor@bath.ac.uk)

Current Opinion in Microbiology 2022, 67:102140

This review comes from a themed issue on **Environmental microbiology**

Edited by **Paul Hoskisson** and **John Bruce**

For complete overview of the section, please refer to the article collection, "**Environmental Microbiology**"

Available online 3rd March 2022

<https://doi.org/10.1016/j.mib.2022.02.002>

1369-5274/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Features that facilitate adaptation of gene regulatory networks

Gene regulatory networks (GRNs) evolve through changes to regulatory connections [1]. 'Wiring' (i.e. the specific connections between regulatory elements) within

a cell's regulatory network is as susceptible to adaptation as genome organisation [2,3]. However, factors that create and determine the evolutionary potential for GRN **rewiring** (for this and other terms used in the review please refer to the Glossary in **Box 1**) are poorly understood. Extensive sequence and structural similarity within protein families exist in prokaryotes between components of different pathways due to past cycles of gene duplication, divergence, and horizontal gene transfer [4]. This process of GRN expansion can also result in many regulators controlling overlapping **regulons** [5–8]. Together this confers 'built-in' promiscuity and modularity of GRNs that facilitates rapid evolution (**Figure 1d(3)**).

Promiscuous binding between regulators and non-cognate binding sites leading to transcriptional activation of non-target genes is termed **crosstalk**. In general, crosstalk between GRNs is low, because misregulation of genes is likely to reduce fitness in an environment where an organism is well adapted, and consequently there are mechanisms in place that aid their insulation [9–11]. However, crosstalk can make new interactions available to selection that (if beneficial) can facilitate the evolution of GRNs [12]. Therefore, there exists a balance between maintaining specificity between TFs and their **cognate binding sites** to ensure network efficiency, and relaxing specificity to enable adaptation of networks. Where this balance lies remains an open question.

The role of TF binding affinity and range in facilitating crosstalk

One required change for evolutionary rewiring of a GRN is for a TF to gain new regulatory function that selection can act upon. As with **neofunctionalisation** in enzymes and other proteins [13], this is commonly thought to occur through the gain of low-level promiscuous side-activity (i.e. crosstalk) [14–16]. If new connections are beneficial, selection can act on mutations in the TF binding domain or on mutations that change the promiscuously regulated promoter sequence, to strengthen the binding potential between the TF and consensus binding sequence — increasing affinity between the pair [17*]. Promoter sequences can evolve through very few mutational changes [18], and shared structural homology between TFs means binding domains can be modified towards new targets readily [19] demonstrating the ease with

Box 1 Glossary:

Autoregulatory loops: where a transcription factor regulates its own expression.

Crosstalk: a change in transcription of non-target genes that results from promiscuous binding. Although there are other mechanisms that can confer crosstalk, they are not relevant to the perspective of this review.

Cognate binding-sites: an established, co-evolved interaction site between transcription factor and DNA-sequence that optimises gene expression in a given environment.

De novo: translating to 'a new', in evolution it describes when a genetic variant or a mutation arises in a population.

DNA-binding domain: a folded protein domain that matches and binds to a region of DNA, allowing protein–DNA interactions.

Global transcriptional regulators: transcription factors that control many essential genes in response to environmental changes.

GRN architecture: the interaction profile of regulatory elements within a GRN.

Neofunctionalisation: when a paralog gene takes on a novel function after duplication.

Orthologous: homologous genes in different species as a product of shared common ancestry. These genes often retain original function.

Paralogous: homologous genes in the same genome as a product of gene duplication and specialisation. These genes often have different functions.

Promiscuous binding: when a transcription factor binds to a non-cognate site

Regulons: a group of genes that are regulated as a single unit.

Rewiring: the emergence of new regulatory interactions between different GRNs.

Two-component regulatory systems: a subset of GRN that enables a cell to sense and respond to changing environmental conditions. Typically, it consists of a sensor histidine-kinase (HK) that will sense an environmental cue, and upon activation phosphorylate an associated response regulator (RR) that will initiate transcriptional changes in response to stimulus.

which beneficial connections can be strengthened under selection [20,21].

TFs engage with DNA sequences via a **DNA-binding domain** [22]. However, these binding sites are not uniform in sequence, instead being variations of consensus sequences [23] with differing binding affinities for the TF involved [24,25]. In particular, A/T-rich consensus sequences show lower binding specificity and therefore promiscuous binding by multiple TFs is more likely [26**] (Figure 1d(4)). This could explain the high incidence of A/T rich promoter regions in horizontally acquired genes that may facilitate expression through crosstalk via non-native TFs [27]. In addition, TFs can also vary in their potential to bind to multiple sequences. Detailed analysis of 182 TF-DNA interactions in *Pseudomonas aeruginosa* showed that the number of binding sites varied substantially, with most TFs binding fewer than

100 targets; however, a considerable proportion (~36%) bound to more, with 8 exceeding 1000 binding sites in the genome [28]. In a similar study from Fan *et al.* looking at TF binding affinity in *Pseudomonas syringae*, out of 100 TFs with identified binding motifs most bound fewer than 200 binding sites with 5 showing more than 500 binding sites [29*]. The consequence of this variation, however, is unknown in terms of whether TF binding affinity and range might afford greater opportunity for GRN adaptation and optimisation in variable environments through the facilitation of crosstalk.


The role of environment in revealing pre-existing crosstalk to selection

The interaction of **GRN architecture** (i.e. abundance and nature of network connections) and transduced environmental signals determines the concentration of active TF for controlling the activity of a particular gene [30–32,33*]. Commonly found in prokaryotes, **two-component regulatory systems** (TCSs) are a subsystem of GRNs that specifically link environmental change to gene expression as they enable organisms to sense and respond to changing environmental conditions.

TCSs comprise a sensor histidine kinase (HK) and their cognate response regulator (RR) (Figure 1a). As with TFs, more complex environments select for systems with more precise control and thus more TCSs [1,34], and they are frequent sites of adaptive mutations during ecological niche shifts [35,36]. These simple GRNs offer a good model system to empirically explore the role of the environment in selecting for novel or rewired GRNs in terms of the mutational targets and architectures that readily facilitate adaptation.

Several elegant experimental studies highlight the role of environmental conditions in determining GRN adaptive routes, by utilising pre-existing indirect links (which may be considered crosstalk). For example, the evolutionary recovery of pili-mediated motility in a *Myxococcus xanthus* Δ *pilR* strain [37]. In wildtype, NmpR (a RR in the multicomponent system NmpRSTU) indirectly modulates *pilA* expression under low oxygen. In normal oxygen, HK NmpU is active and phosphorylates a hybrid RR-HK NmpS, but in low oxygen NmpS adopts its role as a HK to phosphorylate its RR (NmpR) – phosphorylated NmpR then interacts with *pilR* binding sites to express pili associated genes. In a Δ *pilR* mutant under selection for pili-mediated motility, suppressor mutations in HK NmpU were recovered in seven of eight motile mutants and in six of these were the only mutation reported. In addition, activation of NmpR in evolved strains and deletion of *nmpU* in the parental Δ *pilR* strain is sufficient to restore motility, which strongly suggests the *de novo* mutations as the causative agent of motility phenotype [36].

Figure 1



(a)–(d) Environmental and genetic features that create opportunity for crosstalk using a two-component system as an example GRN. (a) Under normal regulation, the sensor histidine kinase (HK) will receive an extra-cellular signal, leading to phosphorylation of the response regulator (RR), which in its activate state can bind to its cognate DNA promoter region to initiate transcription of target genes. In some cases, HK can also act as a phosphatase to de-activate RR. (b) (1) suppressor mutations to phosphatase activity of HK, or (c) (2) loss-of-function mutations to HK repressor genes lead to an elevated concentration of active TF in the cell that increases the chance of non-cognate interactions. In both these cases, sufficient environmental signal is necessary to initiate HK activity (represented by extracellular blue spheres); (d) (3) Sequence/ structural homology between TFs confers ‘built-in’ promiscuity and modularity of GRNs that can facilitate promiscuous binding; (d) (4) Similar

(Figure 1 Legend Continued) promoter sequences, or those with low binding affinities for example, those with high A/T content, show higher propensity to be bound promiscuously by multiple TFs. The strength of transcription is indicated by the thickness of line (with dotted lines representing weak transcription due to non-cognate binding). Created with [BioRender.com](https://www.biorender.com).

Similarly, in a non-flagellate $\Delta fleQ$ strain of *Pseudomonas fluorescens*, promiscuity and rewiring of nitrogen RR NtrC was able to rescue flagellar gene expression after a hyper-activating mutation of its cognate HK NtrB or loss of the NtrB-repressor GlnK [20], which led to increased *ntrBC* expression due to autoregulation. Transcriptome analysis shows nitrogen compound transport genes are downregulated in the $\Delta fleQ$ mutant compared to wildtype, suggesting pre-existing indirect links between the flagellar and nitrogen regulatory networks. Without any mutations to the RR NtrC, $\Delta fleQ$ mutants were able to recover some activity of the flagellar pathway via mutations that increased intracellular levels of active NtrC. Once revealed to selection, second step mutations increased the affinity of NtrC towards FleQ and away from NtrC binding sites, which had the dual benefit of improving flagellar-mediated motility and compensating for an over-active ntr system. These studies suggest that pre-existing links between GRNs, regulated by environmental signals, provided an opportunity for rewiring. It may also suggest that some ecological settings are more likely to result in selection of rewired GRNs than others through activation of promiscuous pathways that, to some extent, are already present.

It is possible that some TFs may be ‘primed’/more accessible for rewiring within a GRN, due to a combination of abundant activating signals and local network architecture that enables a high concentration of a TF in its active state. TCSs are simultaneously promiscuous and stringent [38]. Crosstalk is prevented by HK positive feedback loops, but mutations that remove repression or provide a source of constant activating signal are likely to permit promiscuity through runaway feedback leading to high regulator expression and activity [2,39–42] (Figure 1c(2)). These types of mutation are frequently seen in TCS rewiring events [20,37,43**,44]. Should promiscuous activity be advantageous, by increasing the flexibility and adaptability of GRNs, preexisting activating signals and architectures would likely constrain which regulators are available for evolutionary innovation in response to an environmental change, as these rewiring routes will be revealed to selection first without requiring time to acquire mutations that alter TF binding specificity. An important step towards unlocking promiscuity potential of a RR is increasing its concentration of active form in the cell; this can be achieved through mutation (Figure 1b(1) and c(2)) or through an increase in activating signal (Figure 1c(2)).

Flexible GRNs facilitate adaptation

Recent studies have highlighted rewiring of regulatory networks as an essential adaptive feature for some

examples of host colonisation and establishment. For example, Gopalan-Nair *et al.* found that adaptation of *Ralstonia solanacearum* to a resistant tomato cultivar led to convergent GRN evolution between independently evolved lines [45*]. Although each line acquired different mutations, they resulted in similar rewiring patterns of the virulence regulatory network and associated gene expression profiles. Similarly, Cottalorda *et al.* followed the evolution of 108 clinical *P. aeruginosa* isolates (starting from a single clone type) within the urinary tract between 48 and 488 days and found that adaptive mutations were preferentially located in genes encoding transcriptional regulators, TCSs, and carbon metabolism [46]. Host adaptation is a barrier not only faced by pathogens, but also symbionts. Pankey *et al.* followed the adaptations in *Vibrio fischerii* that enhanced colonisation and establishment in a squid host (*Euprymna scolopes*). They found multiple independent lines had point mutations in HK BinK that promoted traits known to support colonisation and immune evasion, but also altered quorum sensing that resulted in increased luminescence at lower cell densities [47**]. The fact that multiple traits were altered through mutations in one HK suggests that BinK might participate across more than one regulatory pathway that assist a symbiotic lifestyle.

These studies followed the adaptation of bacteria to a host, but what do we know about how a new pathogen emerges from an environmental ancestor? Bryant *et al.* followed the evolution of a recently emerging pathogenic species of nontuberculous mycobacterium, *Mycobacterium abscessus*, from a free-living origin [48*]. Rather than acquiring mutations in regulatory genes *de novo*, they found that horizontal gene transfer of **global transcriptional regulators** was an important mechanism for generating large scale phenotypic variation that created opportunities for adaptive change. Looking at the process in reverse, Yebra *et al.* [49*] discovered that genome remodelling in the bacterium *Staphylococcus aureus* subsp. *anaerobius* (which evolved from an *S. aureus* ancestor) accompanied its transition from a versatile multi-host bacterium to a fastidious niche-restricted pathogen of ruminants. In particular, acquisition of insertion sequence elements interfered with expression of downstream genes; reduced expression was predicted via altered promoter or DNA binding sequences of an inducer, and increased expression by elimination of repressor binding sites. These mechanisms effectively uncoupled target genes from their regulatory elements which enabled substantial rewiring of the genome through disruption of promoter regions and operon structure [49*].

GRNs also play an important role in antimicrobial resistance [50,51] and are a potential target for new antimicrobials. Patel *et al.* [43**] showed that in *Escherichia coli*, mutations in the feedback regulator protein MgrB preceded and facilitated the evolution of drug resistance to the antibiotic trimethoprim. Normally, MgrB attenuates PhoPQ through negative-feedback, however inactivation mutations in MgrB resulted in PhoPQ-mediated trimethoprim tolerance. Of particular interest is the role of the selective environment in the outcome of GRN adaptation under further exposure to trimethoprim. Under strong selection, MgrB mutants acquire resistance via additional mutations in dihydrofolate reductase; however, under weak selection, compensatory mutations in response to an overactive PhoPQ system inactivate its target RpoS [43**]. The role of the selective environment in determining the evolutionary trajectory of GRN rewiring events remains understudied and is of likely crucial importance to understanding bacterial adaptation to ecological shifts more generally.

Conclusions

GRNs are dynamical such that connections are forged and lost frequently across an evolutionary timescale. Recent research shows that GRN restructuring is an essential part of ecological niche switching and highlights the importance of rewiring in newly emerging pathogens and host adaptation. However, we know very little about the evolutionary drivers behind these rewiring events and the role of the environment in shaping GRN arrangement.

Crosstalk is necessary to reveal new TF-DNA binding interactions to selection, an essential process in GRN rewiring. A TF's binding affinity and position within a GRN are emerging as important factors for creating opportunity for crosstalk, but how these factors interplay to drive GRN evolution are far from clear. In particular, there is a lack of data that openly links the ecological setting with opportunity for GRN rewiring via 'tinkering' [52] of pre-existing non-cognate interactions. We have discussed the enormous range of binding potential between different TFs within a bacterium; as an example, *P. aeruginosa* shows the number of binding sites per TF range between 1 and 2407 [28]. We have also mentioned that features of GRN architecture, such as **auto-regulatory loops**, can make some GRNs easier targets for selection [40], especially when pre-existing links exist. But this leads to a number of unanswered questions that suggests exciting opportunities for research in this field: Why does such variation in the average number of TF binding sites exist, is this variation a legacy or a product of selection and what does it mean in combination with architecture and ecological conditions in terms of rewiring potential between GRNs? Combining insights from laboratory studies with natural datasets is an opportunity to uncover both the mechanisms and consequences of

genetic and ecological factors driving opportunity for crosstalk that may ultimately facilitate GRN rewiring and ecological niche-shifts.

Conflict of interest statement

Nothing declared.

Acknowledgements

The Royal Society provided support to T.B.T. (Grant no: DH150169) and M.J.S. (Grant no: RG160491, awarded to T.B.T.). R.W.J. is supported by The JABBS Foundation and BBSRC grants (BB/R014884/1 and BB/T010568/1).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. McAdams HH, Srinivasan B, Arkin AP: **The evolution of genetic regulatory systems in bacteria.** *Nat Rev Genet* 2004, **5**:169-178.
 2. Iglar C, Lagator M, Tkačik G, Bollback JP, Guet CC: **Evolutionary potential of transcription factors for gene regulatory rewiring.** *Nat Ecol Evol* 2018, **2**:1633-1643 210 2018.
 3. Isalan M, Lemerle C, Michalodimitrakis K, Horn C, Beltrao P, Raineri E, Garriga-Canut M, Serrano L: **Evolvability and hierarchy in rewired bacterial gene networks.** *Nature* 2008, **452**:840-845.
 4. Treangen TJ, Rocha EPC: **Horizontal transfer, not duplication, drives the expansion of protein families in prokaryotes.** *PLoS Genet* 2011, **7**.
 5. Huang H, Shao X, Xie Y, Wang T, Zhang Y, Wang X, Deng X: **An integrated genomic regulatory network of virulence-related transcriptional factors in *Pseudomonas aeruginosa*.** *Nat Commun* 2019, **10**:1-13.
 6. Gaballa A, Guariglia-Oropeza V, Wiedmann M, Boor KJ: **Cross talk between SigB and PrfA in *Listeria monocytogenes* facilitates transitions between extra- and intracellular environments.** *Microbiol Mol Biol Rev* 2019, **83**.
 7. Plitnick J, Chevance FFV, Stringer A, Hughes KT, Wade JT: **Regulatory cross talk between motility and interbacterial communication in *Salmonella enterica* serovar typhimurium.** *J Bacteriol* 2021, **203**.
 8. Luterbach CL, Mobley HLT: **Cross talk between MarR-like transcription factors coordinates the regulation of motility in uropathogenic *Escherichia coli*.** *Infect Immun* 2018, **86**.
 9. Willett JW, Tiwari N, Müller S, Hummels KR, Houtman JCD, Fuentes EJ, Kirby JR: **Specificity residues determine binding affinity for two-component signal transduction systems.** *mBio* 2013, **4**.
 10. Capra EJ, Perchuk BS, Skerker JM, Laub MT: **Adaptive mutations that prevent crosstalk enable the expansion of paralogous signaling protein families.** *Cell* 2012, **150**:222-232.
 11. Zarrinpar A, Park SH, Lim WA: **Optimization of specificity in a cellular protein interaction network by negative selection.** *Nature* 2003, **426**:676-680.
 12. Wagner A: **Adaptive gene misregulation.** *Genetics* 2021, **217**.
 13. Guzmán GI, Sandberg TE, LaCroix RA, Nyerges Á, Papp H, de Raad M, King ZA, Hefner Y, Northen TR, Notebaart RA *et al.*: **Enzyme promiscuity shapes adaptation to novel growth substrates.** *Mol Syst Biol* 2019, **15**:1-14.
 14. Pougach K, Voet A, Kondrashov FA, Voordeckers K, Christiaens JF, Baying B, Benes V, Sakai R, Aerts J, Zhu B *et al.*: **Duplication of a promiscuous transcription factor drives the emergence of a new regulatory network.** *Nat Commun* 2014, **5**.

15. Copley SD: **An evolutionary biochemist's perspective on promiscuity.** *Trends Biochem Sci* 2015, **40**:72-78.
16. Voordeckers K, Pougach K, Verstrepen KJ: **How do regulatory networks evolve and expand throughout evolution?** *Curr Opin Biotechnol* 2015, **34**:180-188.
17. Lamrabet O, Plumbridge J, Martin M, Lenski RE, Schneider D, Hindre T: **Plasticity of promoter-core sequences allows bacteria to compensate for the loss of a key global regulatory gene.** *Mol Biol Evol* 2019, **36**:1121-1133
- Utilising *E. coli* isolates from the LTEE, this study looks at adaptation following deletion of *crp* (a major global regulator) in four isolates with different transcriptional regulatory network structures. Genome resequencing revealed compensatory mutations were in promoter regions of two CRP target genes which restored transcription by modifying existing promoters and their regulation by TFs or by producing new promoters.
18. Yona AH, Alm EJ, Gore J: **Random sequences rapidly evolve into de novo promoters.** *Nat Commun* 2018, **9**:1-10.
19. Rosanova A, Colliva A, Osella M, Caselle M: **Modelling the evolution of transcription factor binding preferences in complex eukaryotes.** *Sci Rep* 2017, **7**:1-12.
20. Taylor TB, Mulley G, Dills AH, Alsohim AS, McGuffin LJ, Studholme DJ, Silby MW, Brockhurst MA, Johnson LJ, Jackson RW: **Evolutionary resurrection of flagellar motility via rewiring of the nitrogen regulation system.** *Science* 2015, **347**:1014-1017.
21. La Rosa R, Johansen HK, Molina S: **Convergent metabolic specialization through distinct evolutionary paths in *Pseudomonas aeruginosa*.** *mBio* 2018, **9**.
22. Perez-Rueda E, Hernandez-Guerrero R, Martinez-Nuñez MA, Armenta-Medina D, Sanchez I, Ibarra JA: **Abundance, diversity and domain architecture variability in prokaryotic DNA-binding transcription factors.** *PLoS One* 2018, **13**:1-16.
23. Kotelnikova EA, Makeev VJ, Gelfand MS: **Evolution of transcription factor DNA binding sites.** *Gene* 2005, **347**:255-263.
24. Harari O, Park SY, Huang H, Groisman EA, Zwir I: **Defining the plasticity of transcription factor binding sites by deconstructing DNA consensus sequences: the *phoP*-binding sites among gamma/enterobacteria.** *PLoS Comput Biol* 2010, **6**:10.
25. Barnes SL, Belliveau NM, Ireland WT, Kinney JB, Phillips R: **Mapping DNA sequence to transcription factor binding energy in vivo.** *PLoS Comput Biol* 2018, **15**:1-29.
26. Kurafeiski JD, Pinto P, Bornberg-Bauer E: **Evolutionary potential of cis-regulatory mutations to cause rapid changes in transcription factor binding.** *Genome Biol Evol* 2019, **11**:406-414
- This study investigates the binding behaviour of several hundred TFs across different species. They highlight multiple aspects of TF-binding interactions that determine promiscuous function. In particular, A/T rich binding motifs were bound by multiple different binding domains. This low binding specificity confers high evolvability and potential for genetic innovation.
27. Warman EA, Singh SS, Gubieda AG, Grainger DC: **A non-canonical promoter element drives spurious transcription of horizontally acquired bacterial genes.** *Nucleic Acids Res* 2020, **48**:4891-4901.
28. Wang T, Sun W, Fan L, Hua C, Wu N, Fan S, Zhang J, Deng X, Yan J: **An atlas of the binding specificities of transcription factors in *Pseudomonas aeruginosa* directs prediction of novel regulators in virulence.** *eLife* 2021, **10**.
29. Fan L, Wang T, Hua C, Sun W, Li X, Grunwald L, Liu J, Wu N, Shao X, Yin Y et al.: **A compendium of DNA-binding specificities of transcription factors in *Pseudomonas syringae*.** *Nat Commun* 2020, **11**:1-11
- Using HT-SELEX to identify binding motifs for 100 TFs, this study mapped functional interactions between virulence associated TFs and their targets. They found most TFs targeted fewer than 200 genes, but five bound more than 500 sites in the genome.
30. Friedlander T, Prizak R, Guet CC, Barton NH, Tkačik G: **Intrinsic limits to gene regulation by global crosstalk.** *Nat Commun* 2016, **7**:1-12.
31. Browning DF, Butala M, Busby SJW: **Bacterial transcription factors: regulation by pick "N" mix.** *J Mol Biol* 2019, **431**:4067-4077.
32. Jothi R, Balaji S, Wuster A, Grochow JA, Gsponer J, Przytycka TM, Aravind L, Babu MM: **Genomic analysis reveals a tight link between transcription factor dynamics and regulatory network architecture.** *Mol Syst Biol* 2009, **5**.
33. Nagy-Staron A, Tomasek K, Caruso Carter C, Sonnleitner E, Kavčić B, Paixão T, Guet CC: **Local genetic context shapes the function of a gene regulatory network.** *eLife* 2021, **10**
- Using a synthetic biology approach, this study moves transcriptional units (the coding sequence, promoter and terminator) to new locations within a GRN in *E. coli*, and measures the outcome on phenotype and gene expression. Moving the transcriptional units had dramatic impacts on phenotype, despite interactions within the GRN remaining the same.
34. Rapun-Araiz B, Haag AF, Solano C, Lasa I: **The impact of two-component sensorial network in staphylococcal speciation.** *Curr Opin Microbiol* 2020, **55**:40-47.
35. Capra EJ, Laub MT: **Evolution of two-component signal transduction systems.** *Annu Rev Microbiol* 2012, **66**:325-347.
36. Lozada-Chávez I, Janga SC, Collado-Vides J: **Bacterial regulatory networks are extremely flexible in evolution.** *Nucleic Acids Res* 2006, **34**:3434-3445.
37. Bretl DJ, Ladd KM, Atkinson SN, Müller S, Kirby JR: **Suppressor mutations reveal an NtrC-like response regulator, NmpR, for modulation of Type-IV Pili-dependent motility in *Myxococcus xanthus*.** *PLoS Genet* 2018, **14**.
38. Rowland MA, Deeds EJ: **Crosstalk and the evolution of specificity in two-component signaling.** *Proc Natl Acad Sci U S A* 2014, **111**:5550-5555.
39. Laub MT, Goulian M: **Specificity in two-component signal transduction pathways.** *Genet* 2007, **41**:121-145.
40. Baumstark R, Hänzelmann S, Tsuru S, Schaerli Y, Francesconi M, Mancuso FM, Castelo R, Isalan M: **The propagation of perturbations in rewired bacterial gene networks.** *Nat Commun* 2015, **6**:1-5.
41. Groisman EA: **Feedback control of two-component regulatory systems.** *Annu Rev Microbiol* 2016, **70**:103-124.
42. Xiong K, Gerstein M, Masel J: **Differences in evolutionary accessibility determine which equally effective regulatory motif evolves to generate pulses.** *Genetics* 2021, **219**.
43. Patel V, Matange N: **Adaptation and compensation in a bacterial gene regulatory network evolving under antibiotic selection.** *eLife* 2021, **10**
- This study highlights the role of selection pressure in determining the adaptive trajectory of mutations within a GRN. In *E. coli*, tolerance to trimethoprim evolves via mutations in *mgrB* which attenuates PhoPQ signalling. Under further strong selection, mutations in *folA* result in trimethoprim resistance. Under weak selection, compensatory mutations correct for reduced PhoPQ signalling and *E. coli* remains tolerant (but not resistant) to antibiotic exposure.
44. Chen Q, Stibitz S: **The *BvgASR* virulence regulon of *Bordetella pertussis*.** *Curr Opin Microbiol* 2019, **47**:74-81.
45. Gopalan-Nair R, Jardinaud M-F, Legrand L, Landry D, Barlet X, Lopez-Roques C, Vandecasteele C, Bouchez O, Genin S, Guidot A: **Convergent rewiring of the virulence regulatory network promotes adaptation of *Ralstonia solanacearum* on resistant tomato.** *Mol Biol Evol* 2020, **38**:1792-1808
- Using experimental evolution, the authors look at the adaptation of plant pathogen, *Ralstonia solanacearum*, to infect a resistant tomato cultivar. Over 300 generations they noticed that independently evolved clones had little genetic diversity between them, but across clones they showed convergent changes in gene expression.
46. Cottalorda A, Leoz M, Dahyot S, Gravey F, Grand M, Froidure T, Aujoulat F, Le Hello S, Jumas-Bilak E, Pestel-Caron M: **Within-host microevolution of *Pseudomonas aeruginosa* urinary**

isolates: a seven-patient longitudinal genomic and phenotypic study. *Front Microbiol* 2021, **11**.

47. Pankey MS, Foxall RL, Ster IM, Perry LA, Schuster BM, Donner RA, Coyle M, Cooper VS, Whistler CA: **Host-selected mutations converging on a global regulator drive an adaptive leap towards symbiosis in bacteria.** *eLife* 2017, **6**

This study uses experimental evolution to observe the adaptation of ecologically distinct strains of *Vibrio fischerii*, within the light organs of *Euprymna scolopes*. Adaptive mutations were all within the sensor kinase *binK* that promoted colonisation and immune invasion as well as altered quorum sensing. This suggests preexisting links exist between traits essential for symbiotic life history and provides a mechanism for rapid adaptation to a host.

48. Bryant JM, Brown KP, Burbaud S, Everall I, Belardinelli JM, Rodríguez-Rincon D, Grogono DM, Peterson CM, Verma D, Evans IE *et al.*: **Stepwise pathogenic evolution of *Mycobacterium abscessus*.** *Science* 2021, **372**

Exploring the transition of environmental mycobacterial species to a pathogen, this study conducts genomic analyses on the recent emerging pathogen *M. abscessus*. Initially, horizontal acquisition (of mainly genes with transcriptional effects) drove phenotypic changes with large effects.

Adaptation to the lung environment resulted in parallel evolution; genes under strongest selection were those involved in global regulation.

49. Yebra G, Haag AF, Neamah MM, Wee BA, Richardson EJ, Horcajo P, Granneman S, Tormo-Más MÁ, de la Fuente R, Fitzgerald JR *et al.*: **Radical genome remodelling accompanied the emergence of a novel host-restricted bacterial pathogen.** *PLoS Pathog* 2021, **17**:e1009606

This study explores extensive remodelling of the *S. aureus* genome that facilitates transition away from an opportunistic pathogen to a host-restricted lifestyle in *S. aureus* subsp. *anaerobius*. This life history transition was accompanied by multiple large insertion sequences that altered control of flanking genes critical for host establishment.

50. Tierney ARP, Rather PN: **Roles of two-component regulatory systems in antibiotic resistance.** *Future Microbiol* 2019, **14**:533-552.

51. Bhagirath AY, Li Y, Patidar R, Yerex K, Ma X, Kumar A, Duan K: **Two component regulatory systems and antibiotic resistance in Gram-negative pathogens.** *Int J Mol Sci* 2019, **20**:1781.

52. Jacob F: **Evolution and tinkering.** *Science* 1977, **196**:1161-1166.