

Common Data Elements for Disorders of Consciousness

Edlow, Brian L; Boerwinkle, Varina L; Annen, Jitka; Boly, Melanie; Gosseries, Olivia; Laureys, Steven; Mukherjee, Pratik; Puybasset, Louis; Stevens, Robert D; Threlkeld, Zachary D; Newcombe, Virginia F J; Fernandez-Espejo, Davinia; Curing Coma Campaign and its Contributing Members

DOI:

[10.1007/s12028-023-01794-2](https://doi.org/10.1007/s12028-023-01794-2)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Edlow, BL, Boerwinkle, VL, Annen, J, Boly, M, Gosseries, O, Laureys, S, Mukherjee, P, Puybasset, L, Stevens, RD, Threlkeld, ZD, Newcombe, VFJ, Fernandez-Espejo, D & Curing Coma Campaign and its Contributing Members 2023, 'Common Data Elements for Disorders of Consciousness: Recommendations from the Working Group on Neuroimaging', *Neurocritical Care*. <https://doi.org/10.1007/s12028-023-01794-2>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s12028-023-01794-2>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Common data elements for disorders of consciousness: Recommendations from the working group on neuroimaging

Brian L. Edlow^{1,2*}, Varina L. Boerwinkle^{3*}, Jitka Annen^{4,5}, Melanie Boly^{6,7}, Olivia Gosseries^{4,5}, Steven Laureys^{4,5}, Pratik Mukherjee⁸, Louis Puybasset⁹, Robert Stevens¹⁰, Zachary D. Threlkeld¹¹, Virginia F.J. Newcombe^{12**}, Davinia Fernandez-Espejo^{12**} and the Curing Coma Campaign Collaborators

* co-first authors

** co-senior authors

Author Affiliations

¹ Center for Neurotechnology and Neurorecovery, Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

² Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, MA, USA

³ Clinical Resting State fMRI Laboratory and Service, Department of Neurology, University of North Carolina School of Medicine in Chapel Hill, NC, USA

⁴ Coma Science Group, GIGA Consciousness, University of Liège, Liège, Belgium

⁵ Centre de Cerveau², University Hospital of Liège, Liège, Belgium

⁶ Departments of Neurology, University of Wisconsin, Madison, WI, USA

⁷ Wisconsin Institute for Sleep and Consciousness, Department of Psychiatry, University of Wisconsin-Madison, Madison, WI, USA

⁸ Department of Radiology and Biomedical Imaging, University of California, San Francisco, San Francisco, CA, USA

⁹ Department of Anesthesiology and Intensive Care, Groupe Hospitalier Pitié-Salpêtrière, Assistance Publique-Hôpitaux de Paris, Paris, France.

¹⁰ Departments of Anesthesiology and Critical Care Medicine, Neurology, Neurosurgery, Johns Hopkins University School of Medicine, Baltimore, MD, USA

¹¹ Department of Neurology, Stanford University School of Medicine, Stanford, CA, USA.

¹² University Division of Anaesthesia, University of Cambridge, Addenbrooke's Hospital Cambridge, Cambridge, UK

¹³ School of Psychology and Centre for Human Brain Health, University of Birmingham, Birmingham, Birmingham, UK

Corresponding author:

B.L. Edlow

bedlow@mgh.harvard.edu

Abstract word count: 253

Manuscript word count: 2273

Number of figures: 0

Number of tables: 0

Supplementary Material: Supplementary Appendix of Curing Coma Campaign Collaborators

Keywords: coma, consciousness, common data elements, neuroimaging

Details Page

Authorship Confirmation and Compliance with Journal Policies:

This manuscript has not been published or under consideration elsewhere. This manuscript complies with all instructions to authors. All authors have met requirements for authorship, as outlined below, and approved the final manuscript.

Author Contributions:

B.L.E., V.L.B., V.F.J.N., and D. F-E wrote the initial draft of the manuscript. All co-authors edited the manuscript and approved the final content. All co-authors contributed equally to the Case Report Forms released with the manuscript.

Conflicts of interest:

None

Acknowledgments:

The Curing Coma Campaign Collaborators are listed in the Supplementary Appendix.

Ethical Approval:

New data were not acquired or analyzed for this manuscript, and therefore there was no need for informed consent or approval from an Institutional Review Board.

Funding:

This work was supported by the NIH Director's Office (DP2HD101400), NIH National Institute for Neurological Disorders and Stroke (R21NS113037), James S. McDonnell Foundation, the European Union's Horizon 2020 Framework Programme for Research and Innovation under the Specific Grant Agreement No. 945539 (Human Brain Project SGA3).

Abstract

Background/Objective: Over the past five decades, advances in neuroimaging have yielded insights into the pathophysiologic mechanisms that cause disorders of consciousness (DoC) in patients with severe brain injuries. Structural, functional, metabolic, and perfusion imaging studies have revealed specific neuroanatomic regions, such as the brainstem tegmentum, thalamus, posterior cingulate cortex, medial prefrontal cortex, and occipital cortex, where lesions correlate with the current or future state of consciousness. Advanced imaging modalities such as diffusion tensor imaging (DTI), resting-state functional MRI (rs-fMRI), and task-based fMRI have been used to improve the accuracy of diagnosis and long-term prognosis, culminating in the endorsement of fMRI for the clinical evaluation of patients with DoC in the 2018 United States (task-based fMRI) and 2020 European (task-based and rs-fMRI) guidelines. As diverse neuroimaging techniques are increasingly used for patients with DoC in research and clinical settings, the need for a standardized approach to reporting results is clear. The success of future multicenter collaborations and international trials fundamentally depends upon the implementation of a shared nomenclature and infrastructure.

Methods: To address this need, the Neurocritical Care Society's *Curing Coma Campaign* convened a multidisciplinary panel of DoC neuroimaging experts to propose common data elements (CDEs) for data collection and reporting in this field.

Results: We report the recommendations of this CDE development panel and disseminate CDEs to be used in neuroimaging studies of patients with DoC.

Conclusions: These CDEs will support progress in the field of DoC neuroimaging and facilitate international collaboration.

Introduction

Neuroimaging is essential to the diagnostic and prognostic evaluation of patients with disorders of consciousness (DoC). Acutely in the intensive care unit (ICU), patients with DoC undergo neuroimaging tests to determine the mechanism of altered consciousness and the chances of long-term recovery [1]. In subacute rehabilitation hospitals and chronic nursing facilities, patients with prolonged DoC undergo neuroimaging tests to evaluate for secondary complications such as hydrocephalus and intracranial infections [2, 3]. Across the temporal continuum of DoC, neuroimaging tests are used to guide clinical management, inform prognosis, and support discussions with family members and surrogates about critical decisions such as the continuation of life-sustaining therapy [4, 5].

In the investigational domain, neuroimaging has advanced our understanding of the structural and functional basis of DoC [6, 7]. Volumetry and lesion mapping studies have identified neuroanatomic regions, such as the brainstem tegmentum [8-13], thalamus [14-17], posterior cingulate cortex [17, 18], medial prefrontal cortex [17], and occipital cortex [19], where lesions are associated with reduced levels of consciousness. Structural and functional connectivity studies have delineated brain networks implicated in the pathogenesis of DoC and have demonstrated that reemergence of acutely disrupted networks is associated with recovery of consciousness [20-34]. Furthermore, growing evidence indicates that diffusion MRI [35-37], resting-state functional MRI (rs-fMRI) [38-47], stimulus-based, task-based fMRI [48], and positron emission tomography (PET) [49] studies may predict functional outcomes in patients with DoC.

Advanced neuroimaging tools are also changing the diagnostic landscape for patients with DoC. Functional connectivity mapping with rs-fMRI may identify consciousness-suppressing seizure onset zones in deep brain regions that evade detection by scalp electroencephalography [50-55], raising the possibility that rs-fMRI could be used to identify treatable causes of DoC. PET studies have shown regions of preserved neuronal metabolism in patients who lack behavioral signs of consciousness [49, 56, 57]. Consistent with these PET findings, stimulus- and task-based fMRI studies have revealed cognitive function that evades detection on behavioral assessments [48, 58-62], leading to the creation of a new diagnostic category: covert consciousness (i.e., cognitive motor dissociation [63]). Meta-analyses indicate that 15-20% of patients with severe brain injury thought to be unconscious by clinical exam are actually covertly conscious [64, 65], prompting important new ethical questions about resource allocation and access to state-of-the-art diagnostic tests [66-68].

To address ethical concerns relating to the infrastructure, personnel and resources needed to acquire stimulus-based and task-based fMRI data, there is growing interest in phenotypic differentiation of DoC by rs-fMRI and diffusion MRI [69]. Stimulus-independent resting-state brain activity may ultimately prove to be more feasible for DoC evaluation in clinical settings, due to lack of need for additional task-based staffing, equipment, or reliance on patient mental status. While task-based fMRI is currently the only neuroimaging tool that can definitively detect covert consciousness, rs-fMRI may provide diagnostic information about the likelihood of covert consciousness [51, 70], as patients with complex patterns of functional brain connectivity are more likely to be covertly conscious [29]. Rs-fMRI connectivity also may be more robust than stimulus- and task-

based fMRI in patients receiving pharmacologic sedation, given that the effects of low-level pharmacologic sedation on functional connectivity are relatively small compared to the effect size of severe brain injury [71]. Structural connectivity mapping with diffusion MRI is also robust in the setting of sedation and may provide a complementary screening tool to identify patients with the potential for covert consciousness. Emerging evidence suggests that patients with covert consciousness have a structural connectivity phenotype characterized by disrupted connectivity in the primary motor cortex but preserved connectivity in the supplementary motor area and premotor cortex [72, 73]. Thus, together, structural and functional connectivity mapping techniques have potential to inform DoC patient triage for confirmatory assessments with task-based fMRI.

The translational impact of these neuroimaging discoveries is perhaps best evidenced by the endorsement of task-based fMRI to detect covert consciousness in the 2018 United States [74] and 2020 European [75] Guidelines for the clinical management of patients with DoC. Based on their diagnostic relevance and potential prognostic utility, neuroimaging tests that were once solely in the investigational domain are now being applied for clinical use in neonatal, pediatric, and adult patients [50, 51, 76-78]. Though global implementation has been limited to date [76], support for the clinical utility of advanced neuroimaging tests is increasing, even in countries where national insurance plans do not reliably reimburse for these tests [79]. While current UK guidelines [80] do not recommend advanced neuroimaging as part of standard clinical assessments, they acknowledge this may become a reality in the future.

Informed by this historical backdrop, the Neurocritical Care Society's *Curing Coma Campaign* [81] launched a Common Data Elements (CDE) initiative for DoC in 2020. This

CDE initiative is motivated by the recognition that ongoing progress depends on the development of harmonized and uniform data elements. Experience with other neurological diseases has demonstrated the benefit of collecting data in a systematic and consistent way, an approach championed by the National Institutes of Health (NIH), which provides CDEs for a range of neurological diseases (<https://www.commondataelements.ninds.nih.gov/>). To facilitate a similar CDE development process for patients with DoC, the *Curing Coma Campaign* convened 10 Work Groups to create CDEs for the broad spectrum of DoC research domains. Here, we report the results of the DoC CDE Neuroimaging Work Group. We aim for these neuroimaging CDEs to support progress in DoC neuroimaging and facilitate international collaboration.

Methods

Overview

Our goal was to create neuroimaging CDEs with the following characteristics:

- 1) capable of capturing the broad spectrum of findings reported to date in patients with DoC;
- 2) adaptable based on emerging evidence that might be reported in the future; and
- 3) feasible to implement in hospitals around the world.

Given the rapidly evolving landscape of DoC neuroimaging [6], the CDEs that we report here (version 1.0) are intended to be a starting point for future efforts by the international medical and scientific community to standardize neuroimaging studies. We expect that

the CDEs will be adapted and refined as additional neuroimaging discoveries emerge. These neuroimaging CDEs should be collected in conjunction with other relevant CDEs characterizing clinical characteristics and outcomes. Ultimately, we expect that these DoC neuroimaging CDEs will evolve with ongoing efforts to standardize data acquisition, analysis, and interpretation, with the long-term goal of improving care and outcomes for patients with DoC.

CDE Development Meetings

A 12-member Neuroimaging Work Group was convened as part of the *Curing Coma Campaign* to develop neuroimaging CDEs for patients with DoC. The Work Group met monthly online from 2021 to 2023, with the goal of creating neuroimaging CDEs for patients with DoC. Given that we aim to support innovative single and multi-center studies, we developed the CDEs to capture data from commonly available techniques (e.g., head CT and conventional MRI), as well as from advanced imaging techniques such as fMRI and DTI. Working group members with subspecialized knowledge were self-assigned to modality-specific case report forms (CRFs). Each CRF team, consisting of at least two Work Group members, developed the final product through internal consensus. The full Neuroimaging Work Group evaluated all CRFs for final approval and harmonization across modalities.

Adaptation of Established CDEs for Neuroimaging of DoC

We began by reviewing existing neuroimaging CDEs commissioned by the NINDS (<https://commondataelements.ninds.nih.gov>). Our goal was to leverage these existing

CDEs and, whenever possible, to use CDEs that were already defined according to established standards. These previously published CDEs provide the benefit of user familiarity and prior vetting by neuroimaging experts [82-84].

Consistent with previously published CDEs, we organized the DoC neuroimaging CDEs into CRFs by imaging techniques. Techniques were eligible for inclusion based on prespecified criteria: 1) routine acquisition of the technique in clinical practice; or 2) at least one publication describing the investigational use of the technique in patients with DoC. Importantly, most previously published neuroimaging CDEs pertain to specific neurological diseases (e.g., TBI, ischemic stroke, subarachnoid hemorrhage, COVID-19) [82-85]. DoC, by contrast, represent the spectrum of a syndrome associated with many neurological disorders and types of brain injury. As such, we selected previously published, disease-specific CDEs when relevant, and we proposed new CDEs that capture the unique neuroimaging considerations associated with the DoC patient population, across the age spectrum from neonatal through adulthood.

Proposal for New DoC Neuroimaging CDEs

For neuroimaging techniques described in DoC publications that were not accounted for by previously published CDEs, we created new CDEs based on consensus opinion. We aimed to provide investigators with the flexibility to thoroughly characterize all neuroimaging findings, regardless of brain injury etiology.

Classifying the Pathophysiologic Association of Imaging Findings with DoC

We also provide investigators with an opportunity to enter data about presumed mechanisms of neurological injury and their relatedness to DoC, consistent with recently proposed neuroimaging CDEs for patients with COVID-19 [85]. At the end of each CRF, we created a new CDE pertaining to the presumed pathophysiological cause of the imaging findings. Such data will facilitate epidemiologic and mechanistic studies of DoC, while also providing hypothesis-generating data for future investigations.

CDE Classification

All CDEs were classified as “disease core”, “basic”, “supplemental”, or “exploratory” based on the consensus opinion of the Work Group. This classification nomenclature is consistent with that used in prior NINDS CDE initiatives [82-84]. We assigned the “disease core” designation to all CDEs that are required for all DoC studies. Limiting the number of disease core CDEs was intended to reduce the burden of data entry, which can lead to incomplete CRFs and reduced participation in multi-center international trials. We assigned the “basic” designation to CDEs that are strongly recommended for all DoC studies. We assigned the “supplemental” designation to CDEs that are recommended for specific DoC studies (i.e., depending on the context and goals of the study), and the “exploratory” designation was applied to CDEs that can be considered for use in DoC neuroimaging studies but require further validation. Finally, we assigned the designation “key design element (KDE)” to any methodological parameter that is relevant to the acquisition, processing, or analysis of data.

Results

Adaptation of Previously Proposed CDEs to DoC

The neuroimaging CDEs previously proposed by the NIH that were most relevant to DoC included those developed for ischemic stroke [83], traumatic brain injury [82] and subarachnoid hemorrhage [84]. Based on these previously developed CDEs, we created eight CRFs, each representing a neuroimaging technique: 1) head CT; 2) conventional MRI; 3) T1 volumetrics; 4) diffusion MRI; 5) perfusion imaging (CT and MRI); 6) fMRI (resting-state, passive stimulus-based, and active task-based); 7) PET (resting-state, passive stimulus-based, and active task-based); and 8) MR spectroscopy. These eight CRFs include basic and supplemental CDEs. Exploratory CDEs were not identified. A separate CRF was created for disease core CDEs, which includes CDEs from all Work Groups.

The Neuroimaging Work Group identified additional priorities for the international DoC neuroimaging community that are considered synergistic with the present CDE effort but beyond the scope of the CRFs. Specifically, the Neuroimaging Work Group aims to encourage investigators to: 1) publicly disseminate all code and data processing scripts; 2) openly share data; and 3) label data files using the standard BIDS format [86] to facilitate data pooling.

Dissemination of CDEs for DoC Neuroimaging

We release version 1.0 of the proposed neuroimaging CDEs for patients with DoC as a set of eight CRFs (see Supplementary Materials [Authors' note to Editors/Reviewers: we also plan to disseminate the CRFs on www.zenodo.org after peer review]). The CDEs

underwent a 2-month public feedback period from October to November 2022, which was advertised at the 2022 Neurocritical Care Society annual meeting and via Twitter. Public feedback was received and incorporated into the final CRFs. For the neuroimaging CDEs, feedback pertained to the style and formatting of the CRFs, though no specific content-related changes were recommended.

We encourage ongoing feedback regarding modifications to the CDEs, which can be submitted via email to cde.curingcoma@gmail.com. Suggestions to edit or add to the current list of CDEs will be evaluated by the Neuroimaging Work Group on an as-needed basis, and changes to the CRFs will be posted on the zenodo website with new version numbers. We are committed to an adaptive approach based on emerging evidence, with rapid distribution of modifications using online scientific portals.

Discussion

Global collaboration and data reporting standardization are essential to advance knowledge and improve care for patients with DoC. To support this effort, the *Curing Coma Campaign* convened Work Groups to develop CDEs for DoC research. Here, we disseminate the neuroimaging CDEs that emerged from this international initiative. We designed the DoC neuroimaging CDEs to be widely accessible and practical to implement at both academic medical centers and community hospitals. The DoC neuroimaging CDEs also leverage previous CDE efforts supported by NINDS, ensuring consistency with prior reported efforts to standardize neuroimaging data acquisition. Newly proposed CDEs specific to patients with DoC were added based on a review of DoC neuroimaging studies. All DoC neuroimaging CDEs, organized in eight modality-specific CRFs, are now

publicly available at (see Supplementary Materials [Authors' note to Editors/Reviewers: we also plan to disseminate the CRFs on www.zenodo.org after peer review]).

The CDEs proposed here will support ongoing efforts to identify signatures of atypical (pathological and disrupted) and preserved brain networks [69] and create new opportunities for personalized medicine and targeted therapies [51, 87, 88]. We also expect that these neuroimaging CDEs will support studies that shed new light on fundamental questions about DoC pathophysiology, such as “what is the neuroanatomic basis of covert consciousness?” [72] and “are the neural correlates of consciousness localized to the anterior or posterior regions of the cerebral cortex?” [18]. Furthermore, the CDEs are designed to support large, multicenter studies that test the diagnostic and prognostic utility of advanced imaging techniques, which will be essential for clinical translation. Finally, these CDEs will create new opportunities for personalized medicine by guiding the selection of targeted therapies aimed at promoting recovery of consciousness [51, 87, 88].

This CDE development effort is a dynamic process, and we anticipate revisions that reflect ongoing progress in the field of DoC neuroimaging. Only with a comprehensive, global commitment to data reporting standardization and data sharing can the international community advance knowledge and improve care for patients with DoC.

References

1. Edlow, B.L., et al., *Recovery from disorders of consciousness: mechanisms, prognosis and emerging therapies*. Nat Rev Neurol, 2021. **17**(3): p. 135-156.
2. Arnts, H., et al., *The Dilemma of Hydrocephalus in Prolonged Disorders of Consciousness*. J Neurotrauma, 2020. **37**(20): p. 2150-2156.
3. Wilson, M.R. and K.L. Roos, *Infectious diseases and impaired consciousness*. Neurol Clin, 2011. **29**(4): p. 927-42.
4. Young, M.J. and A. Peterson, *Neuroethics across the Disorders of Consciousness Care Continuum*. Semin Neurol, 2022. **42**(3): p. 375-392.
5. Fins, J.J., *The ethics of measuring and modulating consciousness: the imperative of minding time*. Prog Brain Res, 2009. **177**: p. 371-82.
6. Sanz, L.R.D., et al., *Update on neuroimaging in disorders of consciousness*. Curr Opin Neurol, 2021. **34**(4): p. 488-496.
7. Fischer, D., et al., *Applications of advanced MRI to disorders of consciousness*. Semin Neurol, 2022.
8. Parvizi, J. and A.R. Damasio, *Neuroanatomical correlates of brainstem coma*. Brain, 2003. **126**(Pt 7): p. 1524-36.
9. Edlow, B.L., et al., *Disconnection of the ascending arousal system in traumatic coma*. J Neuropathol Exp Neurol, 2013. **72**(6): p. 505-23.
10. Fischer, D.B., et al., *A human brain network derived from coma-causing brainstem lesions*. Neurology, 2016. **87**(23): p. 2427-2434.
11. Izzy, S., et al., *Revisiting Grade 3 Diffuse Axonal Injury: Not All Brainstem Microbleeds are Prognostically Equal*. Neurocrit Care, 2017. **27**(2): p. 199-207.
12. Snider, S.B., et al., *Cortical lesions causing loss of consciousness are anticorrelated with the dorsal brainstem*. Hum Brain Mapp, 2020. **41**(6): p. 1520-1531.
13. Bianciardi, M., et al., *Location of Subcortical Microbleeds and Recovery of Consciousness After Severe Traumatic Brain Injury*. Neurology, 2021. **97**(2): p. e113-e123.
14. Lutkenhoff, E.S., et al., *Thalamic and extrathalamic mechanisms of consciousness after severe brain injury*. Ann Neurol, 2015. **78**(1): p. 68-76.
15. Fernandez-Espejo, D., et al., *Reductions of thalamic volume and regional shape changes in the vegetative and the minimally conscious states*. J Neurotrauma, 2010. **27**(7): p. 1187-93.
16. Fernandez-Espejo, D., et al., *Diffusion weighted imaging distinguishes the vegetative state from the minimally conscious state*. Neuroimage, 2011. **54**(1): p. 103-12.
17. Annen, J., et al., *Regional brain volumetry and brain function in severely brain-injured patients*. Ann Neurol, 2018. **83**(4): p. 842-853.
18. Boly, M., et al., *Are the Neural Correlates of Consciousness in the Front or in the Back of the Cerebral Cortex? Clinical and Neuroimaging Evidence*. J Neurosci, 2017. **37**(40): p. 9603-9613.
19. Snider, S.B., et al., *Regional Distribution of Brain Injury After Cardiac Arrest: Clinical and Electrographic Correlates*. Neurology, 2022. **98**(12): p. e1238-e1247.

20. Vanhaudenhuyse, A., et al., *Default network connectivity reflects the level of consciousness in non-communicative brain-damaged patients*. Brain, 2010. **133**(Pt 1): p. 161-71.
21. Newcombe, V.F., et al., *Aetiological differences in neuroanatomy of the vegetative state: insights from diffusion tensor imaging and functional implications*. J Neurol Neurosurg Psychiatry, 2010. **81**(5): p. 552-61.
22. Fernandez-Espejo, D., et al., *Combination of diffusion tensor and functional magnetic resonance imaging during recovery from the vegetative state*. BMC Neurol, 2010. **10**: p. 77.
23. Norton, L., et al., *Disruptions of functional connectivity in the default mode network of comatose patients*. Neurology, 2012. **78**(3): p. 175-81.
24. Fernandez-Espejo, D., et al., *A role for the default mode network in the bases of disorders of consciousness*. Ann Neurol, 2012. **72**(3): p. 335-43.
25. Demertzi, A., et al., *Intrinsic functional connectivity differentiates minimally conscious from unresponsive patients*. Brain, 2015. **138**(Pt 9): p. 2619-31.
26. Thengone, D.J., et al., *Local changes in network structure contribute to late communication recovery after severe brain injury*. Sci Transl Med, 2016. **8**(368): p. 368re5.
27. Sair, H.I., et al., *Early Functional Connectome Integrity and 1-Year Recovery in Comatose Survivors of Cardiac Arrest*. Radiology, 2018. **287**(1): p. 247-255.
28. Threlkeld, Z.D., et al., *Functional networks reemerge during recovery of consciousness after acute severe traumatic brain injury*. Cortex, 2018. **106**: p. 299-308.
29. Demertzi, A., et al., *Human consciousness is supported by dynamic complex patterns of brain signal coordination*. Sci Adv, 2019. **5**(2): p. eaat7603.
30. Spindler, L.R.B., et al., *Dopaminergic brainstem disconnection is common to pharmacological and pathological consciousness perturbation*. Proc Natl Acad Sci U S A, 2021. **118**(30).
31. Amiri, M., et al., *Multimodal prediction of residual consciousness in the intensive care unit: the CONNECT-ME study*. Brain, 2023. **146**(1): p. 50-64.
32. Snider, S.B., et al., *Ascending arousal network connectivity during recovery from traumatic coma*. Neuroimage Clin, 2020. **28**: p. 102503.
33. Annen, J., et al., *Function-structure connectivity in patients with severe brain injury as measured by MRI-DWI and FDG-PET*. Hum Brain Mapp, 2016. **37**(11): p. 3707-3720.
34. Panda, R., et al., *Disruption in structural-functional network repertoire and time-resolved subcortical fronto-temporoparietal connectivity in disorders of consciousness*. Elife, 2022. **11**.
35. Galanaud, D., et al., *Assessment of white matter injury and outcome in severe brain trauma: a prospective multicenter cohort*. Anesthesiology, 2012. **117**(6): p. 1300-10.
36. Velly, L., et al., *Use of brain diffusion tensor imaging for the prediction of long-term neurological outcomes in patients after cardiac arrest: a multicentre, international, prospective, observational, cohort study*. Lancet Neurol, 2018. **17**(4): p. 317-326.
37. Puybasset, L., et al., *Prognostic value of global deep white matter DTI metrics for 1-year outcome prediction in ICU traumatic brain injury patients: an MRI-COMA and CENTER-TBI combined study*. Intensive Care Med, 2022. **48**(2): p. 201-212.

38. Norton, L., et al., *Disruptions of functional connectivity in the default mode network of comatose patients*. *Neurology*, 2012. **78**: p. 175-81.
39. Koenig, M.A., et al., *MRI default mode network connectivity is associated with functional outcome after cardiopulmonary arrest*. *Neurocrit Care*, 2014. **20**(3): p. 348-57.
40. Silva, S., et al., *Disruption of posteromedial large-scale neural communication predicts recovery from coma*. *Neurology*, 2015. **85**: p. 1-9.
41. Song, M., et al., *Prognostication of chronic disorders of consciousness using brain functional networks and clinical characteristics*. *Elife*, 2018. **7**.
42. Sair, H.I., et al., *Early functional connectome integrity and 1-year recovery in comatose survivors of cardiac arrest*. *Radiology*, 2018. **287**: p. 247-255.
43. Guo, H., et al., *Evaluation of Prognosis in Patients with Severe Traumatic Brain Injury Using Resting-State Functional Magnetic Resonance Imaging*. *World Neurosurgery*, 2019. **121**: p. e630-e639.
44. Yu, Y., et al., *A multi-domain prognostic model of disorder of consciousness using resting-state fMRI and laboratory parameters*. *Brain Imaging and Behavior*, 2020.
45. Pugin, D., et al., *Resting-state brain activity for early prediction outcome in postanoxic patients in a coma with indeterminate clinical prognosis*. *Am J Neuroradiol*, 2020: p. 1-9.
46. Peran, P., et al., *Functional and Structural Integrity of Frontoparietal Connectivity in Traumatic and Anoxic Coma*. *Crit Care Med*, 2020.
47. Fischer, D., et al., *Intact Brain Network Function in an Unresponsive Patient with COVID-19*. *Ann Neurol*, 2020. **88**(4): p. 851-854.
48. Norton, L., et al., *Functional Neuroimaging as an Assessment Tool in Critically Ill Patients*. *Ann Neurol*, 2023. **93**(1): p. 131-141.
49. Thibaut, A., et al., *Preservation of Brain Activity in Unresponsive Patients Identifies MCS Star*. *Ann Neurol*, 2021. **90**(1): p. 89-100.
50. Boerwinkle, V.L., et al., *Association of network connectivity via resting state functional MRI with consciousness, mortality, and outcomes in neonatal acute brain injury*. *Neuroimage Clin*, 2022. **34**: p. 102962.
51. Boerwinkle, V.L., et al., *Resting-state fMRI in disorders of consciousness to facilitate early therapeutic intervention*. *Neurol Clin Pract*, 2019. **9**(4): p. e33-e35.
52. Boerwinkle, V.L., et al., *Correlating Resting-State Functional Magnetic Resonance Imaging Connectivity by Independent Component Analysis-Based Epileptogenic Zones with Intracranial Electroencephalogram Localized Seizure Onset Zones and Surgical Outcomes in Prospective Pediatric Intractable Epilepsy Study*. *Brain Connect*, 2017. **7**(7): p. 424-442.
53. Chakraborty, A.R., et al., *Resting-state functional magnetic resonance imaging with independent component analysis for presurgical seizure onset zone localization: A systematic review and meta-analysis*. *Epilepsia*, 2020. **61**(9): p. 1958-1968.
54. Guo, J.N., et al., *Impaired consciousness in patients with absence seizures investigated by functional MRI, EEG, and behavioural measures: a cross-sectional study*. *Lancet Neurol*, 2016. **15**(13): p. 1336-1345.
55. Fischer, D., et al., *Ictal fMRI: Mapping Seizure Topography with Rhythmic BOLD Oscillations*. *Brain Sci*, 2022. **12**(12).

56. Stender, J., et al., *Diagnostic precision of PET imaging and functional MRI in disorders of consciousness: a clinical validation study*. Lancet, 2014. **384**(9942): p. 514-22.
57. Menon, D.K., et al., *Cortical processing in persistent vegetative state*. Wolfson Brain Imaging Centre Team. Lancet, 1998. **352**(9123): p. 200.
58. Owen, A.M., et al., *Detecting awareness in the vegetative state*. Science, 2006. **313**(5792): p. 1402.
59. Coleman, M.R., et al., *Towards the routine use of brain imaging to aid the clinical diagnosis of disorders of consciousness*. Brain, 2009. **132**(Pt 9): p. 2541-52.
60. Monti, M.M., et al., *Willful modulation of brain activity in disorders of consciousness*. N Engl J Med, 2010. **362**(7): p. 579-89.
61. Naci, L. and A.M. Owen, *Making every word count for nonresponsive patients*. JAMA Neurol, 2013. **70**(10): p. 1235-41.
62. Edlow, B.L., et al., *Early detection of consciousness in patients with acute severe traumatic brain injury*. Brain, 2017. **140**(9): p. 2399-2414.
63. Schiff, N.D., *Cognitive Motor Dissociation Following Severe Brain Injuries*. JAMA Neurol, 2015. **72**(12): p. 1413-5.
64. Kondziella, D., et al., *Preserved consciousness in vegetative and minimal conscious states: systematic review and meta-analysis*. J Neurol Neurosurg Psychiatry, 2016. **87**(5): p. 485-92.
65. Schnakers, C., et al., *Covert Cognition in Disorders of Consciousness: A Meta-Analysis*. Brain Sci, 2020. **10**(12).
66. Fins, J.J. and J.L. Bernat, *Ethical, palliative, and policy considerations in disorders of consciousness*. Neurology, 2018. **91**(10): p. 471-475.
67. Young, M.J. and B.L. Edlow, *The Quest for Covert Consciousness: Bringing Neuroethics to the Bedside*. Neurology, 2021. **96**(19): p. 893-896.
68. Peterson, A., S. Aas, and D. Wasserman, *What Justifies the Allocation of Health Care Resources to Patients with Disorders of Consciousness?* AJOB Neurosci, 2021. **12**(2-3): p. 127-139.
69. Kondziella, D., et al., *A precision medicine framework for classifying patients with disorders of consciousness: Advanced Classification of Consciousness Endotypes (ACCESS)*. Neurocrit Care, 2021. **35**(Suppl 1): p. 27-36.
70. Boerwinkle, V.L. *Patient Stories: Road to Recovery*. World Coma Day. March 22, 2022. Neurocritical Care Society. <https://bit.ly/Peds-CC1>. 2022.
71. Kirsch, M., et al., *Sedation of Patients With Disorders of Consciousness During Neuroimaging: Effects on Resting State Functional Brain Connectivity*. Anesth Analg, 2017. **124**(2): p. 588-598.
72. Fernandez-Espejo, D., S. Rossit, and A.M. Owen, *A Thalamocortical Mechanism for the Absence of Overt Motor Behavior in Covertly Aware Patients*. JAMA Neurol, 2015. **72**(12): p. 1442-50.
73. Stafford, C.A., A.M. Owen, and D. Fernandez-Espejo, *The neural basis of external responsiveness in prolonged disorders of consciousness*. Neuroimage Clin, 2019. **22**: p. 101791.
74. Giacino, J.T., et al., *Practice guideline update recommendations summary: Disorders of consciousness: Report of the Guideline Development, Dissemination, and*

- Implementation Subcommittee of the American Academy of Neurology; the American Congress of Rehabilitation Medicine; and the National Institute on Disability, Independent Living, and Rehabilitation Research. Neurology, 2018. 91(10): p. 450-460.*
75. Kondziella, D., et al., *European Academy of Neurology guideline on the diagnosis of coma and other disorders of consciousness. Eur J Neurol, 2020. 27(5): p. 741-756.*
 76. Helbok, R., et al., *The Curing Coma Campaign International Survey on Coma Epidemiology, Evaluation, and Therapy (COME TOGETHER). Neurocrit Care, 2022.*
 77. Edlow, B.L. and J.J. Fins, *Assessment of Covert Consciousness in the Intensive Care Unit: Clinical and Ethical Considerations. J Head Trauma Rehabil, 2018. 33(6): p. 424-434.*
 78. Young, M.J., et al., *Toward Uniform Insurer Coverage for Functional MRI Following Severe Brain Injury. J Head Trauma Rehabil, 2023.*
 79. Scolding, N., A.M. Owen, and J. Keown, *Prolonged disorders of consciousness: a critical evaluation of the new UK guidelines. Brain, 2021. 144(6): p. 1655-1660.*
 80. Physicians, R.C.o., *Prolonged disorders of consciousness following sudden onset brain injury. National Clinical Guidelines. London: Royal College of Physicians, 2020.*
 81. Provencio, J.J., et al., *The Curing Coma Campaign: Framing Initial Scientific Challenges- Proceedings of the First Curing Coma Campaign Scientific Advisory Council Meeting. Neurocrit Care, 2020. 33(1): p. 1-12.*
 82. Haacke, E.M., et al., *Common data elements in radiologic imaging of traumatic brain injury. J Magn Reson Imaging, 2010. 32(3): p. 516-43.*
 83. Saver, J.L., et al., *Standardizing the structure of stroke clinical and epidemiologic research data: the National Institute of Neurological Disorders and Stroke (NINDS) Stroke Common Data Element (CDE) project. Stroke, 2012. 43(4): p. 967-73.*
 84. Hackenberg, K.A.M., et al., *Common Data Elements for Radiological Imaging of Patients with Subarachnoid Hemorrhage: Proposal of a Multidisciplinary Research Group. Neurocrit Care, 2019. 30(Suppl 1): p. 60-78.*
 85. Edlow, B.L., et al., *Common Data Elements for COVID-19 Neuroimaging: A GCS-NeuroCOVID Proposal. Neurocrit Care, 2021. 34(2): p. 365-370.*
 86. Gorgolewski, K.J., et al., *The brain imaging data structure, a format for organizing and describing outputs of neuroimaging experiments. Sci Data, 2016. 3: p. 160044.*
 87. Edlow, B.L., et al., *Personalized Connectome Mapping to Guide Targeted Therapy and Promote Recovery of Consciousness in the Intensive Care Unit. Neurocrit Care, 2020. 33(2): p. 364-375.*
 88. Fridman, E.A., et al., *Presynaptic dopamine deficit in minimally conscious state patients following traumatic brain injury. Brain, 2019. 142(7): p. 1887-1893.*