

Effect of Metformin Versus Placebo on New Primary Cancers in Canadian Cancer Trials Group MA.32

Goodwin, Pamela J; Chen, Bingshu E; Gelmon, Karen A; Whelan, Timothy J; Ennis, Marguerite; Lemieux, Julie; Ligibel, Jennifer A; Hershman, Dawn L; Mayer, Ingrid A; Hobday, Timothy J; Bliss, Judith M; Rastogi, Priya; Rabaglio-Poretti, Manuela; Thompson, Alastair M; Rea, Daniel; Stos, Paul M; Shepherd, Lois E; Stambolic, Vuk; Parulekar, Wendy R

DOI:
[10.1200/JCO.23.00296](https://doi.org/10.1200/JCO.23.00296)

License:
Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Goodwin, PJ, Chen, BE, Gelmon, KA, Whelan, TJ, Ennis, M, Lemieux, J, Ligibel, JA, Hershman, DL, Mayer, IA, Hobday, TJ, Bliss, JM, Rastogi, P, Rabaglio-Poretti, M, Thompson, AM, Rea, D, Stos, PM, Shepherd, LE, Stambolic, V & Parulekar, WR 2023, 'Effect of Metformin Versus Placebo on New Primary Cancers in Canadian Cancer Trials Group MA.32: A Secondary Analysis of a Phase III Randomized Double-Blind Trial in Early Breast Cancer', *Journal of Clinical Oncology*. <https://doi.org/10.1200/JCO.23.00296>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.














When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Effect of Metformin Versus Placebo on New Primary Cancers in Canadian Cancer Trials Group MA.32: A Secondary Analysis of a Phase III Randomized Double-Blind Trial in Early Breast Cancer

Pamela J. Goodwin, MD¹ ; Bingshu E. Chen, PhD² ; Karen A. Gelmon, MD³ ; Timothy J. Whelan, MD⁴ ; Marguerite Ennis, PhD⁵ ; Julie Lemieux, MD⁶; Jennifer A. Ligibel, MD⁷ ; Dawn L. Hershman, MD⁸ ; Ingrid A. Mayer, MD⁹ ; Timothy J. Hobday, MD¹⁰ ; Judith M. Bliss, MSc¹¹ ; Priya Rastogi, MD¹²; Manuela Rabaglio-Poretti, MD¹³; Alastair M. Thompson, MD¹⁴; Daniel W. Rea, PhD¹⁵ ; Paul M. Stos, BSc (Hon)²; Lois E. Shepherd, MD² ; Vuk Stambolic, PhD^{16,17}; and Wendy R. Parulekar, MD² 

DOI <https://doi.org/10.1200/JCO.23.00296>

ABSTRACT

Clinical trials frequently include multiple end points that mature at different times. The initial report, typically based on the primary end point, may be published when key planned coprimary or secondary analyses are not yet available. Clinical trial updates provide an opportunity to disseminate additional results from studies, published in JCO or elsewhere, for which the primary end point has already been reported.

Metformin has been associated with lower cancer risk in epidemiologic and preclinical research. In the MA.32 randomized adjuvant breast cancer trial, metformin (v placebo) did not affect invasive disease-free or overall survival. Here, we report metformin effects on the risk of new cancer. Between 2010 and 2013, 3,649 patients with breast cancer younger than 75 years without diabetes with high-risk T1-3, N0-3 M0 breast cancer (any estrogen receptor, progesterone receptor, human epidermal growth factor receptor 2) were randomly assigned to metformin 850 mg orally twice a day or placebo twice a day for 5 years. New primary invasive cancers (outside the ipsilateral breast) developing as a first event were identified. Time to events was described by the competing risks method; two-sided likelihood ratio tests adjusting for age, BMI, smoking, and alcohol intake were used to compare metformin versus placebo arms. A total of 184 patients developed new invasive cancers: 102 metformin and 82 placebo, hazard ratio (HR), 1.25; 95% CI, 0.94 to 1.68; $P = .13$. These included 48 contralateral invasive breast cancers (27 metformin v 21 placebo), HR, 1.29; 95% CI, 0.72 to 2.27; $P = .40$ and 136 new nonbreast primary cancers (75 metformin v 61 placebo), HR, 1.24; 95% CI, 0.88 to 1.74; $P = .21$. Metformin did not reduce the risk of new cancer development in these nondiabetic patients with breast cancer.

ACCOMPANYING CONTENT

 Appendix

 Protocol

Accepted July 20, 2023

Published September 11, 2023

J Clin Oncol 00:1-7

© 2023 by American Society of
Clinical Oncology



View Online
Article

Creative Commons Attribution
Non-Commercial No Derivatives
4.0 License

INTRODUCTION

Metformin, a biguanide commonly used to treat type 2 diabetes, has been associated with lower overall cancer risk in epidemiologic studies conducted in patients with diabetes.^{1,2} Meta-analyses of these studies have inconsistently identified lower risk of some types of cancer when individuals receive metformin.³⁻¹⁴ It has been suggested that data from observational studies of metformin and cancer risk seem largely unreliable, studies using designs least likely to be affected by bias (notably selection and time-related biases) being most likely to yield results that did not support a causal effect of metformin on cancer risk.¹⁵⁻¹⁷ The Diabetes Prevention Program trial, which randomly assigned adults with prediabetes to metformin, a lifestyle-based weight loss intervention or placebo, reported no effect of metformin

versus placebo on cancer mortality at 21-year follow-up (hazard ratio [HR], 1.04; 95% CI, 0.72 to 1.52; $P = .83$).¹⁸

Current understanding of metformin action includes systemic effects resulting from reduced insulin signaling through PI3K/RAS pathways and cancer cell direct effects (notably LKB1-mediated activation of AMPK, a negative regulator of PI3K/Akt/mTOR signaling and protein synthesis). Some preclinical research suggests that metformin may lower cancer risk; however, translation to the clinical setting is not straightforward.^{19,20}

Here, we report the effect of metformin versus placebo on new invasive cancers in MA.32, a phase III adjuvant randomized trial of metformin versus placebo in high-risk T1-3, N0-2, M0 breast cancer. We have previously reported

TABLE 1. Baseline Patient and Tumor Characteristics

Characteristic	Metformin (n = 1,824)	Placebo (n = 1,825)	Total (N = 3,649)
Age, years			
Mean (SD)	52.2 (10.1)	52.6 (10.1)	52.4 (10.1)
BMI, kg/m ²			
Mean (SD)	28.7 (6.6)	28.5 (6.2)	28.6 (6.4)
Sex, No. (%)			
Female	1,821 (99.8)	1,822 (99.8)	3,543 (99.8)
Male	3 (0.2)	3 (0.2)	6 (0.2)
Race, No. (%)			
Asian	51 (2.8)	48 (2.6)	99 (2.7)
Black or African American	83 (4.6)	84 (4.6)	167 (4.6)
American Indian or Alaska Native	9 (0.5)	8 (0.4)	17 (0.5)
Native Hawaiian or Pacific Islander	5 (0.3)	8 (0.4)	13 (0.4)
Non-Hispanic White	1,569 (86.0)	1,557 (85.3)	3,126 (85.7)
Hispanic	84 (4.6)	94 (5.2)	178 (4.9)
Smoking history, No. (%)			
Current smoker	151 (8.3)	142 (7.8)	293 (8.0)
Past smoker	793 (43.5)	756 (41.4)	1,549 (42.4)
Never-smoker	880 (48.2)	927 (50.8)	1,807 (49.5)
Current alcohol consumption, No. (%)			
Yes	1,231 (72.4)	1,320 (72.3)	2,641 (72.4)
No	503 (27.6)	505 (27.7)	1,008 (27.6)
Not reported (or refused) or unknown	23 (1.3)	26 (1.4)	49 (1.3)
Menopausal status, No. (%)			
Premenopausal ^a	714 (39.1)	698 (38.2)	1,412 (38.7)
Postmenopausal ^b	1,110 (60.1)	1,127 (61.8)	2,237 (61.3)
T stage, No. (%)			
T1	620 (34.0)	619 (33.9)	1,239 (34.0)
T2	969 (53.1)	974 (53.4)	1,943 (53.2)
T3	234 (12.8)	232 (12.7)	466 (12.1)
T4	1 (0.0)	0 (0.0)	1 (0.0)
N stage, No. (%)			
N0+N0 (i+)	791 (43.3)	821 (45.0)	1,612 (44.2)
N1+N1mi	724 (39.7)	703 (38.5)	1,427 (39.1)
N2	212 (11.6)	210 (11.5)	422 (11.6)
N3	97 (5.3)	91 (5.0)	188 (5.2)
ER/PgR receptor status, No. (%)			
Positive	1,268 (69.5)	1,265 (69.3)	2,533 (69.4)
Negative	556 (30.5)	560 (30.7)	1,116 (30.6)
HER2 status, No. (%)			
Negative	1,510 (82.8)	1,519 (83.2)	3,029 (83.0)
Positive	314 (17.2)	306 (16.8)	620 (17.2)
Histologic grade, No. (%)			
Grade 1 (low)	170 (9.3)	160 (8.8)	330 (9.0)
Grade 2 (intermediate)	644 (35.3)	645 (35.3)	1,289 (35.3)
Grade 3 (high)	996 (54.6)	998 (54.7)	1,994 (54.6)
Missing/unknown	14 (0.8)	22 (1.2)	36 (1.0)
Adjuvant radiotherapy, No. (%)			
No	491 (26.9)	455 (24.9)	946 (25.9)
Yes	1,333 (73.1)	1,370 (75.1)	2,703 (74.1)

(continued on following page)

TABLE 1. Baseline Patient and Tumor Characteristics (continued)

Characteristic	Metformin (n = 1,824)	Placebo (n = 1,825)	Total (N = 3,649)
Adjuvant chemotherapy, No. (%)			
No	196 (10.7)	195 (10.7)	391 (10.7)
Yes	1,628 (89.3)	1,630 (17.3)	3,258 (89.3)
Adjuvant hormone therapy, No. (%)			
None	700 (38.4)	705 (38.6)	1,405 (38.5)
Tamoxifen	607 (33.3)	624 (34.2)	1,231 (33.7)
Aromatase inhibitor	517 (28.3)	496 (27.2)	1,013 (27.8)
Adjuvant trastuzumab, No. (%)			
No	1,510 (82.8)	1,510 (82.7)	3,020 (82.8)
Yes	314 (17.2)	315 (17.3)	629 (17.2)

Abbreviations: ECOG, Eastern Cooperative Oncology group; ER, estrogen receptor; HER2, human epidermal growth factor receptor 2; N, nodal; PgR, progesterone receptor; SD, standard deviation; T, tumor.

^aPremenopausal: <6 months since last menses and no bilateral oophorectomy OR does not meet postmenopausal criteria and younger than 50 years.

^bPostmenopausal: prior bilateral oophorectomy or >12 months since last menses OR does not meet premenopausal criteria and older than 50 years.

the absence of an effect of metformin on invasive disease-free survival (IDFS) or overall survival in hormone receptor-positive or hormone receptor-negative breast cancer.²¹ Exploratory analyses suggested beneficial effects in human epidermal growth factor receptor 2 (HER2)-positive breast cancer, notably in those with at least one C allele of the rs11212617 single-nucleotide polymorphism (SNP), a SNP associated with enhanced glycemic response and higher metformin blood levels in patients with diabetes.^{22,23}

METHODS

Detailed methods have been reported previously.²¹ Briefly, this investigator-initiated phase III randomized trial recruited patients (2010–2013) age 18–74 years, receiving standard therapy for a high-risk T1–3, N0–3, M0 breast cancer (excluding T1abNo) and diagnosed during the previous year. Exclusions included fasting glucose >7.0 mmol/L, history of diabetes or lactic acidosis, current use of diabetes medication, previous recurrence or other invasive cancer, intake of ≥3 alcoholic drinks daily, or marked organ dysfunction. Random assignment (to metformin 850 mg orally twice a day or placebo orally twice a day for 5 years) was stratified for (1) estrogen receptor and/or progesterone receptor positive (≥1%) versus both negative (<1%), (2) BMI ≤30 kg/m² versus >30 kg/m², (3) HER2 positive versus negative, and (4) any versus no adjuvant chemotherapy. In 2016, after a second interim analysis, futility was declared in hormone receptor-negative patients; study drug was discontinued in this group after a median exposure of 36.7 months, but blinded follow-up and event ascertainment continued.

Our current focus was a secondary analysis of new primary invasive cancers in the contralateral breast (a protocol-specified secondary outcome) and exploratory analyses of

new cancers elsewhere and overall (ipsilateral breast events were considered recurrences) that occurred as first IDFS events, with an exploration of whether effects of metformin differed by baseline BMI, insulin, glucose, homeostasis model assessment (HOMA, reflecting insulin resistance, calculated as glucose [mg/dL] × insulin [pmol/L]/22.5),²⁴ leptin and highly sensitive C-reactive protein (hsCRP) as well as smoking, rs11212617 SNP status, and receipt of adjuvant hormone therapy. Time to new cancer was analyzed by the competing risks method²⁵; two-sided likelihood ratio tests adjusting for age, BMI, smoking (ever v never smoked), and alcohol (any v no intake during the past year), with other IDFS events as competing risk and censoring at the last follow-up without an IDFS event, were performed. MA.32 was approved by the Adult Central Institutional Review Board (National Institutes of Health) and institutional review boards of the participating institutions. Patients provided written informed consent to participate.

RESULTS

We randomly assigned 3,649 patients; all were included in intent-to-treat analyses. Baseline subject, tumor, and treatment characteristics were well balanced between study arms (Table 1). The median follow-up and drug exposure were 95.9 (range, 0–121) months and 58.8 months, respectively.

There were 184 new primary cancers reported as a first IDFS event, 102 on the metformin arm and 82 on the placebo arm (HR, 1.25; 95% CI, 0.94 to 1.68; *P* = .13; Fig 1). Forty-eight were contralateral invasive breast cancers (27 metformin arm, 21 placebo arm), HR, 1.29 (95% CI, 0.72 to 2.27; *P* = .40), and 136 were nonbreast invasive cancers (75 metformin, 61 placebo), HR 1.24 (95% CI, 0.88 to 1.74; *P* = .21). The most common nonbreast cancers were gynecological (n = 31), lung (n = 26), hematological (n = 23), GI




FIG 1. Cumulative incidence curves with respect to metformin and placebo for (A) any invasive cancer, (B) invasive breast cancer, (C) nonbreast invasive cancer, and (D) obesity-associated cancer. ^aObesity-associated cancers include invasive breast, colorectal, pancreatic, gastric cardia, ovary, uterus, kidney, thyroid, and myeloma. HR, hazard ratio.

(*n* = 17), and melanoma (*n* = 13; Appendix Table A1, online only). The effect of metformin versus placebo on the risk of developing a new invasive cancer did not differ by hormone receptor status (positive HR, 1.19 [95% CI, 0.83 to 1.69] *v* negative HR, 1.44 [95% CI, 0.86 to 2.41]) or HER2 status (positive HR, 0.83 [95% CI, 0.39 to 1.81] *v* negative HR, 1.36 [95% CI, 0.99 to 1.86]) of the primary tumor (interaction *P* values .53 and .25, respectively) or by receipt of adjuvant hormonal therapy (tamoxifen HR, 1.23 [95% CI, 0.73 to 2.08]; aromatase inhibitor HR, 1.34 [95% CI, 0.80 to 2.24]; none HR, 1.21 [95% CI, 0.75 to 1.95; interaction *P* value .96). Metformin effects were similar for obesity-associated cancers

(invasive breast, colorectal, pancreatic, gastric cardia, ovary, uterus, kidney, thyroid, and myeloma)²⁶ and other cancers (HR, 1.18; 95% CI, 0.81-1.74; *P* = .39 and HR, 1.37; 95% CI, 0.88 to 2.13; *P* = .16, respectively) and for tobacco-associated (lung, head and neck, kidney, and bladder) and other cancers (HR, 1.77; 95% CI, 0.87 to 3.60; *P* = .11 and HR, 1.18; 95% CI, 0.85 to 1.62; *P* = .32). The effects of metformin did not differ by baseline levels of BMI (<25, 25-30, >30); quartiles of insulin, HOMA, leptin, and hsCRP; smoking (never, ever); and rs11212617 SNP status (any C *v* AA genotype; Table 2). Ten patients developed contralateral ductal carcinoma in situ (eight metformin, two placebo); when these were included as

TABLE 2. Effect of Metformin Versus Placebo on Cancer Risk in Relation to BMI, Metabolic Factors, Smoking, and rs11212617 SNP Status, With (unadjusted) HRs and Interaction *P* Values

Characteristic	Any Invasive Cancer (n = 184 events)	Invasive Breast Cancer (n = 48 events)	Invasive Nonbreast Cancer (n = 136 events)	Obesity-Associated Cancer ^a (n = 104 events)
	Met v Placebo, HR (95% CI)	Met v Placebo, HR (95% CI)	Met v Placebo, HR (95% CI)	Met v Placebo, HR (95% CI)
BMI				
<25 kg/m ² (n = 1,174)	1.17 (0.67 to 2.05)	2.06 (0.52 to 8.14)	1.03 (0.56 to 1.91)	1.13 (0.50 to 2.55)
25-30 kg/m ² (n = 1,206)	1.01 (0.60 to 1.69)	1.17 (0.39 to 3.47)	0.97 (0.54 to 1.73)	1.30 (0.65 to 2.62)
>30 kg/m ² (n = 1,269)	1.57 (1.00 to 2.49)	1.17 (0.54 to 2.52)	1.83 (1.03 to 3.25)	1.13 (0.65 to 2.00)
Intx <i>P</i>	.43	.76	.24	.94
Insulin				
Q1 (0 to <41, n = 864)	1.17 (0.63 to 2.16)	1.42 (0.31 to 6.52)	1.12 (0.57 to 2.19)	0.91 (0.38 to 2.19)
Q2 (41 to <60, n = 843)	0.98 (0.54 to 1.77)	1.13 (0.38 to 3.36)	0.92 (0.46 to 1.86)	0.98 (0.41 to 2.34)
Q3 (69 to <95, n = 886)	1.77 (0.86 to 3.64)	1.67 (0.48 to 5.84)	1.80 (0.75 to 4.34)	2.24 (0.84 to 5.94)
Q4 (95+, n = 886)	1.43 (0.80 to 2.57)	1.34 (0.47 to 3.82)	1.46 (0.72 to 2.96)	1.13 (0.57 to 2.26)
Intx <i>P</i>	.60	.98	.62	.51
HOMA				
Q1 (0 to <1.26, n = 634)	1.12 (0.55 to 2.26)	1.17 (0.23 to 5.80)	1.10 (0.50 to 2.40)	0.92 (0.35 to 2.47)
Q2 (1.26 to <1.91, n = 632)	0.68 (0.34 to 1.37)	1.00 (0.20 to 4.90)	0.62 (0.28 to 1.37)	0.62 (0.21 to 1.89)
Q3 (1.91 to <3.03, n = 640)	1.28 (0.62 to 2.64)	0.82 (0.25 to 2.69)	1.67 (0.66 to 4.26)	1.08 (0.46 to 2.54)
Q4 (3.03+, n = 635)	1.53 (0.79 to 2.96)	1.84 (0.47 to 7.32)	1.42 (0.67 to 3.03)	1.09 (0.49 to 2.42)
Intx <i>P</i>	.39	.85	.34	.86
Leptin				
Q1 (0 to <6.38, n = 873)	1.09 (0.59 to 2.02)	1.11 (0.22 to 5.63)	1.09 (0.56 to 2.11)	1.01 (0.39 to 2.63)
Q2 (6.38 to <12.64, n = 881)	1.18 (0.62 to 2.23)	2.21 (0.58 to 8.43)	0.95 (0.45 to 2.00)	1.12 (0.50 to 2.51)
Q3 (12.64 to <22.03, n = 870)	1.33 (0.75 to 2.37)	1.72 (0.58 to 5.13)	1.20 (0.61 to 2.36)	1.09 (0.52 to 2.29)
Q4 (22.03+, n = 871)	1.53 (0.87 to 2.70)	0.80 (0.30 to 2.15)	2.11 (1.03 to 4.35)	1.36 (0.66 to 2.80)
Intx <i>P</i>	.86	.61	.41	.96
hs-CRP				
Q1 (0 to < 0.5, n = 777)	0.93 (0.46 to 1.88)	4.17 (0.45 to 38.9)	0.73 (0.34 to 1.57)	1.58 (0.56 to 4.49)
Q2 (0.5 to <1.3, n = 918)	0.94 (0.48 to 1.82)	0.40 (0.07 to 2.05)	1.15 (0.55 to 2.42)	0.75 (0.32 to 1.78)
Q3 (1.3 to <3.2, n = 902)	1.32 (0.78 to 2.22)	2.26 (0.79 to 6.50)	1.07 (0.58 to 1.97)	1.26 (0.61 to 2.61)
Q4 (3.2+, n = 898)	1.90 (1.08 to 3.37)	0.99 (0.40 to 2.48)	2.78 (1.29 to 6.00)	1.25 (0.64 to 2.46)
Intx <i>P</i>	.31	.16	.07	.68
Smoking history				
Never-smoker	1.36 (0.89 to 2.07)	1.51 (0.67 to 3.39)	1.30 (0.80 to 2.13)	1.33 (0.79 to 2.23)
Ever smoker	1.19 (0.79 to 1.78)	1.08 (0.50 to 2.48)	1.21 (0.76 to 1.94)	1.02 (0.57 to 1.83)
Intx <i>P</i>	.65	.59	.84	.51
rs11212617				
Any C	1.22 (0.85 to 1.76)	1.34 (0.64 to 2.83)	1.17 (0.78 to 1.80)	1.23 (0.75 to 2.02)
AA	1.53 (0.86 to 2.74)	1.17 (0.43 to 3.24)	1.71 (0.84 to 3.51)	1.10 (0.53 to 2.27)
Intx <i>P</i>	.52	.84	.37	.80

Abbreviations: HOMA, homeostasis model assessment; HR, hazard ratio; hs-CRP, highly sensitive C-reactive protein; Intx *P*, interaction *P* value; Met, metformin; Q, quartile; SNP, single-nucleotide polymorphism.

^aObesity-associated cancers were those categorized as having sufficient evidence of a link between body fatness and increased cancer risk by Lauby-Secretan et al.²⁶ These included GI (colorectal, gastric cardia, pancreatic), gynecologic (uterus, ovarian), and postmenopausal breast, kidney, thyroid, and myeloma.

cancers, the HR for metformin versus placebo for any breast cancer was 1.35 (95% CI, 0.77 to 2.37; $P = .30$) and for any cancer was 1.28 (95% CI, 0.95 to 1.70; $P = .10$).

DISCUSSION

The MA.32 trial provided a unique opportunity to examine the effects of metformin on the risk of developing new primary invasive cancers in a population without diabetes. Metformin did not affect the risk of new contralateral breast cancer, any invasive cancer, cancers outside of the breast, obesity, or tobacco-associated cancers. Point estimates of risk were all above 1, making it unlikely that additional power would have identified a clinically important reduction in risk. Furthermore, metformin did not reduce cancer risk in subgroups defined by BMI, metabolic

factors (insulin, HOMA, leptin, hsCRP), smoking history, or rs11212617 SNP status, factors that could potentially be associated with metformin benefit.

It is unclear whether our results are generalizable to populations with diabetes or prediabetes. Because of its beneficial physiologic effect, it is possible that metformin could reduce cancer risk in populations experiencing hyperglycemia, hyperinsulinemia, or inflammation. However, results of the Diabetes Prevention Program¹⁸ and critical appraisals of the observational studies in populations with diabetes¹⁻¹⁷ discussed above raise concern about metformin benefit.

These findings do not support the use of metformin to reduce risk of new primary cancers in patients without diabetes who have high-risk breast cancer.

AFFILIATIONS

¹Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, and Department of Medicine, University of Toronto, Toronto, ON, Canada

²Canadian Cancer Trials Group, Queen's University, Kingston, ON, Canada

³University of British Columbia, BC Cancer Agency, Vancouver, BC, Canada

⁴McMaster University, Juravinski Cancer Centre, Hamilton, ON, Canada

⁵Applied Statistician, Markham, ON, Canada

⁶CHU de Québec-Université Laval, Québec, QC, Canada

⁷Dana-Farber Cancer Institute, Boston, MA

⁸Herbert Irving Comprehensive Cancer Center, Columbia University Medical Center, New York, NY

⁹Vanderbilt University, Nashville, TN

¹⁰Mayo Clinic, Rochester, MN

¹¹ICR-CTS, Institute of Cancer Research (UK), London, United Kingdom

¹²NRG Oncology and University of Pittsburgh Medical Center, Pittsburgh, PA

¹³IBCSG and Department of Oncology, Bern University Hospital, University of Bern, Berne, Switzerland

¹⁴Baylor College of Medicine, Houston, TX

¹⁵Cancer Research UK Clinical Trials Unit (CRCTU), Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK

¹⁶Princess Margaret Cancer Centre, University Health Network, Toronto, ON, Canada

¹⁷Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada

CORRESPONDING AUTHOR

Pamela J. Goodwin, MD, Mount Sinai Hospital, 1085-600 University Ave, Toronto, ON M5G 1X5, Canada; e-mail: pamela.goodwin@sinahealth.ca

DISCLAIMER

The study sponsors have no role in the design of the study; the collection, analysis, and interpretation of the data; the writing of the manuscript; and the decision to submit the manuscript for publication.

SUPPORT

Supported by the Canadian Cancer Society Research Institute (No. 021039); National Cancer Institute (US; Nos. CA077202, CA180868,

CA180822); The Breast Cancer Research Foundation (New York); Canadian Breast Cancer Foundation—Ontario Region, Ontario Institute for Cancer Research (No. 10NOV-467); Apotex Canada (in kind donation of placebo and metformin); and Hold'em for Life Charity.

CLINICAL TRIAL INFORMATION

[NCT01101438](https://clinicaltrials.gov/ct2/show/study/NCT01101438)

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.23.00296>.

DATA SHARING STATEMENT

The primary efficacy analysis will be available from the Canadian Cancer trials Group (Kingston, ON). The associated data have been uploaded to the NCI data archive website: <http://nctn-data-archive.nci.nih.gov/view-trials> and will be searchable via ClinicalTrials.gov identifier ([NCT1101438](https://clinicaltrials.gov/ct2/show/study/NCT1101438)). Further information regarding that analysis and the data analyzed in this substudy can be obtained from the corresponding author.

AUTHOR CONTRIBUTIONS

Conception and design: Pamela J. Goodwin, Bingshu E. Chen, Karen A. Gelmon, Jennifer A. Ligibel, Dawn L. Hershman, Ingrid A. Mayer, Alastair M. Thompson, Vuk Stambolic, Wendy R. Parulekar

Provision of study materials or patients: Pamela J. Goodwin, Karen A. Gelmon, Julie Lemieux, Jennifer A. Ligibel, Dawn L. Hershman, Ingrid A. Mayer, Timothy J. Hobday, Priya Rastogi, Manuela Rabaglio-Poretti, Alastair M. Thompson, Daniel W. Rea, Wendy R. Parulekar

Collection and assembly of data: All authors

Data analysis and interpretation: Pamela J. Goodwin, Bingshu E. Chen, Karen A. Gelmon, Timothy J. Whelan, Marguerite Ennis, Julie Lemieux, Dawn L. Hershman, Ingrid A. Mayer, Timothy J. Hobday, Judith M. Bliss, Priya Rastogi, Alastair M. Thompson, Daniel W. Rea, Paul M. Stos, Vuk Stambolic, Wendy R. Parulekar

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

REFERENCES

1. Zhang K, Bai P, Dai H, et al: Metformin and risk of cancer among patients with type 2 diabetes mellitus: A systematic review and meta-analysis. *Prim Care Diabetes* 15:52-58, 2021
2. Mekuria AN, Ayele Y, Tola A, et al: Monotherapy with metformin versus sulfonylureas and risk of cancer in type 2 diabetic patients: A systematic review and meta-analysis. *J Diabetes Res* 2019:1-8, 2019
3. Tang GH, Satkunam M, Pond G, et al: Association of metformin with breast cancer incidence and mortality in patients with type II diabetes: A GRADE-assessed systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 27:627-635, 2018
4. Hevroni G, Skwiersky S, Zhyvotovska A, et al: Metformin use and the risk of gastrointestinal malignancies in diabetic populations: A meta-analysis. *Int J Clin Endocrinol Metab* 6:035-041, 2020
5. Ng C-AW, Jiang AA, Toh EMS, et al: Metformin and colorectal cancer: A systematic review, meta-analysis and meta-regression. *Int J Colorectal Dis* 35:1501-1512, 2020
6. Yao L, Liu M, Huang Y, et al: Metformin use and lung cancer risk in diabetic patients: A systematic review and meta-analysis. *Dis Markers* 2019:6230162, 2019
7. Shi J, Liu B, Wang H, et al: Association of metformin use with ovarian cancer incidence and prognosis: A systematic review and meta-analysis. *Int J Gynecol Cancer* 29:140-146, 2019
8. Tian J, Liang Y, Qu P: Antidiabetic medications and the risk of endometrial cancer in patients. *Gynecol Obstet Invest* 84:455-462, 2019
9. Wen Q, Zhao Z, Wen J, et al: The association between metformin therapy and risk of gynecological cancer in patients: Two meta-analyses. *Eur J Obstet Gynecol Reprod Biol* 237:33-41, 2019
10. Cui H, Wang Y, Yang S, et al: Antidiabetic medications and the risk of prostate cancer in patients with diabetes mellitus: A systematic review and meta-analysis. *Pharmacol Res* 177:106094, 2022
11. Feng Z, Zhou X, Liu N, et al: Metformin use and prostate cancer risk. A meta-analysis of cohort studies. *Medicine* 98:e14955, 2019
12. Xiao K, Liu F, Liu J, et al: The effect of metformin on lung cancer risk and survival in patients with type 2 diabetes mellitus: A meta-analysis. *J Clin Pharm Ther* 45:783-792, 2020
13. Hu J, Chen J, Cui Y, et al: Association of metformin intake with bladder cancer risk and oncologic outcomes in type 2 diabetes mellitus patients. A systematic review and meta-analysis. *Medicine* 97:e11596, 2018
14. Chang MS, Hartman RI, Xue J, et al: Risk of skin cancer associated with metformin use: A meta-analysis of randomized controlled trials and observational studies. *Cancer Prev Res* 14:77-84, 2021
15. Suissa S, Azoulay L: Metformin and the risk of cancer. *Diabetes Care* 35:2665-2673, 2012
16. Li X, Celotto S, Pizzol D, et al: Metformin and health outcomes: An umbrella review of systematic reviews with meta-analyses. *Eur J Clin Invest* 51:e13536, 2021
17. Farmer RE, Ford D, Forbes HJ, et al: Metformin and cancer in type 2 diabetes: A systematic review and comprehensive bias evaluation. *Int J Epidemiol* 46:745-744, 2017
18. Lee CG, Hechman-Stoddard B, Dabela D, et al: Effect of metformin and lifestyle interventions on mortality in the Diabetes Prevention Program and Diabetes Prevention Program Outcomes Study. *Diabetes Care* 44:2775-2782, 2021
19. Saini N, Yang X: Metformin as an anti-cancer agent: Actions and mechanisms targeting cancer stem cells. *Acta Biochim Biophys Sin (Shanghai)* 50:133-143, 2018
20. Dowling RJ, Niraula S, Stambolic V, et al: Metformin in cancer: Translational challenges. *J Mol Endocrinol* 48:R31-R43, 2012
21. Goodwin PJ, Chen BE, Gelmon KA, et al: Effect of metformin vs placebo on invasive disease-free survival in patients with breast cancer. The MA.32 randomized clinical trial. *JAMA* 327:1963-1973, 2022
22. van Leeuwen N, Nijpels G, Becker ML, et al: A gene variant near ATM is significantly associated with metformin treatment response in type 2 diabetes: A replication and meta-analysis of five cohorts. *Diabetologia* 55:1971-1977, 2012
23. Luizon MR, Eckalbar WL, Wang Y, et al: Genomic characterization of metformin hepatic response. *PLoS Genet* 12:e1006449, 2016
24. Bonora E, Targher G, Alberiche M, et al: Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: Studies in subjects with various degrees of glucose tolerance and insulin sensitivity. *Diabetes Care* 23:57-63, 2000
25. Fine JP, Gray RJ: A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc* 94:496-509, 1999
26. Lauby-Secretan B, Scoccianti C, Loomis D, et al: Body fatness and cancer—viewpoint of the IARC working group. *N Engl J Med* 375:794-798, 2016

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**Effect of Metformin Versus Placebo on New Primary Cancers in Canadian Cancer Trials Group MA.32: A Secondary Analysis of a Phase III Randomized Double-Blind Trial in Early Breast Cancer**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Pamela J. Goodwin

Research Funding: Epic Sciences (Inst)

Karen A. Gelmon

Honoraria: AstraZeneca, Merck Sharp & Dohme, Seagen, Novartis Canada Pharmaceuticals Inc, Pfizer, Lilly, Gilead Sciences

Consulting or Advisory Role: Pfizer, Novartis, Astrazeneca, Merck, Lilly, Roche, Mylan, Ayala Pharmaceuticals, Gilead Sciences

Research Funding: Pfizer (Inst), Bristol Myers Squibb (Inst), AstraZeneca (Inst)

Expert Testimony: Genentech

Timothy J. Whelan

Research Funding: Exact Sciences (Inst)

Julie Lemieux

Consulting or Advisory Role: Novartis, Lilly, AstraZeneca, Gilead Sciences, Merck

Research Funding: Imagia

Patents, Royalties, Other Intellectual Property: Patent related to health

Dawn L. Hershman

Consulting or Advisory Role: AIM Specialty Health

Ingrid A. Mayer

Employment: AstraZeneca

Leadership: AstraZeneca

Stock and Other Ownership Interests: AstraZeneca

Consulting or Advisory Role: AstraZeneca, Lilly, Genentech, MacroGenics, Pfizer, Seagen, Sanofi

Research Funding: Novartis (Inst), Pfizer (Inst), Genentech (Inst)

Timothy J. Hobday

Research Funding: Novartis (Inst)

Judith M. Bliss

Research Funding: AstraZeneca (Inst), Merck Sharp & Dohme (Inst), Puma Biotechnology (Inst), Pfizer (Inst), Roche (Inst), GlaxoSmithKline/Novartis (Inst), Lilly (Inst), Janssen-Cilag (Inst), Clovis Oncology (Inst)
Travel, Accommodations, Expenses: Pfizer

Priya Rastogi

Travel, Accommodations, Expenses: Genentech/Roche, Lilly, AstraZeneca

Alastair M. Thompson

Employment: Lilly

Stock and Other Ownership Interests: Lilly

Travel, Accommodations, Expenses: Endomagnetics

Daniel W. Rea

Honoraria: Roche, Novartis, Pfizer, Lilly, AstraZeneca

Research Funding: Celgene (Inst), Roche (Inst), bioTheranostics (Inst)

Travel, Accommodations, Expenses: Pfizer, Daiichi Sankyo, eisai, Novartis

No other potential conflicts of interest were reported.

APPENDIX

TABLE A1. New Invasive Cancers Reported as a First Invasive Disease-Free Survival Event

Cancer	Metformin, No.	Placebo, No.	Total, No.
Any invasive cancer	102	82	184
Breast	27	21	48
Brain	2	0	2
GI			
Pancreatic	2	1	3
Gastric cardia	3	0	3
Colorectal	6	5	11
Genitourinary			
Bladder	1	1	2
Kidney	1	1	2
Gynecological			
Fallopian tube	1	0	1
Ovarian	5	4	9
Uterus	8	9	17
Vaginal	1	0	1
Cervical	2	1	3
Head and neck	3	0	3
Hematological			
Leukemia	9	3	12
Lymphoma	1	6	7
Myeloma	1	3	4
Lung	16	10	26
Lymphoid disorder	1	0	1
Melanoma	5	8	13
Neuroendocrine	1	2	3
Peritoneum	1	0	1
Salivary gland	1	3	4
Thyroid	3	4	7
Unknown primary (adenocarcinoma)	1	0	1