

A note on colour-bias perfect matchings in hypergraphs

Balogh, Jozsef; Treglown, Andrew; Zarate Gueren, Camila

License:
Creative Commons: Attribution (CC BY)

Document Version
Peer reviewed version

Citation for published version (Harvard):
Balogh, J, Treglown, A & Zarate Gueren, C 2024, 'A note on colour-bias perfect matchings in hypergraphs', *SIAM Journal on Discrete Mathematics*.

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

A NOTE ON COLOUR-BIAS PERFECT MATCHINGS IN HYPERGRAPHS

JÓZSEF BALOGH, ANDREW TREGLOWN AND CAMILA ZÁRATE-GUERÉN

ABSTRACT. A result of Balogh, Csaba, Jing and Pluhár yields the minimum degree threshold that ensures a 2-coloured graph contains a perfect matching of significant colour-bias (i.e., a perfect matching that contains significantly more than half of its edges in one colour). In this note we prove an analogous result for perfect matchings in k -uniform hypergraphs. More precisely, for each $2 \leq \ell < k$ and $r \geq 2$ we determine the minimum ℓ -degree threshold for forcing a perfect matching of significant colour-bias in an r -coloured k -uniform hypergraph.

1. INTRODUCTION

A *perfect matching* in a hypergraph H is a collection of vertex-disjoint edges of H which covers the vertex set $V(H)$ of H . In recent decades there has been significant interest in the problem of establishing *minimum degree* conditions that force a perfect matching in a k -uniform hypergraph. More precisely, given a k -uniform hypergraph H and an ℓ -element vertex set $S \subseteq V(H)$ (where $\ell \in [k-1]$) we define $d_H(S)$ to be the number of edges containing S . The *minimum ℓ -degree* $\delta_\ell(H)$ of H is the minimum of $d_H(S)$ over all ℓ -element sets of vertices in H . We refer to $\delta_1(H)$ as the *minimum vertex degree* of H and $\delta_{k-1}(H)$ as the *minimum codegree* of H .

Suppose that $\ell, k, n \in \mathbb{N}$ such that $\ell \leq k-1$ and k divides n . Let $m_\ell(k, n)$ denote the smallest integer m such that every k -uniform hypergraph H on n vertices with $\delta_\ell(H) \geq m$ contains a perfect matching.

A simple consequence of Dirac's theorem is that $m_1(2, n) = n/2$ for all even $n \in \mathbb{N}$. Improving earlier asymptotically exact bounds given in [12, 17], Rödl, Ruciński and Szemerédi [18] determined the minimum codegree threshold for perfect matchings in k -uniform hypergraphs. That is, they showed that if $n \in \mathbb{N}$ is sufficiently large, then $m_{k-1}(k, n) = n/2 - k + C$, where $C \in \{3/2, 2, 5/2, 3\}$ depends on the values of n and k .

The value of $m_\ell(k, n)$ is known for various pairs (k, ℓ) when n is sufficiently large. For example, after an earlier asymptotic result of Pikhurko [15], Treglown and Zhao [19] determined the value of $m_\ell(k, n)$ for $\ell \geq k/2$ and n sufficiently large. However, the minimum vertex degree case of the problem is wide open in general, and the only cases where the asymptotic or exact value of $m_1(k, n)$ is known is when $k = 2, 3, 4, 5$. See, e.g., [16, 21] for discussions on further results in the area.

Given any $1 \leq \ell < k$ it is known that

$$(1) \quad m_\ell(k, n) \geq \max \left\{ \frac{1}{2} - o(1), 1 - \left(\frac{k-1}{k} \right)^{k-\ell} - o(1) \right\} \binom{n}{k-\ell}.$$

See, e.g., the introduction of [20] for the two families of hypergraphs that demonstrate (1). It is widely believed that the inequality in (1) is asymptotically sharp for all choices of k, ℓ , see [11, 13].

JB: University of Illinois Urbana-Champaign, 1409 W. Green Street, Urbana IL 61801, United States. Research supported in part by NSF grants DMS-1764123 and RTG DMS-1937241, FRG DMS-2152488, the Arnold O. Beckman Research Award (UIUC Campus Research Board RB 24012), the Langan Scholar Fund (UIUC).

AT: University of Birmingham, United Kingdom, a.c.treglown@bham.ac.uk. Research supported by EPSRC grant EP/V002279/1.

CZG: University of Birmingham, United Kingdom, ciz230@student.bham.ac.uk. Research supported by EPSRC.

Moreover, Treglown and Zhao [20] gave a conjecture on the exact value of $m_\ell(k, n)$ for sufficiently large $n \in k\mathbb{N}$.

The aim of this paper is to study the *colour-bias* version of this problem. The topic of colour-bias structures in graphs was first raised by Erdős in the 1960s (see [5, 6]). Sparked by work of Balogh, Csaba, Jing and Pluhár [1], there has been renewed interest in the topic, particularly in establishing minimum degree conditions that force a colour-bias copy of a graph F . More precisely, if a graph G contains a copy of F , then however the edges of G are 2-coloured, one can clearly ensure that G contains a copy of F with at least $e(F)/2$ edges of the same colour. The question then is how large does the minimum degree $\delta(G)$ of G need to be to guarantee that G contains a copy of F with significantly more than $e(F)/2$ edges of the same colour, no matter how one 2-colours the edges of G ? The following result resolves this problem in the case when F is a Hamilton cycle.

Theorem 1.1 (Balogh, Csaba, Jing and Pluhár [1]). *Let $0 < c < 1/4$ and $n \in \mathbb{N}$ be sufficiently large. If G is an n -vertex graph with*

$$\delta(G) \geq (3/4 + c)n,$$

then given any 2-colouring of $E(G)$ there is a Hamilton cycle in G with at least $n/2 + cn/32$ edges of the same colour. Moreover, if $n \in 4\mathbb{N}$, there is an n -vertex graph G' with $\delta(G') = 3n/4$ and a 2-colouring of $E(G')$ for which every Hamilton cycle in G' has precisely $n/2$ edges in each colour.

Note that Theorem 1.1 shows that the minimum degree threshold for forcing a *colour-bias* Hamilton cycle in a graph is significantly higher than the threshold for just forcing a Hamilton cycle. Indeed, Dirac's theorem tells us that any n -vertex graph G with $\delta(G) \geq n/2$ contains a Hamilton cycle.

Since a Hamilton cycle on an even number of vertices is the union of two perfect matchings, Theorem 1.1 implies the following result.

Theorem 1.2 (Balogh, Csaba, Jing and Pluhár [1]). *Let $0 < c < 1/4$ and $n \in 2\mathbb{N}$ be sufficiently large. If G is an n -vertex graph with*

$$\delta(G) \geq (3/4 + c)n,$$

then given any 2-colouring of $E(G)$ there is a perfect matching in G with at least $n/4 + cn/64$ edges of the same colour. Moreover, if $n \in 4\mathbb{N}$, there is an n -vertex graph G' with $\delta(G') = 3n/4$ and a 2-colouring of $E(G')$ for which every perfect matching in G' has precisely $n/4$ edges in each colour.

Let $n \in 4\mathbb{N}$. We define the graph G' in Theorem 1.2 as follows: $V(G')$ consists of the disjoint union of two vertex classes A and B of sizes $n/4$ and $3n/4$, respectively; $E(G')$ contains all possible red edges whose endpoints are both in B and all possible blue edges with one endpoint in A and one endpoint in B . Thus, $\delta(G') = 3n/4$ and every perfect matching in G' has precisely $n/4$ edges in each colour.

Since [1] appeared, a number of analogues of Theorem 1.1 have been established for other types of spanning structures. Given graphs G and F , an F -factor in G is a collection of vertex-disjoint copies of F in G that together cover $V(G)$. In [2], the minimum degree threshold for forcing a colour-bias K_r -factor was determined.¹ More recently, this result was extended to F -factors for every fixed graph F ; see [4]. For $k \geq 2$, the minimum degree threshold for forcing a colour-bias k th power of a Hamilton cycle in a graph was established in [3].

Other variants of the problem have also been studied. In [7, 10] an r -colour version of Theorem 1.1 was proven: in this setting now one r -colours $E(G)$ and seeks a Hamilton cycle with significantly more than n/r edges of the same colour. Colour-bias problems have also been considered for random graphs [9]. Recently, Mansilla Brito [14] gave a minimum codegree result for forcing a colour-bias

¹Recall K_r denotes the complete graph on r vertices.

copy of a tight Hamilton cycle in a 3-uniform hypergraph. We remark that all of these colour-bias results can be phrased in the equivalent language of *discrepancy*; see, e.g., [1, 2, 3, 4, 10].

Our main result determines the minimum ℓ -degree threshold for forcing a colour-bias perfect matching in a k -uniform hypergraph for all $\ell \geq 2$ and $k \geq 3$. To state our result we need the following definitions. Given integers $1 \leq \ell < k$, let $\mathcal{C}_{k,\ell}$ be the set of all $c > 0$ such that $m_\ell(k, n) \leq c \binom{n}{k-\ell}$ for all sufficiently large $n \in k\mathbb{N}$. Set $c_{k,\ell}$ to be the infimum of $\mathcal{C}_{k,\ell}$. In particular, note that the general conjecture on the asymptotic value of $m_\ell(k, n)$ equivalently states that

$$c_{k,\ell} = \max \left\{ \frac{1}{2}, 1 - \left(\frac{k-1}{k} \right)^{k-\ell} \right\}.$$

Theorem 1.3. *Let $k, \ell, r \in \mathbb{N}$ where $2 \leq \ell < k$ and $r \geq 2$. Given any $\eta > 0$ where $c_{k,\ell} + \eta < 1$, there exists an $n_0 \in \mathbb{N}$ such that the following holds. Let H be a k -uniform hypergraph on $n \geq n_0$ vertices, where $n \in k\mathbb{N}$. If*

$$\delta_\ell(H) \geq (c_{k,\ell} + \eta) \binom{n}{k-\ell},$$

then given any r -colouring of $E(H)$ there is a perfect matching in H with at least $\frac{n}{rk} + \frac{\eta n}{8r(r-1)k^k(k^2+k)}$ edges of the same colour.

We remark that Theorem 1.3 holds even in the cases in which we do not know the value of $c_{k,\ell}$. By definition of $c_{k,\ell}$, the minimum ℓ -degree condition in Theorem 1.3 is essentially best possible. Indeed, for $c < c_{k,\ell}$, a minimum ℓ -degree condition of $\delta_\ell(H) \geq c \binom{n}{k-\ell}$ does not even guarantee a perfect matching, let alone one of significant colour-bias. So in this sense the colour-bias and ‘standard’ versions of the problem are aligned when $\ell \geq 2$.

In contrast, the same phenomenon does not occur for the minimum vertex degree version of the problem. Indeed, Theorem 1.2 tells us that the minimum degree threshold for a colour-bias perfect matching in a *graph* is different to the minimum degree threshold for a perfect matching in a graph. Furthermore, in Section 4 we describe a similar phenomenon in the 3-uniform hypergraph setting.

Remark. Whilst finalising a manuscript that gave the proof of Theorem 1.3 in the case when $\ell = k - 1$ and $r = 2$, we learnt of simultaneous and independent work of Gishboliner, Glock and Sgueglia [8]. In [8] they determine the *minimum codegree threshold* for forcing a tight Hamilton cycle of significant colour-bias in an r -coloured k -uniform hypergraph (where $r \geq 2$ and $k \geq 3$). As an immediate consequence of their result they also establish the corresponding *minimum codegree threshold* for perfect matchings.

We therefore decided to seek a generalisation of our minimum codegree result to other degree conditions, i.e., Theorem 1.3. In doing so, we found an argument much cleaner than our original approach.

Notation. Let H be a hypergraph. The *neighbourhood* $N_H(X)$ of a set $X \subseteq V(H)$ is the family of sets $S \subseteq V(H) \setminus X$ such that $S \cup X \in E(H)$. If $X = \{x\}$ we define $N_H(x) := N_H(X)$. Given a vertex $x \in V(H)$ and set $Y \subseteq V(H)$ we sometimes write xY or Yx to denote $\{x\} \cup Y$. Given a colouring c of $E(H)$, we call an edge $e \in E(H)$ a C -edge if e is coloured C in c . Given a set $X \subseteq V(H)$, we write $H[X]$ for the *induced subhypergraph of H with vertex set X* . We define $H \setminus X := H[V(H) \setminus X]$.

Given a hypergraph F with an r -colouring $c : E(F) \rightarrow \{C_1, \dots, C_r\}$, its *colour profile* is (x_1, \dots, x_r) where x_i is the number of C_i -edges in F for each $i \in [r]$. Two colour profiles (x_1, \dots, x_r) , (y_1, \dots, y_r) are said to be *different with respect to the colour C_i* if $x_i \neq y_i$.

2. PRELIMINARIES AND USEFUL RESULTS

2.1. Proof overview and key definitions. Throughout this section, we will suppose that H is a k -uniform hypergraph on n vertices with an r -colouring $c : E(H) \rightarrow \{C_1, \dots, C_r\}$.

Our general strategy for the proof of Theorem 1.3 is as follows. Our aim is to find certain *gadgets* inside of H . A gadget is just a subhypergraph of H with some given structure. A gadget G is *good* if G contains two perfect matchings that have different colour profiles with respect to the r -colouring c .

For a certain well chosen $t \in \mathbb{N}$, we will prove that there are t vertex-disjoint good gadgets G_1, \dots, G_t in H and a $j \in [r]$ so that, for each good gadget G_i , the two perfect matchings M_i and M'_i in G_i have colour profiles that are different with respect to the colour C_j .

We will then be able to easily find a perfect matching in H of significant colour-bias. Indeed, removing the vertices of G_1, \dots, G_t from H will result in a k -uniform hypergraph H' that contains a perfect matching M . The flexibility of the good gadgets then allows us to extend M into a perfect matching in H with significant colour-bias, whatever the colour profile of M is.

We next state the definitions required to formally introduce the notion of a good gadget.

Definition 2.1. Let $u, v \in V(H)$ be distinct and $T \in N_H(u) \cap N_H(v)$. We say

- **uTv is S** if $c(T \cup \{u\}) = c(T \cup \{v\})$;
- **uTv is $C_i C_j$** if $c(T \cup \{u\}) = C_i$ and $c(T \cup \{v\}) = C_j$.

Let $C_i C_j(uv)$ denote the collection of sets $T \in N_H(u) \cap N_H(v)$ for which uTv is $C_i C_j$. Define $S(uv)$ analogously.

Note that $C_i C_j(uv) = C_j C_i(vu)$ for all distinct $u, v \in V(H)$.

Definition 2.2. Let $D > 0$ and let $u, v \in V(H)$ be distinct. We say that $N_H(u) \cap N_H(v)$ is

- **type S(D)** if $|S(uv)| \geq Dn^{k-2}$;
- **type $C_i C_j$ (D)** if $i \neq j$ and $|C_i C_j(uv)| \geq Dn^{k-2}$.

We remark that it may be the case that $N_H(u) \cap N_H(v)$ has more than one type.

Definition 2.3. Let $e = \{e_1, \dots, e_k\}$ and $f = \{f_1, \dots, f_k\}$ be two edges in H . A **$(k^2 + k, \mathbf{e}, \mathbf{f})$ -gadget** G is a subhypergraph of H on $k^2 + k$ vertices so that:

- $V(G)$ is the disjoint union of e, f and T_1, \dots, T_k where $T_i \in N_H(e_i) \cap N_H(f_i)$ for each $i \in [k]$;
- $e, f \in E(G)$;
- $e_i T_i, f_i T_i \in E(G)$ for all $i \in [k]$.

A $(k^2 + k, e, f)$ -gadget in which every $e_i T_i f_i$ is S will be called an **S- $(k^2 + k, \mathbf{e}, \mathbf{f})$ -gadget**.

A **$(3k, \mathbf{e}, \mathbf{f})$ -gadget** G is a subhypergraph of H on $3k$ vertices so that:

- $e_i = f_i$, for all $i \in \{3, \dots, k\}$;
- $V(G)$ is the disjoint union of e, f_1, f_2, T_1 and T_2 , where $T_i \in N_H(e_i) \cap N_H(f_i)$ for each $i \in [2]$;
- $e, f \in E(G)$;
- $e_1 T_1, f_1 T_1, e_2 T_2, f_2 T_2 \in E(G)$.

Given $t \in \{3k, k^2 + k\}$, we say that a (t, e, f) -gadget G is **good** if it contains two perfect matchings with different colour profiles (with respect to the r -colouring of G induced by the r -colouring c of H).

Note that e and f are vertex-disjoint in a $(k^2 + k, e, f)$ -gadget but intersect in $k - 2$ vertices in a $(3k, e, f)$ -gadget; see Figure 1.




FIGURE 1. On the left, a $(12, e, f)$ -gadget. On the right, a $(9, e, f)$ -gadget.

2.2. Tools for the proof of Theorem 1.3. The following well-known result allows one to deduce a lower bound on $\delta_\ell(H)$ given a lower bound on $\delta_{\ell'}(H)$, for any $\ell \leq \ell'$.

Proposition 2.4. *Let $1 \leq \ell \leq \ell' < k$ and H be a k -uniform hypergraph on n vertices. If $\delta_{\ell'}(H) \geq x \binom{n-\ell'}{k-\ell'}$ for some $0 \leq x \leq 1$, then $\delta_\ell(H) \geq x \binom{n-\ell}{k-\ell}$. \square*

The next result gives a sufficient condition for finding a good $(3k, e, f)$ -gadget in a k -uniform hypergraph of large minimum 2-degree.

Lemma 2.5. *Let $k \geq 3$ and $D := 3k$. Let H be a k -uniform hypergraph on n vertices with an r -colouring $c : E(H) \rightarrow \{C_1, \dots, C_r\}$. Suppose there exists $i \neq j \in [r]$ and distinct $v_1, v_2, v_3, v_4 \in V(H)$ such that $N_H(v_1) \cap N_H(v_2)$ and $N_H(v_3) \cap N_H(v_4)$ are both type $C_i C_j(D)$. If*

$$\delta_2(H) > \frac{1}{2} \binom{n}{k-2},$$

then there exists a good $(3k, e, f)$ -gadget in H , for some $e, f \in E(H)$.

Proof. By the minimum 2-degree condition, there exists a set $X \subseteq V(H)$ of size $k-2$ such that $A = X \cup \{v_1, v_3\}$ and $B = X \cup \{v_2, v_4\}$ are both in $E(H)$. We show that we can construct a $(3k, A, B)$ -gadget and afterwards we prove that it is good.

Given that $N_H(v_1) \cap N_H(v_2)$ is type $C_i C_j(D)$, there are at least $3kn^{k-2}$ sets $T_{1,2} \in N_H(v_1) \cap N_H(v_2)$ such that $c(v_1 T_{1,2}) = C_i$ and $c(v_2 T_{1,2}) = C_j$. As $|A \cup B| = k+2 < 3k$, we may choose such a set $T_{1,2}$ so that it is also vertex-disjoint from $A \cup B$. Similarly, there is a set $T_{3,4} \in N_H(v_3) \cap N_H(v_4)$ such that $c(v_3 T_{3,4}) = C_i$, $c(v_4 T_{3,4}) = C_j$ and $T_{3,4}$ is vertex-disjoint from A, B and $T_{1,2}$.

Then, define a gadget G as follows:

- $V(G)$ is the union of $A, B, T_{1,2}$ and $T_{3,4}$;
- $A, B, v_1 T_{1,2}, v_2 T_{1,2}, v_3 T_{3,4}$ and $v_4 T_{3,4}$ are in $E(G)$.

By definition, G is a $(3k, A, B)$ -gadget.

To prove that G is good, we need to find two perfect matchings in G with different colour profiles. Define $M_A := \{A, v_2 T_{1,2}, v_4 T_{3,4}\}$ and $M_B := \{B, v_1 T_{1,2}, v_3 T_{3,4}\}$. Both M_A and M_B are perfect matchings in G . While M_A has at least two C_j -edges ($v_2 T_{1,2}$ and $v_4 T_{3,4}$), M_B has at least two C_i -edges ($v_1 T_{1,2}$ and $v_3 T_{3,4}$). Thus, M_A and M_B have different colour profiles, as desired. \square

The next lemma ensures a hypergraph H as in Theorem 1.3 contains a good gadget or a perfect matching of huge colour-bias.

Lemma 2.6. *Let $2 \leq \ell < k$ and $\eta > 0$. There exists an $n_0 \in \mathbb{N}$ such that the following holds for all $n \geq n_0$ with $n \in k\mathbb{N}$. Let H be a k -uniform hypergraph on n vertices with an r -colouring $c : E(H) \rightarrow \{C_1, \dots, C_r\}$ and*

$$\delta_\ell(H) \geq (c_{k,\ell} + \eta) \binom{n}{k-\ell}.$$

Suppose that H does not have a perfect matching containing at least $n/k - \binom{r}{2}$ edges of the same colour. Then

- there exists a good $(3k, e, f)$ -gadget in H , for some $e, f \in E(H)$; or
- there exists a good $(k^2 + k, e, f)$ -gadget in H , for some $e, f \in E(H)$.

Proof. Let H and c be as in the lemma and suppose n is sufficiently large. Let $D := k^2 + k \geq 3k$. Note that, given our minimum ℓ -degree condition, Proposition 2.4 implies that

$$(2) \quad \delta_1(H) \geq (c_{k,\ell} + \eta) \binom{n-1}{k-1} > \left(\frac{1}{2} + \frac{\eta}{2}\right) \binom{n}{k-1} \quad \text{and} \quad \delta_2(H) \geq (c_{k,\ell} + \eta) \binom{n-2}{k-2} > \frac{1}{2} \binom{n}{k-2}.$$

Here the inequalities follow as $c_{k,\ell} \geq 1/2$ by (1).

As n is sufficiently large, and by definition of $c_{k,\ell}$, the minimum ℓ -degree condition ensures a perfect matching M in H .

Let $L := \binom{r}{2} + 1$. By the hypothesis of the lemma, M does not contain $n/k - \binom{r}{2}$ edges of the same colour; so there exist distinct edges $e_1, \dots, e_L, f_1, \dots, f_L \in M$ such that $c(e_i) \neq c(f_i)$ for each $i \in [L]$.

Given any distinct $x, y \in V(H)$, (2) implies that $|N_H(x) \cap N_H(y)| \geq \eta \binom{n}{k-1}$. In particular, this means that $N_H(x) \cap N_H(y)$ is of type $S(D)$ or of type $C_i C_j(D)$ for some distinct $i, j \in [r]$.

Suppose there exists $i \neq j \in [r]$ and distinct $x, y, z, w \in V(H)$ such that $N_H(x) \cap N_H(y)$ and $N_H(z) \cap N_H(w)$ are both type $C_i C_j(D)$. Then by Lemma 2.5, there exists a good $(3k, e, f)$ -gadget in H , for some $e, f \in E(H)$.

So we may assume no such $i \neq j \in [r]$ and $x, y, z, w \in V(H)$ exist. In particular, for each of the $\binom{r}{2} = L - 1$ choices for $i \neq j \in [r]$, there is at most one pair (e_s, f_s) such that there exist $u \in e_s$ and $v \in f_s$ so that either $N_H(u) \cap N_H(v)$ or $N_H(v) \cap N_H(u)$ is type $C_i C_j(D)$. Thus, the following claim holds.

Claim 2.7. *There is a pair (e_s, f_s) such that for each $u \in e_s$ and $v \in f_s$ we have that $N_H(u) \cap N_H(v)$ is type $S(D)$.*

Let $e_s = \{u_1, \dots, u_k\}$ and $f_s = \{v_1, \dots, v_k\}$. For each $i \in [k]$, we choose a set T_i so that

- $T_i \in S(u_i v_i)$;
- $T_1, \dots, T_k, e_s, f_s$ are all vertex-disjoint.

Note we can guarantee (ii) since $|S(u_i v_i)| \geq D n^{k-2} = (k^2 + k) n^{k-2}$ for each $i \in [k]$.

We construct a $(k^2 + k, e_s, f_s)$ -gadget G as follows:

- $V(G)$ is the union of $e_s, f_s, T_1, \dots, T_k$;
- e_s and f_s are edges in G ;
- $u_i T_i, v_i T_i$ are edges in G for all $i \in [k]$.

By definition, G is an S - $(k^2 + k, e_s, f_s)$ -gadget with $c(e_s) \neq c(f_s)$. This implies that G is a good $(k^2 + k, e_s, f_s)$ -gadget. Indeed, $M_e := \{e_s, v_1 T_1, \dots, v_k T_k\}$ and $M_f := \{f_s, u_1 T_1, \dots, u_k T_k\}$ are perfect matchings in G with different colour profiles. \square

3. PROOF OF THEOREM 1.3

Let H be a sufficiently large n -vertex k -uniform hypergraph as in the statement of the theorem. Let $c : E(H) \rightarrow \{C_1, \dots, C_r\}$ be an r -colouring of $E(H)$. If H contains a perfect matching with at least $n/k - \binom{r}{2}$ edges of the same colour, then we are done.

So, suppose no perfect matching in H contains at least $n/k - \binom{r}{2}$ edges of the same colour. By Lemma 2.6, we can find either a good $(3k, e, f)$ -gadget or a good $(k^2 + k, e, f)$ -gadget in H . Call this gadget G_1 .

Next consider $H_1 := H \setminus V(G_1)$. Clearly $\delta_\ell(H_1) \geq (c_{k,\ell} + \eta/2) \binom{n}{k-\ell}$. Suppose H_1 contains a perfect matching M_1 with at least $|H_1|/k - \binom{r}{2}$ edges of the same colour. Thus, by taking any perfect matching in G_1 and adding it to M_1 , we obtain a perfect matching in H containing at least $|H_1|/k - \binom{r}{2} \geq n/k - |G_1|/k - \binom{r}{2} \geq n/k - k - 1 - \binom{r}{2}$ edges of the same colour, as desired.

Hence, we may assume H_1 does not contain such a perfect matching M_1 . By Lemma 2.6, we can find either a good $(3k, e, f)$ -gadget or a good $(k^2 + k, e, f)$ -gadget in H_1 . Call this gadget G_2 and set $H_2 := H_1 \setminus V(G_2)$.

Repeating this argument, we either obtain a perfect matching in H of significant colour-bias, or a collection of $t := \frac{\eta n}{4rk^k(k^2+k)}$ vertex-disjoint gadgets G_1, \dots, G_t where, given any $i \in [t]$, G_i is either a good $(3k, e, f)$ -gadget or a good $(k^2 + k, e, f)$ -gadget in H . In particular, note that each gadget we select has size at most $k^2 + k$, and if one removes $t(k^2 + k)$ vertices from H one still has that $\delta_\ell(H) \geq (1/2 + \eta) \binom{n}{k-\ell} - t(k^2 + k)n^{k-\ell-1} \geq (1/2 + \eta/2) \binom{n}{k-\ell}$. Thus, we can indeed repeatedly apply Lemma 2.6 to obtain these gadgets G_1, \dots, G_t .

Set $\mathcal{G} := \{G_1, \dots, G_t\}$. For each colour C_i , consider the set \mathcal{G}_i of all the gadgets in \mathcal{G} that contain two perfect matchings with different colour profiles with respect to the colour C_i . Clearly there exists some $j \in [r]$ such that \mathcal{G}_j contains at least t/r gadgets.

For each gadget G_i in \mathcal{G}_j consider the perfect matching M_i in G_i with the largest possible number of edges coloured C_j ; let M'_i be the perfect matching in G_i with the fewest possible edges coloured C_j . So M_i has at least one more C_j -edge than M'_i .

Let M^+ denote the union of all these M_i and let M^- denote the union of all these M'_i . So M^+ contains at least $t/r = \frac{\eta n}{4rk^k(k^2+k)}$ more C_j -edges than M^- .

Let $V(\mathcal{G}_j)$ denote the set of vertices in H that lie in one of the gadgets in \mathcal{G}_j . Note that $\delta_\ell(H \setminus V(\mathcal{G}_j)) \geq (c_{k,\ell} + \eta/2) \binom{n}{k-\ell}$ so there exists a perfect matching M in $H \setminus V(\mathcal{G}_j)$. Thus, $M \cup M^+$ and $M \cup M^-$ are both perfect matchings in H .

If $M \cup M^-$ contains at least $\frac{n}{rk} + \frac{\eta n}{8r(r-1)k^k(k^2+k)}$ edges of the same colour then the theorem holds. Thus, we may assume this is not the case. This immediately implies the following claim.

Claim 3.1. *For every $i \in [r]$, the number of C_i -edges in $M \cup M^-$ is at least $\frac{n}{rk} - \frac{\eta n}{8rk^k(k^2+k)}$.*

In particular, $M \cup M^-$ contains at least $\frac{n}{rk} - \frac{\eta n}{8rk^k(k^2+k)}$ C_j -edges. Since there are at least $\frac{\eta n}{4rk^k(k^2+k)}$ more C_j -edges in M^+ than in M^- , we obtain that $M \cup M^+$ contains at least $\frac{n}{rk} + \frac{\eta n}{8rk^k(k^2+k)}$ C_j -edges, as desired. \square

4. CONCLUDING REMARKS

In this paper we have determined the minimum ℓ -degree threshold for forcing a colour-bias perfect matching in a k -uniform hypergraph for all $2 \leq \ell < k$. The only remaining open case of the problem is the minimum *vertex* degree version.

A result of Hàn, Person and Schacht [11] yields that $m_1(3, n) = (5/9 + o(1)) \binom{n-1}{2}$. The following example shows that the corresponding colour-bias problem has a significantly higher minimum vertex degree threshold.

Example 4.1. *Given any $n \in 6\mathbb{N}$, there exists an n -vertex 3-uniform hypergraph H with*

$$\delta_1(H) \geq \frac{3}{4} \binom{n-1}{2}$$

and a 2-colouring of $E(H)$ so that every perfect matching in H has precisely $n/6$ edges in each colour.

Proof. Define H so that (i) $V(H)$ is the disjoint union of two vertex classes A and B , both of size $n/2$; (ii) $E(H)$ consists of all those 3-uniform edges containing at least one vertex from each

of A and B . Thus,

$$\delta_1(H) = \binom{n/2}{2} + \frac{n}{2} \left(\frac{n}{2} - 1 \right) \geq \frac{3}{4} \binom{n-1}{2}.$$

Colour each edge containing 2 vertices from A red; each edge containing 2 vertices from B blue. It is easy to see that every perfect matching in H uses the same number of red and blue edges. \square

We suspect that this example is extremal for the minimum vertex degree problem in 3-uniform hypergraphs.

Question 4.2. *Given any $\eta > 0$ does there exist a $\gamma > 0$ so that the following holds for all sufficiently large $n \in 3\mathbb{N}$? Suppose that H is an n -vertex 3-uniform hypergraph with*

$$\delta_1(H) \geq \left(\frac{3}{4} + \eta \right) \binom{n-1}{2}.$$

Then given any 2-colouring of $E(H)$ there is a perfect matching in H with at least $n/6 + \gamma n$ edges of the same colour.

Remark. Question 4.2 is answered in the affirmative in a forthcoming paper of Hiép Hàn, Richard Lang, João Pedro Marciano, Matías Pavez-Signé, Nicolás Sanhueza-Matamala, and the second and third authors. In fact, this new work resolves the minimum vertex degree problem fully (i.e., for all choices of the uniformity $k \geq 3$ and number of colours $r \geq 2$).

By tweaking the proof of Theorem 1.3, one can show that given any $k \geq 3$ and $r \geq 2$, there is a constant C such that every sufficiently large r -coloured n -vertex k -uniform hypergraph H with $\delta_{k-1}(H) \geq n/2 + C$ contains a perfect matching with at least $(n/rk) + 1$ edges of the same colour. Moreover, the lower bound on the colour-bias grows linearly as one increases the minimum codegree further. The PhD thesis of the third author will contain a rigorous proof of this.

ACKNOWLEDGMENTS

Part of the research in this paper was carried out during a visit by the first author to the University of Birmingham in July 2023. The authors are grateful to the BRIDGE strategic alliance between the University of Birmingham and the University of Illinois at Urbana-Champaign, which partially funded this visit. The authors are also grateful to the referees for their careful reviews.

Open access statement. This research was funded in part by EPSRC grant EP/V002279/1. For the purpose of open access, a CC BY public copyright licence is applied to the Author Accepted Manuscript version of this paper.

Data availability statement. There are no additional data beyond that contained within the main manuscript.

REFERENCES

- [1] J. Balogh, B. Csaba, Y. Jing and A. Pluhár, On the discrepancies of graphs, *Electron. J. Combin.*, **27** (2020), P2.12.
- [2] J. Balogh, B. Csaba, A. Pluhár and A. Treglown, A discrepancy version of the Hajnal–Szemerédi theorem, *Combin. Probab. Comput.*, **30** (2021), 444–459.
- [3] D. Bradač, Powers of Hamilton cycles of high discrepancy are unavoidable, *Electron. J. Combin.*, **29** (2022), P3.22.
- [4] D. Bradač, M. Christoph and L. Gishboliner, Minimum Degree Threshold for H -factors with High Discrepancy, arXiv:2302.13780.
- [5] P. Erdős, Ramsey és Van der Waerden tételével kapcsolatos kombinatorikai kérdésekről, *Mat. Lapok.*, **14** (1963), 29–37.
- [6] P. Erdős and J.H. Spencer, Imbalances in k -colorations, *Networks*, **1** (1971/72), 379–385.

- [7] A. Freschi, J. Hyde, J. Lada and A. Treglown, A note on color-bias Hamilton cycles in dense graphs, *SIAM J. Discr. Math.* **35** (2021), 970–975.
- [8] L. Gishboliner, S. Glock and A. Sgueglia, Tight Hamilton cycles with high discrepancy, arXiv:2312:09976.
- [9] L. Gishboliner, M. Krivelevich and P. Michaeli, Color-biased Hamilton cycles in random graphs, *Random Structures & Algorithms*, **60** (2022), 289–307.
- [10] L. Gishboliner, M. Krivelevich and P. Michaeli, Discrepancies of spanning trees and Hamilton cycles, *J. Combin. Theory Ser. B* **154** (2022), 262–291.
- [11] H. Hàn, Y. Person and M. Schacht, On perfect matchings in uniform hypergraphs with large minimum vertex degree, *SIAM J. Discrete Math.* **23** (2009), 732–748.
- [12] D. Kühn and D. Osthus, Matchings in hypergraphs of large minimum degree, *J. Graph Theory* **51** (2006), 269–280.
- [13] D. Kühn and D. Osthus. Embedding large subgraphs into dense graphs. In *Surveys in combinatorics 2009*, volume 365 of *London Math. Soc. Lecture Note Ser.*, pages 137–167. Cambridge Univ. Press, Cambridge, 2009.
- [14] C.J. Mansilla Brito, Discrepancia de ciclos hamiltonianos en hipergrafos 3-uniformes, Master’s thesis, Universidad de Concepción, 2023.
- [15] O. Pikhurko, Perfect matchings and K_4^3 -tilings in hypergraphs of large codegree, *Graphs Combin.* **24** (2008), 391–404.
- [16] V. Rödl and A. Ruciński, Dirac-type questions for hypergraphs – a survey (or more problems for Endre to solve), *An Irregular Mind*, Bolyai Soc. Math. Studies **21** (2010), 561–590.
- [17] V. Rödl, A. Ruciński and E. Szemerédi, Perfect matchings in uniform hypergraphs with large minimum degree, *European J. Combin.* **27** (2006), 1333–1349.
- [18] V. Rödl, A. Ruciński and E. Szemerédi, Perfect matchings in large uniform hypergraphs with large minimum collective degree, *J. Combin. Theory Ser. A* **116** (2009), 613–636.
- [19] A. Treglown and Y. Zhao, Exact minimum degree thresholds for perfect matchings in uniform hypergraphs II, *J. Combin. Theory A* **120** (2013), 1463–1482.
- [20] A. Treglown and Y. Zhao. A note on perfect matchings in uniform hypergraphs. *Electron. J. Combin.*, **23** (2016), P1.16.
- [21] Y. Zhao. Recent advances on Dirac-type problems for hypergraphs, *Recent Trends in Combinatorics*, the IMA Volumes in Mathematics and its Applications 159. Springer, New York, 2016. Vii 706.