

Minimal cover groups

Cameron, Peter; Craven, David; Dorbidi, Hamid Reza; Harper, Scott; Sambale, Benjamin

DOI:

[10.1016/j.jalgebra.2024.06.038](https://doi.org/10.1016/j.jalgebra.2024.06.038)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Cameron, P, Craven, D, Dorbidi, HR, Harper, S & Sambale, B 2024, 'Minimal cover groups', *Journal of Algebra*, vol. 660, pp. 345-372. <https://doi.org/10.1016/j.jalgebra.2024.06.038>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Contents lists available at ScienceDirect

Journal of Algebra

journal homepage: www.elsevier.com/locate/jalgebra

Research Paper

Minimal cover groups

Peter J. Cameron^{a,*}, David Craven^b, Hamid Reza Dorbidi^c,
Scott Harper^a, Benjamin Sambale^d^a School of Mathematics and Statistics, University of St Andrews, St Andrews,
Fife KY16 9SS, UK^b School of Mathematics, University of Birmingham, Edgbaston, Birmingham
B15 2TT, UK^c Department of Mathematics, Faculty of Science, University of Jiroft, Jiroft
78671-61167, Iran^d Institut für Algebra, Zahlentheorie und Diskrete Mathematik, Leibniz Universität
Hannover, 30167 Hannover, Germany

ARTICLE INFO

Article history:

Received 27 February 2024

Available online 23 July 2024

Communicated by Tim Burness

Keywords:

Cauchy's theorem

Cayley's theorem

Simple groups

Abelian groups

ABSTRACT

Let \mathcal{F} be a set of finite groups. A finite group G is called an \mathcal{F} -cover if every group in \mathcal{F} is isomorphic to a subgroup of G . An \mathcal{F} -cover is called *minimal* if no proper subgroup of G is an \mathcal{F} -cover, and *minimum* if its order is smallest among all \mathcal{F} -covers. We prove several results about minimal and minimum \mathcal{F} -covers: for example, every minimal cover of a set of p -groups (for p prime) is a p -group (and there may be finitely or infinitely many, for a given set); every minimal cover of a set of perfect groups is perfect; and a minimum cover of a set of two nonabelian simple groups is either their direct product or simple. Our major theorem determines whether $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ has finitely many minimal covers, where q and r are distinct primes. Motivated by this, we say that n is a *Cauchy number* if there are only finitely many groups which are minimal (under inclusion) with respect to having order divisible by n , and we determine all such numbers. This extends Cauchy's theorem.

* Corresponding author.

E-mail address: pjc20@st-andrews.ac.uk (P.J. Cameron).

We also define a dual concept where subgroups are replaced by quotients, and we pose a number of problems.

© 2024 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Cayley's celebrated theorem asserts that every group of order n is isomorphic to a subgroup of the symmetric group S_n . This motivates the following problem: given a finite set \mathcal{F} of finite groups, find a group G such that every group in \mathcal{F} is isomorphic to a subgroup of G . We call a group G with this property an \mathcal{F} -cover. When \mathcal{F} consists of all the groups of order n up to isomorphism, which is the case occurring in Cayley's theorem, we refer to an \mathcal{F} -cover as an n -cover.

It is natural to ask what the smallest \mathcal{F} -cover is, and to make this precise we introduce the following definitions. For a finite set \mathcal{F} of finite groups, we say that an \mathcal{F} -cover is

- *minimal* if no proper subgroup of G is an \mathcal{F} -cover;
- *co-minimal* if no proper quotient of G is an \mathcal{F} -cover;
- *strongly minimal* if it is both minimal and co-minimal;
- *minimum* if no \mathcal{F} -cover has smaller order.

Note that a minimum cover is strongly minimal.

In this paper, we will mainly focus on minimal and minimum covers, guided by the following two questions.

Question A. For which finite sets \mathcal{F} of finite groups are there only finitely many minimal \mathcal{F} -covers up to isomorphism?

Question B. Given a group theoretic property \mathbf{P} , is it true that if every group in \mathcal{F} has property \mathbf{P} , then every (or perhaps some) minimum (or minimal) \mathcal{F} -cover has property \mathbf{P} ?

Our first theorem, which we prove in Section 3, fully answers Question A in an important special case.

Theorem 1. *Let $q < r$ be primes. The set $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ has only finitely many minimal covers if and only if $q = 2$ and r is a Fermat prime, in which case there are exactly three: \mathbb{Z}_{2r} , D_{2r} and $\mathbb{Z}_2^{2^a} : \mathbb{Z}_r$ where $r = 2^a + 1$.*

Cauchy's theorem asserts that if a prime p divides the order of a finite group G , then G has an element of order p , or, equivalently, has a subgroup isomorphic to \mathbb{Z}_p . More generally, Sylow's theorem asserts that if a prime power p^a divides the order of G , then

G has a subgroup isomorphic to one of the (finitely many) groups of order p^a . These theorems are sharp, in the sense that if every group of order divisible by $n > 1$ necessarily has an element of order n , then n is prime, and if it necessarily has a subgroup of order n , then n is a prime power (see [21]). Nevertheless, Theorem 1 shows that for $n = 2p$ where p is a Fermat prime, there is a set \mathcal{W} of three groups of order divisible by n such that if n divides the order of G , then G has a subgroup isomorphic to a group in \mathcal{W} . More generally, we say that n is a *Cauchy number* if there is a finite set \mathcal{W} of groups of order divisible by n such that if n divides the order of G , then G has a subgroup isomorphic to a group in \mathcal{W} . Cauchy numbers form the focus of Section 4, where we completely determine them.

Theorem 2. *Let n be a natural number. Then n is a Cauchy number if and only if one of the following holds:*

- (a) n is a prime power;
- (b) $n = 6$;
- (c) $n = 2p^a$, where $p > 3$ is a Fermat prime and $a \geq 1$.

Remark 3. Theorem 1 guarantees that every group of order divisible by 6 contains a subgroup isomorphic to \mathbb{Z}_6 , S_3 or A_4 . While our proof of Theorem 1 uses the Classification of Finite Simple Groups, as we explain in Remark 3.10, this important special case can be proved without it.

Remark 4. We say that G is an n -*witness* if n divides the order of G but n does not divide the order of any proper subgroup of G . Then n is a Cauchy number if and only if there are finitely many n -witnesses up to isomorphism. If $n = p^a$, then the n -witnesses are the groups of order n , and if $n = 2p$, for a Fermat prime p , then the n -witnesses are \mathbb{Z}_{2p} , D_{2p} , $\mathbb{Z}_2^{2a} : \mathbb{Z}_p$ as in Theorem 1. Now consider $n = 2p^a$ for a Fermat prime $p > 3$ and $a > 1$. Proposition 4.5 shows that any n -witness has order dividing $2^d p^a$, where d is the maximum degree of a (not necessarily faithful) irreducible representation of a group of order p^a over \mathbb{F}_2 . This is sharp, since if P is a group of order p^a and V is an irreducible $\mathbb{F}_2 P$ -module, then $V:P$ is a n -witness.

In the process of proving Theorem 2 we establish the following, which could be of independent interest (see Proposition 3.5(a)).

Theorem 5. *Every finite simple group with order divisible by 60 has a subgroup isomorphic to A_5 .*

Turning to Question B, in Section 5, we begin with simple groups.

Theorem 6. *Let M and N be nonabelian finite simple groups. If G is a minimum cover of $\{M, N\}$ then either $G = M \times N$ or G is simple and $|G| \leq |M| \cdot |N|$.*

Remark 7. Both possibilities in Theorem 6 arise. In fact, they can arise simultaneously. By Corollary 5.7, A_{12} and $A_7 \times M_{12}$ are both minimum covers of $\{A_7, M_{12}\}$. In particular, $\{M, N\}$ can have nonisomorphic minimum covers, but we do not know whether $\{M, N\}$ can have nonisomorphic minimum *simple* covers. Section 5 gives further results and questions in this direction.

Section 6 considers p -groups. Every minimal cover for a set of p -groups is a p -group, and we focus on p^n -covers, i.e. \mathcal{F} -covers where \mathcal{F} is the set of all groups of order p^n . We prove that there are only finitely many minimal p^2 -covers, but infinitely many minimal 2^3 -covers (see Theorem 6.5), and we determine (rather weak) upper and lower bounds for the order of a minimum p^n -cover: the upper bound is $p^{(p^n-1)/(p-1)}$, and the lower bound $p^{(c+o(1))n^2}$ with $c = 2/27$ (see Theorem 6.9).

Many other properties are inherited by minimal or minimum covers. For instance, if every group in \mathcal{F} is perfect, then every minimal \mathcal{F} -cover is perfect (see Example 2.8(a)), and if every group in \mathcal{F} is nilpotent, then every minimum \mathcal{F} -cover is nilpotent (see Theorem 7.3). However, for other properties, this fails. For instance, if every group in \mathcal{F} is soluble, then there need not exist a soluble \mathcal{F} -cover (see Example 7.8).

We leave open the question of whether if \mathcal{F} is a finite set of finite abelian groups, then there necessarily exists an abelian minimum \mathcal{F} -cover. However, Section 7 does present a number of results in this direction. In particular, the main result of that section is an algorithm to find the smallest abelian group containing a given set of finite abelian groups. Since abelian groups are direct products of their Sylow subgroups, it suffices to solve the problem for abelian p -groups. Our algorithm has the following consequence (see Corollary 7.6).

Theorem 8. *The smallest abelian group which embeds all abelian groups of order p^n is unique up to isomorphism and has order $p^{f(n)}$, where*

$$f(n) = \sum_{k=1}^n \lfloor n/k \rfloor.$$

This function f was studied by Dirichlet who found its asymptotic behavior. It is sequence A006218 in the On-Line Encyclopedia of Integer Sequences [24], where several further areas where it arises are given.

We conclude the paper with a dual version of the ideas introduced in the paper and number of further directions and open problems.

Notation. Our notation is fairly standard, but note that we write \mathbb{Z}_n for the cyclic group of order n and D_{2n} for the dihedral group of order $2n$. For sporadic simple groups we follow the ATLAS of Finite Groups [5]. We write n_p for the p -part of n , and $o_p(n)$ for the order of n modulo p .

Acknowledgments. We are grateful to Jon Awbrey and Michael Kinyon for helpful comments. The fourth author is an EPSRC Postdoctoral Fellow (EP/X011879/1). In order to meet institutional and research funder open access requirements, any accepted manuscript arising shall be open access under a Creative Commons Attribution (CC BY) reuse licence with zero embargo.

2. General results

We begin with a simple result to illustrate the central concepts of the paper, showing in particular that any finite set of finite groups has a cover.

Proposition 2.1. *Let \mathcal{F} be a finite set of finite groups, and let G be an \mathcal{F} -cover. Then the following hold:*

- (a) *We have that $\text{lcm}\{|F| : F \in \mathcal{F}\}$ divides $|G|$.*
- (b) *If G is a minimum \mathcal{F} -cover, then $|G| \leq \prod_{F \in \mathcal{F}} |F|$.*
- (c) *If the orders of the groups in \mathcal{F} are pairwise coprime, then equality holds in (b).*

Proof. (a) is immediate from Lagrange's Theorem; (b) follows from the fact that the direct product of the groups in \mathcal{F} is a cover; and (c) is immediate from (a) and (b). \square

Note that if the orders of the groups in \mathcal{F} are powers of distinct primes, then a minimum \mathcal{F} -cover is a group which has these as its Sylow subgroups.

Example 2.2. A minimum cover of $\{\mathbb{Z}_3, (\mathbb{Z}_2)^2, \mathbb{Z}_5\}$ has order 60; any group of order 60 having these three groups as its Sylow subgroups is an example. This holds for seven of the 13 groups of order 60, including $S_3 \times D_{10}$, $A_4 \times \mathbb{Z}_5$, and A_5 .

Proposition 2.1 raises the following natural problem.

Problem 2.3. Characterize sets \mathcal{F} of finite groups for which the order of a minimum \mathcal{F} -cover is the least common multiple of the orders of the groups in \mathcal{F} .

Proposition 2.1 shows that the order of a minimum \mathcal{F} -cover is bounded by a function of \mathcal{F} (namely, the product of the orders of the groups in \mathcal{F}). In particular, there are only finitely many minimum \mathcal{F} -covers. However, there may be infinitely many minimal \mathcal{F} -covers, which motivates Question A in the introduction. In this context, the following result will be useful.

Proposition 2.4. *Let π be a finite set of primes, and \mathcal{F} a finite set of π -groups. If there exists a minimal \mathcal{F} -cover which is not a π -group, then there exist infinitely many minimal \mathcal{F} -covers.*

Proof. Let G be a minimal \mathcal{F} -cover whose order is divisible by a prime $p \notin \pi$. By a theorem of Gaschütz (see [8, Chapter B, Theorem 11.8]), there exists a *Frattini extension* H of G , which has a non-trivial elementary abelian normal p -subgroup $E \leq \Phi(H)$ such that $H/E \cong G$. Let $M < H$ be a maximal subgroup. Then $E \leq \Phi(H) \leq M$. Suppose by way of contradiction that M is an \mathcal{F} -cover. For every $F \in \mathcal{F}$, we may assume that $F \leq M$. Since F is a π -group, it follows that $F \cap E = 1$ and

$$F \cong FE/E \leq M/E < H/E \cong G.$$

But this implies that M/E is an \mathcal{F} -cover. This contradiction shows that H is a minimal \mathcal{F} -cover. Now we can repeat this process with H instead of G . This yields an infinite series of minimal \mathcal{F} -covers. \square

Let us now turn to Question B on whether minimal covers inherit group-theoretic properties. We begin with some simple observations.

Proposition 2.5. *Let p be a prime, and let \mathcal{F} be a finite set of finite p -groups. Then every minimal \mathcal{F} -cover is a p -group.*

Proof. Let G be a minimal \mathcal{F} -cover, and P a Sylow p -subgroup of G . Then every group in \mathcal{F} is embedded in G , and so conjugate to a subgroup of P . Hence P is an \mathcal{F} -cover. By minimality, $P = G$. \square

Proposition 2.6. *Let \mathcal{A} be a class of finite groups which is closed under the taking of subgroups and direct products. If \mathcal{F} is a finite subset of \mathcal{A} then there exists an \mathcal{A} -group which is a minimal \mathcal{F} -cover.*

Proof. Let $G = \prod_{H \in \mathcal{F}} H$. Then G is an \mathcal{A} -group which is also an \mathcal{F} -cover. So G contains a minimal \mathcal{F} -cover which is an \mathcal{A} -group. \square

It is not clear when we can obtain a minimum \mathcal{F} -cover in this way. We will see in Section 7 that this is the case for nilpotent groups, but we have been unable to decide the apparently easier case of abelian groups.

The following two theorems give further results in this direction.

Proposition 2.7. *Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, none of which has a non-trivial \mathcal{X} -group as a quotient, and let G be a minimal \mathcal{F} -cover. Then G has no non-trivial \mathcal{X} -group as a quotient.*

Proof. Suppose that $G/N \in \mathcal{X}$. Then for any group $H \in \mathcal{F}$ with $H \leq G$, we have $H/(H \cap N) \cong HN/N \leq G/N \in \mathcal{X}$. So $H/(H \cap N) \in \mathcal{X}$ which implies $H \subseteq N$. By minimality of G , we have $N = G$. \square

Example 2.8. Let us record an application of Proposition 2.7. Let \mathcal{X} be the class of finite abelian groups. The condition that G has no non-trivial homomorphism to an \mathcal{X} -group means that G is perfect. So we deduce that, if every group in \mathcal{F} is perfect, then any minimal \mathcal{F} -cover is perfect.

There is a dual result, as follows.

Proposition 2.9. *Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, and suppose that no group in \mathcal{F} has a non-trivial normal \mathcal{X} -subgroup. Let G be a co-minimal cover of \mathcal{F} . Then G has no non-trivial normal \mathcal{X} -subgroup.*

Proof. Let G be a co-minimal cover of \mathcal{F} , and suppose that G has a normal \mathcal{X} -subgroup N . For each group $H \in \mathcal{F}$, we have $H \cap N \in \mathcal{X}$. So $H \cap N = \{1\}$. Hence $H \cong H/(H \cap N) \cong HN/N \leq G/N$. By co-minimality, $G/N \cong G$, so $N = \{1\}$. \square

Example 2.10. We now give an application of Proposition 2.9. Let \mathcal{X} be the class of finite abelian groups. If no group in \mathcal{F} has a non-trivial abelian normal subgroup, then a co-minimal \mathcal{F} -cover has no non-trivial abelian normal subgroup.

We conclude with some remarks on n -covers, which, recall, are \mathcal{F} -covers where \mathcal{F} consists of the groups of order n .

Proposition 2.11. *Let $n = p_1^{\alpha_1} \cdots p_k^{\alpha_k}$ and G be a minimal n -cover. If P_i is a Sylow p_i -subgroup of G then P_i is a $p_i^{\alpha_i}$ -cover.*

Proof. Let P be a group of order $p_i^{\alpha_i}$. Then $H = P \times \prod_{j \neq i} \mathbb{Z}_{p_j^{\alpha_j}}$ is a group of order n . So H is isomorphic to a subgroup of G . Hence P is isomorphic to a subgroup of P_i . \square

Question 2.12. In the above situation, when is P_i a minimal $p_i^{\alpha_i}$ -cover group?

Remark 2.13. If \mathcal{F} is a set of soluble π -groups, then every soluble minimal \mathcal{F} -cover is a π -group, by Hall’s theorem. Moreover, if G is a cover of a set \mathcal{F} of π -groups, then every \mathcal{F} -subgroup of G embeds in $G/O_{\pi'}(G)$; so every co-minimal \mathcal{F} -cover (and *a fortiori* every minimum \mathcal{F} -cover) G satisfies $O_{\pi'}(G) = 1$.

3. Two cyclic groups of prime order

In this section, we examine minimal covers for $\{\mathbb{Z}_q, \mathbb{Z}_r\}$, where q and r are distinct primes. In particular, we will prove Theorem 1, which determines when there are finitely many such covers.

Our first result handles the generic case.

Proposition 3.1. *Let q and r be distinct odd primes with at least one strictly greater than 5. Then there are infinitely many minimal $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -covers.*

Proof. Without loss of generality we may assume that $q < r$, so $r \geq 7$. By Dirichlet’s theorem, there are infinitely many primes p satisfying $p \equiv 1 \pmod{q}$ and $p \equiv -1 \pmod{r}$. We claim that, for any such prime p , the group $G = \text{PSL}_2(p)$ is a minimal \mathcal{F} -cover, where $\mathcal{F} = \{\mathbb{Z}_q, \mathbb{Z}_r\}$. To see this, we consult the list of subgroups of $\text{PSL}_2(p)$ given in Suzuki [26, Theorem 3.6.25]. The subgroup A_5 has order not divisible by $r \geq 7$, and $r \nmid p(p-1)$, so the only subgroup of order divisible by r is dihedral of order $p+1$. But this cannot be divisible by $q \mid (p-1)$, so no maximal subgroup of G has order divisible by qr . \square

Remark 3.2. Since $\text{PSL}_2(p)$ is simple, we see that the minimal covers exhibited in the proof of Proposition 3.1 are strongly minimal.

To handle the remaining pairs of primes, we first prove a general result, which reduces our study to insoluble groups.

Proposition 3.3. *Let G be a minimal $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -cover. One of the following holds:*

- (a) *G is soluble, has an elementary abelian normal subgroup N , and G/N is cyclic. Either N is a q -group and $|G/N| = r$, or N is an r -group and $|G/N| = q$. There are three such groups, namely \mathbb{Z}_{qr} , $\mathbb{Z}_q^a : \mathbb{Z}_r$ and $\mathbb{Z}_r^b : \mathbb{Z}_q$, where a is the multiplicative order of q modulo r and b is the multiplicative order of r modulo q .*
- (b) *G is not soluble and has a unique maximal normal subgroup N . (This includes the case where G is simple.) The group N is equal to $\Phi(G)$ (and in particular N is nilpotent), and the quotient G/N is a nonabelian simple group that is also a minimal $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -cover. Moreover, $|N|$ is coprime to qr , but if $|N| \neq 1$, it is not coprime to $|G/N|$.*

Proof. Suppose that G is soluble, so by Hall’s theorem there exists a Hall $\{q, r\}$ -subgroup. Thus by minimality G is a $\{q, r\}$ -group. Let N be a minimal normal subgroup of G , which without loss of generality we may assume to be an elementary abelian q -group. If x has order r in G then $\langle N, x \rangle$ is a $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -cover, so by minimality $G = \langle N, x \rangle$.

To complete the proof of (a) we need to determine a and b . By minimality N is an irreducible module for \mathbb{Z}_r over \mathbb{F}_q . If it is the trivial module we obtain \mathbb{Z}_{qr} , so we may assume that it is non-trivial. Since $\text{Aut}(\mathbb{Z}_r)$ acts transitively on the set of isomorphism classes of non-trivial modules over an algebraically closed field (of characteristic different from r), we see that $\text{Aut}(\mathbb{Z}_r)$ acts transitively on all non-trivial irreducible modules over any field, in particular \mathbb{F}_q , and this implies there is a unique group $(\mathbb{Z}_q)^a : \mathbb{Z}_r$ up

to isomorphism. So it suffices to determine a : but if d is the multiplicative order of q modulo r then \mathbb{Z}_r is a subgroup of $\text{GL}_d(q)$ (since the cyclotomic polynomial $\Phi_d(q)$ divides $|\text{GL}_d(q)|$, and $r \mid \Phi_d(q)$). It is also not a subgroup of any smaller linear group, so $d = a$. This proves (a).

Thus we may assume that G is not soluble. Let N be any maximal normal subgroup. By minimality of G , at least one of q and r does not divide $|N|$. Suppose exactly one does, so $q \mid |N|$ without loss of generality. If $x \in G$ has order r , then x acts on the set of Sylow q -subgroups of N . The number of these subgroups is prime to r , so x normalizes one, say P . Thus $\langle P, x \rangle$ is a soluble $\{q, r\}$ -group contained in G , contradicting minimality.

Thus each of q and r does not divide $|N|$. In particular, G/N must be nonabelian simple. If G/N is not a minimal $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -cover then we may choose a proper subgroup M/N of G/N with order divisible by qr , whence its preimage M also has order divisible by qr , contradicting minimality of G .

To prove uniqueness of N , let M be any other maximal normal subgroup of G , and note the above discussion also holds for M . Clearly $MN = G$, so $G/N \cong M/(M \cap N)$. But then $|M/(M \cap N)|$ and so $|M|$ are divisible by qr , and this contradicts the minimality of G .

If H is any maximal subgroup of G , then $NH < G$, since otherwise $H/(H \cap N) \cong G/N$ and so H is a cover. But this implies that $N \leq H$. So $N \leq \Phi(G)$, and we must have equality as G/N is simple. Finally, if $\text{gcd}(|N|, |G/N|) = 1$ then the Schur–Zassenhaus theorem implies that N has a complement, which is a cover. \square

Corollary 3.4. *If there are no nonabelian simple minimal $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -covers, then all minimal $\{\mathbb{Z}_q, \mathbb{Z}_r\}$ -covers are soluble.*

We next consider $\{q, r\} = \{3, 5\}$. Here there is a unique simple minimal cover, but infinitely many insoluble covers.

Proposition 3.5.

- (a) *The only simple minimal $\{\mathbb{Z}_3, \mathbb{Z}_5\}$ -cover is the alternating group A_5 .*
- (b) *There are infinitely many insoluble $\{\mathbb{Z}_3, \mathbb{Z}_5\}$ -covers.*

Proof. (a) Let G be a simple group, and we use the classification of finite simple groups. First, it is easy that A_5 actually *is* a minimal $\{\mathbb{Z}_3, \mathbb{Z}_5\}$ -cover so we assume that G is not A_5 . We will show that G always possesses a proper subgroup of order divisible by 15, or $|G|$ is not divisible by 15.

If G is an alternating group then clearly G contains A_5 and we are done. If G is a sporadic group then all maximal subgroups of G are known and we may check the ATLAS of Finite Groups [5] (except for the Monster, where all maximal subgroups have only recently been identified [6]; however, $2 \cdot B$ is a subgroup of M , and we are done).

Thus let G be a group of Lie type. Note that G cannot be a Suzuki group (3 does not divide its order) or a small Ree group (5 does not divide its order). For the large Ree groups, they all contain the Tits group, which contains a maximal subgroup $A_6.2^2$ (see [5, p. 74] for example).

Thus we may assume that G is a simple Chevalley or Steinberg group, in characteristic p . Let $G = G(p^a)$ be the group; in all cases there is a Levi subgroup of G that is either $\text{PSL}_2(p^a)$ or $\text{SL}_2(p^a)$. If $p^a \equiv 0, \pm 1 \pmod{5}$ then 15 divides the order of $\text{PSL}_2(p^a)$, and we are done, unless $G = \text{PSL}(2, p^a)$. But, if $p = 5$, then G contains $\text{PSL}_2(5)$, whereas if $p \equiv \pm 1 \pmod{5}$ then G contains $\text{PSL}(p)$, which contains A_5 because of the congruence condition.

So we may suppose that this is not the case. In particular, $p \equiv \pm 2 \pmod{5}$ and a is odd. These conditions imply that p has order 4 modulo 5. A standard fact about cyclotomic polynomials is that if $5 \mid \Phi_d(p^a)$ then d is a power of 5 times 4. Thus if $5 \mid |G|$, then $\Phi_4(p^a) = p^{2a} + 1$ (or Φ_{20}, Φ_{100} , etc.) divides $|G(p^a)|$, so this excludes PSL_2 , PSL_3 , PSU_3 , 3D_4 and G_2 from consideration. The groups $\text{PSP}_4(p^a)$ contain groups of the form $\text{PSL}_2(p^{2a})$, so these cannot be minimal covers. Since both $\text{PSL}_4(p^a)$ and $\text{PSU}_4(p^a)$ contain $\text{PSP}_4(p^a)$, these cannot be minimal covers either.

The remaining groups we have not considered are symplectic and orthogonal groups (which all contain $\text{PSP}_4(p^a) = \Omega_5(p^a)$) or exceptional groups $F_4(p^a)$, $E_6(p^a)$, $E_7(p^a)$ and $E_8(p^a)$ (each of which contains the previous one and the first one contains $\text{PSp}_4(p^a)$) and ${}^2E_6(p^a)$ (which contains $F_4(p^a)$).

This completes the list of groups of Lie type, so A_5 is the only simple minimal $\{\mathbb{Z}_3, \mathbb{Z}_5\}$ -cover, as claimed.

(b) This follows from (a) and Proposition 2.4 applied to $\pi = \{3, 5\}$. \square

Note that Theorem 5 is an immediate consequence of Proposition 3.5(a).

We now consider the case where $q = 2$. Let us outline our strategy in this case. Suppose that G is a simple minimal $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ -cover. If M is any maximal subgroup of G then M is not divisible by $2r$, so any maximal subgroup of order divisible by r has odd order. Of course, if R denotes a Sylow r -subgroup of G then $N_G(R)$ must be contained in a maximal subgroup of G , which therefore must have odd order. Fortunately, there are very few odd-order maximal subgroups of simple groups, and they are enumerated in [18, Table 2]. The following lemma, which makes use of [18], is stated in greater generality than required here so that it can be used in the next section also.

Lemma 3.6. *Let A be a finite almost simple group with socle S . Let p be a Fermat prime. Let M be a maximal subgroup of A such that $N_S(P) \leq M$ where P is a Sylow p -subgroup of S . Then M has even order, or $p = 3$ and $S = \text{PSL}_2(3^a)$ for an odd integer $a > 1$.*

Proof. Suppose that M has odd order. According to [18, Table 2], Table 1 gives the only possibilities for S and $M \cap S$ where S is a finite nonabelian simple group and M is an odd-order maximal subgroup of an almost simple group with socle S . (Not all examples

Table 1
Possibilities for an odd-order maximal subgroup M of an almost simple group with socle S .

S	$M \cap S$	conditions
A_n	$n: \left(\frac{n-1}{2}\right)$	$n \equiv 3 \pmod{4}$ prime
$\text{PSL}_2(q)$	$q: \left(\frac{q-1}{2}\right)$	$q \equiv 3 \pmod{4}$ prime power
$\text{PSL}_d(q)$	$\left(\frac{1}{(d,q-1)} \frac{q^d-1}{q-1}\right) : d$	d odd prime
$\text{PSU}_d(q)$	$\left(\frac{1}{(d,q+1)} \frac{q^d+1}{q+1}\right) : d$	d odd prime
M_{23}	23:11	
Th	31:15	
\mathbb{B}	47:23	
\mathbb{M}	59:29	
\mathbb{M}	71:35	

in the table necessarily arise. In particular, at the time of writing [18], it was not known if the subgroups in the final two rows were maximal subgroups of the Monster. They have since been proved not to be [15,16].) In the first row, we require $p = n$ but we must have $n \geq 5$ and $n \equiv 3 \pmod{4}$, which is impossible since p is a Fermat prime. In the second row, we require $q = p^a$ but we must have $q \geq 4$ and $q \equiv 3 \pmod{4}$, which, since p is a Fermat prime, forces $p = 3$ and $a > 1$ to be odd; this gives the exception in the statement. For a contradiction, suppose that one of the remaining rows holds. The restriction that p is a Fermat prime means that we must have one of the following:

1. $S = \text{PSL}_d(q)$ and $M = \left(\frac{1}{(d,q-1)} \frac{q^d-1}{q-1}\right) : d$ for an odd prime d .
2. $S = \text{PSU}_d(q)$ and $M = \left(\frac{1}{(d,q+1)} \frac{q^d+1}{q+1}\right) : d$ for an odd prime d .

Since $M \cap S$ contains $N_S(P)$, we may apply the theory of Sylow tori (see [20, Theorem 25.19] for instance). In Case 1, this implies that p is a primitive prime divisor of $q^d - 1$, so $o_p(q) = d$, while in Case 2, this implies that p is a primitive prime divisor of $q^{2d} - 1$, so $o_p(q) = 2d$. In both cases, d divides $p - 1$, which is impossible since d is odd and p is a Fermat prime. \square

Proposition 3.7. *Let r be an odd prime. If r is a Fermat prime then there are no simple minimal $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ -covers, and if r is not a Fermat prime then there are infinitely many simple minimal $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ -covers.*

Proof. First assume that r is a Fermat prime. For a contradiction, suppose that S is a simple minimal $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ -cover. As indicated above, let R be a Sylow r -subgroup of S and let M be a maximal subgroup of S containing $N_S(R)$. Since r divides $|M|$ we must have that $|M|$ is odd. By Lemma 3.6, we deduce that $r = 3$ and $S = \text{PSL}_2(3^a)$ for odd $a > 1$. However, $\text{PSL}_2(3) \cong A_4$ is a subgroup of $S = \text{PSL}_2(3^a)$ that has order divisible by 2 and 3. This establishes that S is not a minimal $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ -cover.

Now assume that r is not a Fermat prime, and let d be an odd prime divisor of $r - 1$. Let p be a prime at least 5 that has order d modulo r , of which there are infinitely

many options by Dirichlet’s theorem on primes in arithmetic progressions, and let $G = \text{PSL}_d(p)$. Then r divides $(p^d - 1)/(p - 1)$ and so divides $|G|$, and if R is a Sylow r -subgroup of G then R is cyclic and $N_G(R)$ has odd order. We claim that $N_G(R)$ is the only maximal subgroup of G containing x , a non-trivial element of R .

To see this we use the work of Guralnick–Penttilä–Praeger–Saxl on ppd-elements [12]. (The prime r is a primitive prime divisor of $q^e - 1$ if it does not divide $q^i - 1$ for $i < e$. An element of $\text{GL}(d, q)$ is a ppd(d, q, e) element if its order is divisible by a primitive prime divisor of $q^e - 1$, where $d/2 < e \leq d$. In our situation, we have $d = e$.) We have set things up so that x is a ppd($d, p; d$)-element. The possible subgroups of $\text{GL}_d(p)$ containing x are enumerated in [12, Examples 2.1–2.9], and almost all of them can immediately be ignored since in our case d is a prime and d appears twice in the phrase ‘ppd($d, p; d$)’. We check each set of examples from [12] in turn.

- Example 2.1 are classical groups, and since $d = e$ is an odd prime, and we are over a prime field, none of these applies.
- Examples 2.2 and 2.3 do not apply since they require $e < d$.
- Examples 2.4 and 2.5 do not apply as d is an odd prime.
- Example 2.6(a) does not apply since it requires $r - 1 = d$.
- For Examples 2.6(b) and 2.6(c), there are three tables of examples; we need that $d = e$ is a prime, and the only option is $3 \cdot A_7 \leq \text{SL}_3(25)$ with $r = 7$. This is an example, but we required that G be over a prime field, so this may be excluded.
- For Example 2.7 there is a table of examples, and we find the options $G = M_{11}$, $(d, p, r) = (5, 3, 11)$ and $G = M_{23}, M_{24}$, $(d, p, r) = (11, 2, 23)$. Since we have (not coincidentally) chosen $p \geq 5$, these are not examples.
- In Example 2.8, e is always even, but in our case $e = d$ is odd.
- In Example 2.9, there are no examples with d a prime and $d = e$.

We thus find that $G = \text{PSL}_d(p)$ is a minimal $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ -cover. As there are infinitely many options for p , we find infinitely many such groups. \square

Remark 3.8. Using [12] we could actually determine *exactly* which simple groups are minimal covers of $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ for r a non-Fermat prime.

We therefore have the following corollary.

Corollary 3.9. *If r is a Fermat prime then the set $\{\mathbb{Z}_2, \mathbb{Z}_r\}$ has just three minimal covers, namely \mathbb{Z}_{2r} , D_{2r} and $(\mathbb{Z}_2)^{2^a} : \mathbb{Z}_r$ where $r = 2^a + 1$.*

Proof. Combining Corollary 3.4 with Proposition 3.7, we see that there are no insoluble covers. Proposition 3.3(a) shows that the three groups in the statement are the only possible soluble covers, and Cauchy’s theorem establishes that they actually are covers. \square

Propositions 3.1, 3.5, 3.7 and Corollary 3.9 together give Theorem 1.

Remark 3.10. Our proofs use the Classification of Finite Simple Groups; however, the fact that there are only three $\{\mathbb{Z}_2, \mathbb{Z}_3\}$ -covers (namely \mathbb{Z}_6 , S_3 and A_4) can be proved without the Classification. Three papers [9,11,23] in 1977 independently determined the finite simple groups with no elements of order 6; and it is straightforward to show that, apart from the Suzuki groups (whose orders are not divisible by 6), they all contain subgroups isomorphic to S_3 or A_4 .

4. Cauchy numbers

Building on the previous section, we now determine the Cauchy numbers, thus proving Theorem 2. Recall that n is a *Cauchy number* if there is a finite set \mathcal{W} of groups of order divisible by n such that if n divides the order of G , then G has a subgroup isomorphic to a group in \mathcal{W} . The main result of the previous section, Theorem 1, asserts that if n is the product of two distinct primes, then n is a Cauchy number if and only if n is twice a Fermat prime.

We will use the term “ n -group” (for positive integer n) for a group whose order is divisible by n , and “ n -witness” for an n -group which has no proper subgroup which is an n -group. So n is a Cauchy number if and only if the set of n -witnesses (up to isomorphism) is finite.

Remark 4.1. It was pointed out to us by Michael Kinyon that there is already a sequence of “Cauchy numbers”, arising in the Laplace summation formula, see [22]. “Cauchy numbers” also occur in the theory of compressible flow in continuum mechanics. However, we think that these topics are sufficiently different that no confusion will ensue.

Proposition 4.2. *A divisor of a Cauchy number is a Cauchy number.*

Proof. It suffices to show that if m is a positive integer and p a prime such that pm is a Cauchy number, then m is a Cauchy number. Let G be an m -witness. We claim that either G or $G \times \mathbb{Z}_p$ is a pm -witness. Suppose that G is not a pm -group. Then $G \times \mathbb{Z}_p$ is a pm -group. Let H be a proper subgroup of $G \times \mathbb{Z}_p$ whose order is divisible by pm , and consider the projection π to the first factor of the direct product. If π restricted to H is not a monomorphism, then $H = K \times \mathbb{Z}_p$ for some $K < G$; then K is not an m -group, so H is not a pm -group. Otherwise, $H \leq G$, contrary to assumption. On the other hand, if G is a pm -group, it is clearly minimal.

Now let \mathcal{W}_m be the set of m -witnesses, and define a map F from \mathcal{W}_m to \mathcal{W}_{mp} given by

$$F(G) = \begin{cases} G & \text{if } G \text{ is a } pm\text{-group,} \\ G \times \mathbb{Z}_p & \text{otherwise.} \end{cases}$$

We claim that F is one-to-one. By the Krull–Schmidt theorem, we need to show that we cannot have m -witnesses G_1 and G_2 with $F(G_1) = G_1 \times \mathbb{Z}_p = G_2 = F(G_2)$. But if this holds, then $G_1 < G_2$, contradicting the minimality of G_2 .

Therefore, $|\mathcal{W}_m| \leq |\mathcal{W}_{pm}|$, whence $|\mathcal{W}_m|$ is finite and m is a Cauchy number. \square

Our aim is to determine the Cauchy numbers. First we show that every positive integer satisfies the analogous condition when we restrict to soluble groups.

Lemma 4.3. *If G is an n -witness with normal subgroup N , then G/N is an $n/\gcd(n, |N|)$ -witness.*

Proof. $|G/N|$ is divisible by $n/\gcd(n, |N|)$. If it is not minimal with this property, let H be a proper subgroup of G containing N with H/N divisible by $n/\gcd(n, |N|)$; then n divides $|H|$, contrary to minimality of G . \square

Proposition 4.4. *For any natural number n , there are only finitely many finite soluble groups which are n -witnesses.*

Proof. The proof is by induction on n ; the induction begins by noting that the theorem is true when n is a prime power. So assume that n is not a prime power.

Let N be a minimal normal subgroup of the soluble n -witness G . Then N is an elementary abelian p -group, for some p dividing n . So $1 < |N| < |G|$. By the inductive hypothesis and Lemma 4.3, G/N is a $n/\gcd(n, |N|)$ -witness, so its order is bounded. Also, as N is a minimal normal subgroup, it is generated by any G/N -orbit on non-identity elements, so $|N| \leq p^{|G/N|}$ is also bounded, and we are done. \square

Proof of Theorem 2. Combining Proposition 4.2 with Theorem 1, we deduce that if n is a Cauchy number which is not a prime power, then $n = 2^b p^a$, where p is a Fermat prime and a and b positive integers. The next step is to show that either $n = 6$, or $n = 2p^a$ where $a \geq 1$ and p is a Fermat prime greater than 3. For this we have to exclude $n = 18$ and $n = 4p$ where p is a Fermat prime (the cases $p = 3$ and $p = 5$ require separate arguments).

- (a) For $n = 12$, let f be an odd prime, let $q = 2^f$ and let $G = \text{PSL}_2(q)$. Then G is a 12-witness. The maximal subgroups of G are, up to conjugacy, $D_{2(q+1)}$, $D_{2(q-1)}$, $2^f : (q-1)$. Since f is odd, we know that $q \equiv 2 \pmod{3}$, so 3 does not divide $2(q-1)$ or $2^f : (q-1)$; and 4 does not divide $2(q+1)$.
- (b) For $n = 18$, again let f be an odd prime, and take $q = 3^f$ and $G = \text{PSL}_2(q)$. Then G is an 18-witness. For its only subgroups of order divisible by 3 are $\text{PSL}_2(3) \cong D_6$ and $q : (q-1)/2$; and $(q-1)/2$ is odd.
- (c) For $n = 20$, note that A_5 is a 20-witness, and so Proposition 2.4 shows that 20 is not a Cauchy number.

- (d) Let $n = 4p$, with p a prime greater than 5. By the Chinese Remainder Theorem, there is an arithmetic progression of numbers r such that $r \equiv 1 \pmod{4}$ and $r \equiv -1 \pmod{p}$. Therefore, by Dirichlet’s Theorem, there are infinitely many such prime numbers r . Fix such a prime r . Let $G = \text{PSL}_2(r)$, which has order $\frac{1}{2}(r - 1)r(r + 1)$. We claim that G is an n -witness. First note that n divides $|G|$, since 4 divides $r - 1$ and p divides $\frac{1}{2}(r + 1)$. We now claim that n divides the order of no proper subgroup of G . For a contradiction, suppose that H is a maximal subgroup of order divisible by n . Since $p > 5$, we know that $H \neq A_5$. Since $p > 2$ and p divides $r + 1$, we know that p does not divide $r - 1$. Therefore, consulting the list of maximal subgroups of $\text{PSL}_2(r)$, we see that $H \cong D_{r+1}$, but 4 does not divide $r + 1$ since $r \equiv 1 \pmod{4}$. This contradicts the fact that n divides $|H|$.

Now we have proved that a number not of the form described in the theorem is not a Cauchy number. So from now on we assume that $n = 2p^a$, where p is a Fermat prime greater than 3 and a a positive integer, and have to show that n is a Cauchy number. By Proposition 4.4 it suffices to show that all n -witnesses are soluble.

Assume that there exists an insoluble n -witness, and let G be one with a chosen minimal, and subject to this $|G|$ minimal. Let N be a proper non-trivial normal subgroup of G . Since G/N is a $n/\text{gcd}(n, |N|)$ -witness, the minimality of n and G shows that G/N is soluble.

We claim that G/N is a p -group. To see this, first note that N is insoluble since G/N is soluble, so, in particular, $|N|$ is even. Fix $N \leq P \leq G$ such that P/N is a Sylow p -subgroup of G/N . Then $n = 2p^a$ divides $|P|$. Since G is a n -witness, we deduce that $G = P$. Therefore, $G/N = P/N$ is a p -group.

Since every proper quotient of G is a p -group, $O^p(G)$ is the unique minimal normal subgroup of G . Write $O^p(G) = S^k$ where S is a nonabelian simple group and $k \geq 1$.

Let P be a Sylow p -subgroup of S and let H be a maximal subgroup of G containing $N_G(P^k)$. Let \hat{P} be a Sylow p -subgroup of G containing P^k . Then $\hat{P} \leq N_G(P^k) \leq H$ since $P^k = \hat{P} \cap S^k \leq \hat{P}$. Therefore, p^a divides $|H|$. In particular, $|H|$ is odd since G is a $2p^a$ -witness, and $S^k \not\leq H$ since H is a proper subgroup of G . In particular, the action of G on G/H is faithful and primitive and the point stabilizer H has odd order. Therefore, the main corollary in [18] restricts the possibilities for G and H . Namely, if we identify S with the first factor of S^k , then $H \cap S = M \cap S$ where M is an odd-order maximal subgroup of an almost simple group with socle S . However, $N_S(P) \leq N_G(P^k) \leq H$, so $N_S(P) \leq M \cap S$, which means that $|H|$ is even, by Lemma 3.6. This contradiction completes the proof. \square

We conclude this section by commenting on the set of n -witnesses when n is a Cauchy number. If $n = p^a$, then the n -witnesses are the groups of order n , and if $n = 2p$, for a Fermat prime p , then Corollary 3.9 implies that the n -witnesses are $\mathbb{Z}_{2p}, D_{2p}, \mathbb{Z}_2^{2a} : \mathbb{Z}_p$ where $p = 2^a + 1$. The following result handles the remaining case.

Proposition 4.5. *Let $n = 2p^a$ for a Fermat prime $p > 3$ and $a > 1$.*

- (a) *Let P be a group of order p^a and let V is an irreducible \mathbb{F}_2P -module. Then $V:P$ is a n -witness.*
- (b) *Any n -witness has order dividing $2^d p^a$ where d is the maximum degree of an representation of group of order p^a over \mathbb{F}_2 .*

Proof. First consider part (a). Let $G = V:P$. Clearly $n = 2p^a$ divides $|G|$. The maximal subgroups of G are P , of order p^a , and $V:H$ where H is a maximal subgroup of P , of order $2p^{a-1}$. This proves that G is an n -witness.

Now consider part (b). Corollary 3.9 gives the result when $a = 1$. Now assume that $a > 1$ and proceed by induction on a . Let G be an n -witness. We know that G is soluble and hence a $\{2, p\}$ -group, by Hall’s theorem. Let M be a minimal normal subgroup of G . If M is a 2-group, then by minimality, G/M must be a group of order p^a , and since M is a minimal normal subgroup, G/M must act irreducibly on M . Now assume that M is a p -group. Suppose that $|M| \geq p^a$. Then by minimality, $|G/M| = 2$. However, M is a minimal normal subgroup, so G/M must act irreducibly on M , which forces $|M| = p$, so $a = 1$, contrary to our assumption. Therefore, $|M| < p^a$, so, by minimality, G/M is a $2p^a/|M|$ -witness, and the claim holds by induction. \square

5. Simple groups

Let us now turn to covers of finite simple groups, beginning with the following general result.

Theorem 5.1. *Let \mathcal{F} be a finite set of finite simple groups with $|\mathcal{F}| = n$, and let G be a minimum \mathcal{F} -cover. Let $N_0 < N_1 < \dots < N_k = G$ be a composition series of G and $\mathcal{F}_i = \{H \in \mathcal{F} : H \text{ is isomorphic to a subgroup of } N_i/N_{i-1}\}$. Then*

- (a) $k \leq n$.
- (b) $\bigcup_{i=1}^k \mathcal{F}_i = \mathcal{F}$.
- (c) N_i/N_{i-1} is a minimum \mathcal{F}_i -cover.
- (d) $\prod_{i=1}^k N_i/N_{i-1}$ is a minimum \mathcal{F} -cover.

Proof. The proof is by induction on k . For $k = 1$ all the statements are trivial. Now assume $k \geq 2$. Let $\mathcal{F}' = \{H \in \mathcal{F} : H \text{ is isomorphic to a subgroup of } N_{k-1}\}$. Then every simple subgroup of G is isomorphic to a subgroup of N_{k-1} or a subgroup of G/N_{k-1} . So $\mathcal{F} = \mathcal{F}' \cup \mathcal{F}_k$. Since G is a minimum \mathcal{F} -cover, \mathcal{F}' and \mathcal{F}_k are nonempty. Also N_{k-1} is an \mathcal{F}' -cover and G/N_{k-1} is an \mathcal{F}_k -cover. If N_{k-1} is not a minimum \mathcal{F}' -cover and M is a minimum \mathcal{F}' -cover then $M \times G/N_{k-1}$ is an \mathcal{F} -cover whose order is less than $|G|$ which is a contradiction. Similarly G/N_{k-1} is a minimum \mathcal{F}_k -cover. So by induction $k - 1 \leq |\mathcal{F}'| \leq n - 1$ which implies $k \leq n$. Also by induction $\bigcup_{i=1}^{k-1} \mathcal{F}_i = \mathcal{F}'$ and N_i/N_{i-1}

is a minimum \mathcal{F}_i -cover for $1 \leq i \leq k - 1$ and $\prod_{i=1}^{k-1} N_i/N_{i-1}$ is a minimum \mathcal{F}' -cover. So $\prod_{i=1}^k N_i/N_{i-1}$ is a minimum \mathcal{F} -cover. \square

We now consider Theorem 6, which we repeat below.

Theorem 6. *Let M and N be nonabelian finite simple groups. If G is a minimum cover of $\{M, N\}$ then either $G = M \times N$ or G is simple and $|G| \leq |M| \cdot |N|$.*

In particular, if there is an $\{M, N\}$ -cover of order less than $|M| \cdot |N|$, then any minimum $\{M, N\}$ -cover is simple.

Before proving Theorem 6, let us note that both possibilities in the theorem can occur as the following two examples demonstrate. These examples can be verified using the ATLAS of Finite Groups [5].

Example 5.2. Let $M = A_5$ and $N = \text{PSL}_2(8)$. The orders of these groups are 60 and 504. Their least common multiple is 2520 and their product is 30240. The only simple groups with order divisible by 2520 and not greater than 30240 are A_7 , A_8 and $\text{PSL}_3(4)$; none of these embeds $\text{PSL}_2(8)$. By the theorem, the unique minimum $\{M, N\}$ -cover is $M \times N$.

Example 5.3. Let $M = A_6$ and $N = \text{PSL}_2(7)$. Their orders are 360 and 168, with least common multiple 2520. There is a unique simple group of order 2520, namely A_7 , which embeds both M and N . Therefore, A_7 is the unique minimum $\{M, N\}$ -cover.

Proof of Theorem 6. By Theorem 5.1, either G is simple, or it has a composition series of length 2 with composition factors M and N .

Suppose, without loss of generality, that G has a normal subgroup isomorphic to M with quotient isomorphic to N . Hence $C_G(M) \triangleleft G$ and $M \cap C_G(M) = Z(M) = \{1\}$. Each element of G acts on M by conjugation. A consequence of the Classification of Finite Simple Groups is that the outer automorphism group of M is soluble. Since G has no non-trivial soluble quotient, we see that each element of G induces an inner automorphism of M . Let $g \in G$. Then there exists $m \in M$ such that for all $x \in M$, we have $g x g^{-1} = m x m^{-1}$. So $m^{-1} g \in C_G(M)$ which implies $g \in M C_G(M)$. Hence $G = M C_G(M)$. Thus G has normal subgroups M and $C_G(M)$ intersecting trivially (and commuting), so is their direct product. \square

Corollary 5.4. *There is a function f such that, if $\mathcal{F} = \{M, N\}$ where M and N are nonabelian finite simple groups with $|N| > f(|M|)$, then $M \times N$ is the unique minimum \mathcal{F} -cover.*

Proof. Suppose not, and let G be a minimum \mathcal{F} -cover. Then G is simple, and has a subgroup N with index at most $|M|$. Now G acts faithfully on the cosets of N , and so it

is embeddable in the symmetric group of degree $|M|$, with N as the point stabilizer. So N is embeddable in the symmetric group of degree $|M| - 1$, and $|N| \leq (|M| - 1)!$. \square

We conclude this section by asking: Is it possible for a set of two nonabelian finite simple groups to have two nonisomorphic minimum covers?

In light of Theorem 6, if M and N are nonabelian finite simple groups and $\mathcal{F} = \{M, N\}$ has two nonisomorphic minimum covers, then one of the following must hold:

- (a) there are two simple groups of the same order (smaller than $|M| \cdot |N|$) which are minimum \mathcal{F} -covers; or
- (b) there is a simple group of order $|M| \cdot |N|$ which is a minimum \mathcal{F} -cover.

For (a), with the Classification of Finite Simple Groups, the only pairs of nonisomorphic finite simple groups of the same order are $\{\text{PSL}_3(4), A_8\}$ and $\{\text{PSP}_{2m}(q), \text{P}\Omega_{2m+1}(q)\}$ for $m \geq 3$ and q odd. Using the ATLAS [5], we can show that the first pair are not both minimal covers of any pair of simple groups. We suspect that there are no examples for the second pair either.

For (b), the following question arises which is of independent interest.

Question 5.5. Classify the triples (M, N, G) of nonabelian finite simple groups such that $|M| \cdot |N| = |G|$.

If G is a finite simple group that has a sharply t -transitive action of degree n , then $|A_{n-t}| \cdot |G| = |A_n|$. The following result addresses this special case.

Proposition 5.6. *Let G be a finite simple group with a sharply t -transitive action of degree n . Then A_n is a minimum cover of $\{A_{n-t}, G\}$ if and only if one of the following holds*

- (a) $G = \text{PSL}_2(2^f)$ and $(n, t) = (2^f + 1, 3)$ where $f \geq 3$
- (b) $G = M_{12}$ and $(n, t) = (12, 5)$

Proof. First assume that $t = 1$, so G acts regularly on n points. Since G has a faithful action on strictly fewer points, G embeds in A_{n-1} . In particular, A_{n-1} is a minimum cover of $\{A_{n-1}, G\}$.

Now assume that $t > 1$. Consulting [3, Theorem 4.11], for example, the only sharply t -transitive actions of a finite simple group G are:

1. $G = \text{PSL}_2(2^f)$ and $(n, t) = (2^f + 1, 3)$ where $f \geq 3$
2. $G = M_{11}$ and $(n, t) = (11, 4)$
3. $G = M_{12}$ and $(n, t) = (12, 5)$

It is straightforward to rule out Case 2: M_{23} is a cover of $\{A_7, M_{11}\}$ and $|M_{23}| = 10200960 < 19958400 = |A_{11}|$. For Case 3, it is also easy to verify (in MAGMA [2], say) that A_{12} is a minimum cover of $\{A_7, M_{12}\}$. (For comparison with Case 2, while M_{24} is a cover of $\{A_7, M_{12}\}$ we have $|M_{24}| = 244823040 > 239500800 = |A_{12}|$.)

Case 1 remains. Fix $f \geq 3$, write $q = 2^f$ and let $G = \text{PSL}_2(q)$. We claim that A_{q+1} is the smallest simple group to embed A_{q-2} and G . If $f \in \{3, 4\}$, then this is easily verified in MAGMA [2], so let us assume that $f \geq 5$. Let H be a simple group embedding both A_{q-1} and G . Since H embeds $\text{PSL}_2(q)$, if $H = A_d$, then $d \geq q + 1$ (see [17, Theorem 5.2.2]). Since H embeds A_{q-2} with $q - 2 \geq 30$, we deduce that H is not a sporadic group (see [17, Theorem 5.2.9]). Now assume that H is a classical group. Since H embeds A_{q-2} , the dimension of the natural module for H must be at least $q - 4$ (see [17, Theorem 5.3.7]). Consulting the order formulae for these groups, it is easy to see that this implies that $|H| \geq 2^{(q-4)(q-6)/2}$. This means

$$\log_2 |H| \geq \frac{1}{2}(q - 4)(q - 6) \geq q \log_2(q + 1) \geq \log_2 |A_{q+1}|.$$

Finally assume that H is an exceptional group of Lie type. In light of the previous cases, consulting the possible maximal subgroups of H [19, Theorem 8], we see that H does not embed A_{q-2} with $q - 2 \geq 30$. Therefore, in all cases, $|H| \geq |A_{q+1}|$, so A_{q+1} is a minimum cover of $\{A_{q-2}, G\}$, as claimed. \square

Corollary 5.7. *A set of two nonabelian finite simple groups can have two nonisomorphic minimum covers. For example,*

- (a) A_{2^f+1} and $A_{2^f-2} \times \text{PSL}_2(2^f)$ are minimum covers of $\{A_{2^f-2}, \text{PSL}_2(2^f)\}$ whenever $f \geq 3$;
- (b) A_{12} and $A_7 \times M_{12}$ are minimum covers of $\{A_7, M_{12}\}$.

Remark 5.8. In an earlier version of the paper, we asked whether any prime divisor of the order of a minimum cover of a set of finite groups must divide the order of one of the groups in the set. However, this is false, since M_{23} is the unique minimum cover of $\{\text{PSL}_3(4), A_8\}$, two simple groups of the same order 20160 (this can be seen from the ATLAS [5]), but $|M_{23}|$ is divisible by 11 and 23, neither of which divides 20160.

6. Groups of prime-power order

In this section we examine sets of p -groups, for p prime.

Remark 6.1. It is well known that the only groups of order 8 are $\mathbb{Z}_8, \mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2, D_8, Q_8$. So the only 4-cover groups of order 8 are $\mathbb{Z}_4 \times \mathbb{Z}_2$ and D_8 . Similarly there are five groups of order p^3 for an odd prime p , which are $\mathbb{Z}_{p^3}, \mathbb{Z}_{p^2} \times \mathbb{Z}_p, \mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p$, and the two non-abelian groups

$$G_p = \left\{ \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix} : a \in 1 + p\mathbb{Z}_{p^2}, b \in \mathbb{Z}_{p^2} \right\},$$

$$\text{and Heis}(p) = \left\{ \begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} : a, b, c \in \mathbb{Z}_p \right\}$$

with exponent p^2 and p respectively. This implies the only p^2 -cover groups of order p^3 are G_p and $\mathbb{Z}_p \times \mathbb{Z}_{p^2}$

Proposition 6.2. *Let p be a prime number.*

- (a) *If $p = 2$, then the only minimal 4-cover groups are $\mathbb{Z}_4 \times \mathbb{Z}_2$ and D_8 ;*
- (b) *If $p > 2$, then the only minimal p^2 -cover groups are $\mathbb{Z}_{p^2} \times \mathbb{Z}_p$ and G_p .*

Proof. Let G be a minimal p^2 -cover group. Then G is a p -group by Proposition 2.5. It is well known that \mathbb{Z}_{p^2} and $\mathbb{Z}_p \times \mathbb{Z}_p$ are the only two groups of order p^2 up to isomorphism. Let H and $K = \langle a \rangle$ be subgroups of G which are isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$ and \mathbb{Z}_{p^2} . Also G contains a normal subgroup N of order p^2 . If N is cyclic then there is an element $g \in H \setminus N$. Then $\langle g \rangle N$ is a p^2 -cover of order p^3 and the proof is complete by Remark 6.1. So assume $N \cong \mathbb{Z}_p \times \mathbb{Z}_p$. First assume $|N \cap K| = 1$. Then NK is a p^2 -cover of order p^4 . Hence $G = NK$. Since $N_G(N)/C_G(N)$ is isomorphic to a subgroup of $\text{Aut}(N) \cong \text{GL}(2, p)$, so $|N_G(N)/C_G(N)| \mid p$. Thus $a^p \in C_G(N)$. So $a^p \in Z(G)$. Also $N \cap Z(G)$ is non-trivial so there is an element $g \in Z(G) \cap N$ of order p . Hence $\langle g \rangle K$ is a p^2 -cover of order p^3 which is a contradiction by minimality of G . Hence $|N \cap K| = p$. So $|NK| = p^3$ and NK is a p^2 -cover. By Remark 6.1, the proof is complete. \square

In the next case, we have the following.

Theorem 6.3.

- (a) *There are two minimum 2^3 -cover groups, both of order 2^5 .*
- (b) *For prime $p > 2$, there is no p^3 -cover of order p^5 , but there is one of order p^6 ; so a minimum p^3 -cover has order p^6 .*

Proof. Part (a) can be proved by computer: we used the computer algebra system GAP [10]. The two 8-covers are the groups `SmallGroup(32, 40)` and `SmallGroup(32, 43)` in the GAP library. (There is no 8-cover of order 16. For suppose that G was an 8-cover of order 16. Then G contains subgroups $A \cong \mathbb{Z}_8$ and $B \cong (\mathbb{Z}_2)^3$; clearly $|A \cap B| \leq 2$, so $|AB| = |A| \cdot |B| / |A \cap B| \geq 32$.)

For (b), the case $p = 3$ can also be shown using GAP. For $p \geq 5$, we proceed as follows. Let G be a p^3 -cover of order p^5 . Then G has nilpotency class at most 4, so smaller than p ; hence G is a regular p -group [13, p. 183]. Now [13, Theorem 12.4.5] shows that the

elements of order p in G , together with the identity, form a subgroup H of G . Now since G contains both the elementary abelian group $(\mathbb{Z}_p)^3$ of order p^3 and the nonabelian group $\text{Heis}(p)$ of order p^3 and exponent p ; so the subgroup H must satisfy $|H| > p^3$, so $|H| \geq p^4$. Now G must also contain the cyclic group $K = \mathbb{Z}_{p^3}$, and $|H \cap K| \leq p$, so $|HK| \geq p^4 \cdot p^3/p = p^6$. So no p^3 -cover of order p^5 can exist.

For the example, we start with the group $E = \text{Heis}(p)$ of order p^3 and exponent p :

$$E = \langle x, y, z \mid x^p = y^p = z^p = [x, z] = [y, z] = 1, [x, y] = z \rangle.$$

It is easy to see that E has an automorphism α such that $\alpha(x) = x$ and $\alpha(y) = xy$.

Now we use the following result [13, Theorem 15.3.1]:

Lemma 6.4. *Let N be a finite group, $\alpha \in \text{Aut}(N)$, and $m \in \mathbb{N}$. Then the following assertions are equivalent:*

- (a) *There exists a finite group H such that $N \trianglelefteq H$, $H/N = \langle hN \rangle = \mathbb{Z}_m$ and $\alpha(x) = h x h^{-1}$ for all $x \in N$;*
- (b) *There exists $n \in N$ such that $\alpha(n) = n$ and $\alpha^m(a) = n a n^{-1}$ for all $a \in N$.*

We apply the lemma with $m = p^2$ and $n = z$, giving a non-split extension H of E such that $H/E = \langle aE \rangle \cong \mathbb{Z}_{p^2}$ with $a^{p^2} = z$ and $e^a = \alpha(e)$ for all $e \in E$. For $b := a^p y^{-1} \in H$ we compute $b^x = a^p [x, y] y^{-1} = bz = ba^{p^2} = b^{1+p}$. Hence, H has subgroups $E \cong \text{Heis}(p)$, $\langle a \rangle \cong \mathbb{Z}_{p^3}$, $\langle a^p, x \rangle \cong \mathbb{Z}_{p^2} \times \mathbb{Z}_p$ and $\langle b, x \rangle \cong G_p$.

Finally, set $G = H \times \mathbb{Z}_p$; it is clear that G also contains $(\mathbb{Z}_p)^3$. \square

In addition, computation with GAP shows that, for $p = 3$, there are many examples of p^3 -covers of order p^6 .

Next we show that there are infinitely many minimal 2^3 -covers, and that these may be taken to be strongly minimal; indeed, having no proper subquotient which is a 2^3 -cover.

For the proof we use the *semidihedral group*

$$SD_{2^n} = \langle a, b \mid a^{2^{n-1}} = b^2 = 1, b^{-1} a b = a^{2^{n-2}-1} \rangle$$

of order 2^n , for $n \geq 4$. We make a couple of observations about this group.

- Its center is cyclic of order 2, generated by $a^{2^{n-2}}$.
- $(ab)^2 = a \cdot a^{2^{n-2}-1} = z$, so ab has order 4.
- It is a 2-generated 2-group, and so has three maximal subgroups of index 2. These are $\langle a \rangle$ (cyclic), $\langle a^2, b \rangle$ (dihedral) and $\langle a^2, ab \rangle$ (generalized quaternion).

Theorem 6.5. *Let $n \geq 4$, $A = SD_{2^n}$, and $C = \langle c \rangle = \mathbb{Z}_2$. Then the group $A \times C$ of order 2^{n+1} is a minimal and co-minimal 8-cover.*

Proof. First we show that all groups of order 8 are subgroups of G . We have $\langle a^{2^{n-4}} \rangle \cong \mathbb{Z}_8$, $\langle a^{2^{n-3}}, c \rangle \cong \mathbb{Z}_4 \times \mathbb{Z}_2$, $\langle a^{2^{n-2}}, b, c \rangle \cong (\mathbb{Z}_2)^3$, $\langle a^{2^{n-3}}, b \rangle \cong D_8$, and $\langle a^{2^{n-3}}, ab \rangle \cong Q_8$.

Now we show that no proper subgroup of G is an 8-cover. It suffices to consider a maximal subgroup H . Let ϕ be the projection of $A \times C$ onto the first factor. If $c \notin H$, then the restriction of ϕ to H is an isomorphism to A , so H is semidihedral. If $c \in H$, then $H = K \times C$ where K is a maximal subgroup of A , and so is cyclic, dihedral or generalized quaternion. None of these groups is an 8-cover.

Finally we show that no proper quotient of G is a 2-cover. Again it suffices to consider maximal quotients G/N , where N is a normal subgroup of G . Then $N \leq Z(G) = \langle z, c \rangle$; so $N = \langle z \rangle$, $\langle c \rangle$ or $\langle zc \rangle$. If $z \notin N$, then the restriction of the projection $G \rightarrow G/N$ to A is an isomorphism, so G/N is semidihedral. Otherwise $G/N = (A/\langle z \rangle) \times C \cong D_{2^{n-1}} \times \mathbb{Z}_2$. Again none of these groups, or any of their quotients, is an 8-cover. \square

Remark 6.6. A slightly more elaborate argument shows that in fact no quotient of a subgroup of G , apart from G itself, is an 8-cover.

The numbers of 8-covers of order 2^n are given below, together with the numbers of minimal and strongly minimal 8-covers, for $5 \leq n \leq 8$.

Order	32	64	128	256
Number of groups	51	267	2328	56092
Number of 8-covers	2	45	745	14798
Number of minimal 8-covers	2	18	85	969
Number of strongly minimal 8-covers	2	14	3	7

Remark 6.7. A minimum 16-cover has order 2^8 , and `SmallGroup(256,384)` in the GAP library is an example.

In contrast to the upper bound for the order of a minimum p^m -cover (the order of the Sylow subgroup of S_{p^m}), we give a lower bound $p^{\Omega(m^2)}$, which is probably rather weak.

We begin with a brief note. The fraction $|\text{GL}(n, p)|/p^{n^2}$ is the probability that an $n \times n$ matrix over the field of order p is invertible. It can be written as

$$\prod_{i=1}^n (1 - p^{-i}).$$

The theory of infinite products shows that, as $n \rightarrow \infty$ with p fixed, it decreases to a positive limit $\theta(p)$, which is an evaluation of a Jacobi theta-function. It is easily seen that $\theta(p)$ is an increasing function of p . The value of $\theta(2)$ is $0.2887\dots$. So the probability that an $n \times n$ matrix over the p -element field is invertible is at least $\theta(2)$ for any n and p .

Lemma 6.8. *Let G be a group of order p^n . Then the number of n -tuples of elements of G , which generate G is at least cp^{n^2} , where c is an absolute constant.*

Proof. Let $|G/\Phi(G)| = p^k$. By Burnside’s basis theorem, a k -tuple g_1, \dots, g_k generates G if and only if the images of g_1, \dots, g_k in $G/\Phi(G)$ form a basis for this quotient, which is isomorphic to a k -dimensional vector space over the p -element field. The number of such bases is the order of $\text{GL}(k, p)$, which as noted is at least cp^{k^2} , for an absolute constant c . For each basis element, there are p^{n-k} elements of the corresponding coset of $\Phi(G)$ in G . Also, we can complete the n -tuple by choosing arbitrary elements of G , each in p^n ways. So the number of n -tuples is

$$|\text{GL}(k, p)| \cdot p^{k(n-k)} \cdot p^{(n-k)n} \geq cp^{n^2},$$

as required. \square

Theorem 6.9. *The order of a minimum p^n -cover is at least $p^{(2/27+o(1))n^2}$ as $n \rightarrow \infty$ (for fixed p).*

Proof. Suppose that G is a minimum p^n -cover, of order p^N . There are p^{Nn} n -tuples of elements of G ; among them are generating tuples for all groups of order p^n . By Lemma 6.8, each group of order p^n has at least cp^{n^2} generating n -tuples. So the number of groups of order p^n is at most

$$p^{nN}/(cp^{n^2}) = c^{-1}p^{n(N-n)}.$$

However, it was proved by Higman and Sims [14,25] (see also [1]) that the number of different groups of order p^n is $p^{(2/27+o(1))n^3}$. So

$$c^{-1}p^{n(N-n)} \geq p^{(2/27+o(1))n^3},$$

from which we find that $N \geq (2/27 + o(1))n^2$. \square

Question 6.10. Find better bounds for the order of a minimum p^n -cover. In particular, is there an upper bound of the form $p^{F(n)}$, where F is independent of p ?

In the next section, we will find the smallest abelian group which contains all abelian groups of order p^n ; its order is roughly $p^{n \log n}$.

7. Cyclic, abelian and nilpotent groups

The observation that A_5 is a minimum cover for $\{\mathbb{Z}_3, (\mathbb{Z}_2)^2, \mathbb{Z}_5\}$ shows that it is not true that, if all groups in \mathcal{F} are abelian, nilpotent, or soluble, then every minimum \mathcal{F} -cover has the same property. Moreover, $\{\mathbb{Z}_2, \mathbb{Z}_3\}$ has two minimum covers, \mathbb{Z}_6 and S_3 . So the best we can hope is that, if all groups in \mathcal{F} have a certain property, then at least one minimum \mathcal{F} -cover has this property. This is the case for cyclic groups:

Theorem 7.1. *Let n_1, \dots, n_k be positive integers and $N = \text{lcm}(n_1, \dots, n_k)$. Then \mathbb{Z}_N is a minimum cover for $\mathcal{F} = \{\mathbb{Z}_{n_1}, \dots, \mathbb{Z}_{n_k}\}$.*

Proof. Clearly \mathbb{Z}_N is an \mathcal{F} -cover, and by Proposition 2.1 it is minimum. \square

Perhaps the next simplest example of Question B is one which we have not been able to settle. The case where $|\mathcal{F}| = 2$ is easy but even $|\mathcal{F}| = 3$ seems hard to us.

Question 7.2. Let \mathcal{F} be a set of abelian p -groups, for some prime p . Is there a minimum \mathcal{F} -cover which is an abelian p -group?

Theorem 7.3. *Suppose that \mathcal{F} is a finite set of finite nilpotent groups. Then there is a minimum \mathcal{F} -cover which is nilpotent. If Question 7.2 has an affirmative answer, then the same holds with “abelian” replacing “nilpotent”.*

Proof. Let $\mathcal{F} = \{F_1, \dots, F_r\}$ be a finite set of finite nilpotent groups, and let G be a minimum \mathcal{F} -cover. For each prime p , let $F_i(p)$ be the Sylow p -subgroup of F_i , and let $\mathcal{F}(p) = \{F_1(p), \dots, F_r(p)\}$. Let $H(p)$ be a minimum $\mathcal{F}(p)$ -cover, and note that $H(p)$ is a p -group, by Proposition 2.5. Let H be the direct product of the groups $H(p)$. Now a Sylow p -subgroup $G(p)$ of G is an $\mathcal{F}(p)$ -cover, so $|G(p)| \geq |H(p)|$, and thus $|G| \geq |H|$. But since each group in \mathcal{F} is the direct product of its Sylow subgroups, it is embeddable in H , and thus by minimality $|G| = |H|$. So H is a nilpotent cover of \mathcal{F} with smallest possible order.

Now suppose that all the groups in \mathcal{F} are abelian, and that Question 7.2 has an affirmative answer. The same argument then applies, using the fact that a nilpotent group is abelian if and only if all its Sylow subgroups are. \square

What is the order of the minimum cover? The theorem shows that it is enough to find the order of a minimum cover of a set of p -groups. We can answer this question in the case of abelian p -groups, again assuming that Question 7.2 has an affirmative answer.

Suppose that $\mathcal{F} = \{F_1, \dots, F_r\}$ is a set of abelian p -groups. We can write each one in canonical form:

$$F_i = \mathbb{Z}_{p^{a(i,1)}} \times \dots \times \mathbb{Z}_{p^{a(i,k)}},$$

where $a(i, 1) \geq a(i, 2) \geq \dots \geq a(i, k)$; by adding extra zero terms if necessary we can assume that the value of k is the same for each group. Let

$$c(j) = \max\{a(1, j), a(2, j), \dots, a(r, j)\}$$

for $j = 1, \dots, k$. We claim that

$$c(1) \geq c(2) \geq \dots \geq c(k).$$

For suppose that $c(j + 1) = a(i, j + 1)$. Then $c(j) \geq a(i, j) \geq a(i, j + 1) = c(j + 1)$.

Let

$$P = \mathbb{Z}_{p^{c(1)}} \times \cdots \times \mathbb{Z}_{p^{c(k)}}.$$

The above claim shows that this is the canonical form for P .

Proposition 7.4. *With the above notation, P is the smallest abelian \mathcal{F} -cover.*

The proof depends on the following lemma:

Lemma 7.5. *Let*

$$A = \mathbb{Z}_{p^{a(1)}} \times \cdots \times \mathbb{Z}_{p^{a(k)}} \text{ and } B = \mathbb{Z}_{p^{b(1)}} \times \cdots \times \mathbb{Z}_{p^{b(k)}}$$

be abelian p -groups in canonical form. Then B is embeddable in A if and only if $b(j) \leq a(j)$ for $j = 1, \dots, k$.

Proof. Suppose that the inequalities hold. Then for each j we can choose a subgroup $\mathbb{Z}_{p^{b(j)}}$ of $\mathbb{Z}_{p^{a(j)}}$; the direct product of these subgroups is isomorphic to B .

Conversely, suppose that B is embeddable in A . Then B contains a subgroup $(\mathbb{Z}_{p^{b(j)}})^j$; in order to embed this in A , we require that at least j of $a(1), \dots, a(k)$ are greater than or equal to $b(j)$ for each j . Since the a are non-increasing, this requires $a(j) \geq b(j)$. \square

Now, to complete the proof of Proposition 7.4, note that in the notation before the proposition, P embeds all the F_i if and only if $c(j) \geq a(i, j)$ for all i . So P is the smallest abelian \mathcal{F} -cover. \square

We can use this result to find the smallest abelian group containing every abelian group of order p^n .

Define a function f by the rule

$$f(n) = \sum_{k=1}^n \lfloor n/k \rfloor.$$

Corollary 7.6. *Let \mathcal{F} be the set of all abelian groups of order p^n . There is a unique smallest abelian \mathcal{F} -cover; its order is $p^{f(n)}$. If the answer to Question 7.2 is affirmative, it is a minimum \mathcal{F} -cover.*

Proof. In the notation introduced before Proposition 7.4, we have $c(k) = \lfloor n/k \rfloor$ for $k = 1, \dots, n$. For if the factors in the canonical decomposition of an abelian group of order p^n have orders $p^{a(1)}, p^{a(2)}, \dots$, then

$$ka(k) \leq a(1) + \cdots + a(k) \leq n,$$

so $a(k) \leq \lfloor n/k \rfloor$; but there is a group of order p^n with k invariant factors which are of nearly equal orders (that is, orders $p^{\lfloor n/k \rfloor}$ or $p^{\lceil n/k \rceil}$); the k th of these in non-increasing order has order $p^{\lfloor n/k \rfloor}$.

So the smallest abelian group covering \mathcal{F} has order $p^{\sum \lfloor n/k \rfloor} = p^{f(n)}$, as required. \square

For $p^n = 2^2$ and 2^3 , this gives respectively 8 and 32 for the smallest abelian group containing all abelian groups of order p^n . We have seen that these are also the orders of minimal covers for all groups of these orders. Furthermore, no smaller group can cover all abelian groups of these orders, by the proof of Theorem 6.3. So Question 7.2 has an affirmative answer in these cases.

We note that the order of the smallest abelian cover of the class of abelian groups of order p^n is roughly $p^{n \log n}$, which can be contrasted with the lower bound of p^{cn^2} for a group covering every group of order p^n . More precisely, $f(n) = n(\log n + 2\gamma - 1) + O(\sqrt{n})$, where γ is the Euler–Mascheroni constant (Dirichlet [7]).

The sequence of values of the function f is sequence A006218 in the On-Line Encyclopedia of Integer Sequences [24]. This gives many interpretations of the sequence, but the one given here appears to be new.

Combining the result for the prime factors of an integer n , we obtain the following result.

Theorem 7.7. *Let $n = p_1^{m_1} \cdots p_r^{m_r}$, where p_1, \dots, p_r are distinct primes. Then the order of the smallest abelian group which embeds all abelian groups of order n is*

$$A(n) = p_1^{f(m_1)} \cdots p_r^{f(m_r)},$$

where $f(m) = \sum_{k=1}^m \lfloor m/k \rfloor$.

Note that $A(n) = n$ if n is squarefree, while $A(n)$ is roughly $n^{\log \log n}$ if n is a power of 2. The values of $A(n)$ are not those given by sequence A102361 in the OEIS [24], although the first fifteen terms agree (and these sequences agree at all fourth-power-free integers).

We could ask whether similar results exist for soluble groups. But there is an easy example to show that a set of soluble groups may have no soluble minimum cover:

Example 7.8. Let A be the alternating group A_4 and B the dihedral group D_{10} of order 10. Then $\text{lcm}(|A|, |B|) = 60$, and both groups are embeddable in A_5 , so A_5 is a minimum cover. There is no other cover of order 60. For such a group G would act on the five cosets of A ; it is easily seen that either the action is faithful (whence $G \cong A_5$) or A lies in the kernel (in which case $G \cong A_4 \times \mathbb{Z}_5$); but in the second case G does not embed the dihedral group.

8. Further directions

We conclude by highlighting some further directions to be pursued, in addition to the various questions already posed in the paper.

First, the main definitions of this paper can be naturally dualized. Let \mathcal{F} be a set of finite groups. A group G is a *dual \mathcal{F} -cover* if every group in \mathcal{F} is isomorphic to a quotient of G . We say that a dual \mathcal{F} -cover G is

- *minimal* if no proper quotient of G is a dual \mathcal{F} -cover;
- *co-minimal* if no proper subgroup of G is a dual \mathcal{F} -cover;
- *strongly minimal* if it is both minimal and co-minimal;
- *minimum* if no dual \mathcal{F} -cover has smaller order.

Note that a minimum dual cover is strongly minimal.

We have not investigated this concept except to note that, in the class of abelian groups, subgroups and quotients coincide (because of duality for abelian groups), so, for example, an abelian group is a minimum dual cover of a class of abelian groups if and only if it is a minimum cover.

Question 8.1. Investigate dual covers along the lines we have followed for covers.

Next, the work in Section 7 leads to the following question.

Question 8.2. For which classes \mathcal{X} of groups, closed under the taking of subgroups and direct product, is it true that, if \mathcal{F} is a finite set of \mathcal{X} -groups, then there is a minimum \mathcal{F} -cover which is an \mathcal{X} -group?

As we have seen, this is true for cyclic groups and for nilpotent groups, but it is false for soluble groups (see Example 7.8). We have been unable to resolve this question for abelian groups.

Finally, we give the answer to a question asked in an earlier version of the paper. The question asked: For which groups G is it the case that G is not a minimal cover of the set of its proper subgroups (equivalently, the set of its maximal subgroups)?

Theorem 8.3. *If G is not a minimal cover of the set of its proper subgroups, then G is a p -group for some prime p , and all its maximal subgroups are isomorphic.*

Proof. If G is not of prime-power order, then such a cover must contain all the Sylow subgroups of G , and so must be at least as large as G . If G is a p -group, then all maximal subgroups have index p ; so, if two are nonisomorphic, then again a cover must be at least as large as G . \square

Remark 8.4. The 2-groups with the property of the theorem were determined by Čepulić [4].

Data availability

No data were used for the research described in the article.

References

- [1] Simon R. Blackburn, Peter M. Neumann, Geetha Venkataraman, *Enumeration of Finite Groups*, Cambridge University Press, Cambridge, 2007.
- [2] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I. The user language, *J. Symb. Comput.* 24 (1997) 235–265.
- [3] Peter J. Cameron, *Permutation Groups*, London Mathematical Society Student Texts, vol. 45, Cambridge University Press, 1999.
- [4] V. Čepulić, On finite 2-groups all of whose subgroups are mutually isomorphic, *Sci. China Ser. A: Math.* 52 (2009) 254–260.
- [5] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker, R.A. Wilson, *ATLAS of Finite Groups*, Oxford University Press, Oxford, 1985.
- [6] Heiko Dietrich, Melissa Lee, Tomasz Popiel, The maximal subgroups of the Monster, arXiv:2304.14646.
- [7] P.G.L. Dirichlet, *Werke*, in: L. Kronecker (Ed.), Vol. ii, 1849, pp. 49–66, Reprinted Wentworth Press, 2018.
- [8] Klaus Doerk, Trevor Hawkes, *Finite Soluble Groups*, Walter de Gruyter, Berlin, 1992.
- [9] L.F. Fletcher, B. Stellmacher, W.B. Stewart, Endliche gruppen, die kein element der ordnung 6 enthalten, *Q. J. Math. Oxf. (2)* 28 (1977) 143–154.
- [10] The GAP Group, GAP – groups, algorithms, and programming, version 4.12.2, <https://www.gap-system.org>, 2022.
- [11] L.M. Gordon, Finite simple groups with no elements of order six, *Bull. Aust. Math. Soc.* 17 (1977) 235–246.
- [12] Robert Guralnick, Tim Penttila, Cheryl Praeger, Jan Saxl, Linear groups with orders having certain large prime divisors, *Proc. Lond. Math. Soc. (3)* 78 (1999) 167–214.
- [13] Marshall Hall Jr., *The Theory of Groups*, MacMillan, New York, 1959.
- [14] Graham Higman, Enumerating p -groups. I: inequalities, *Proc. Lond. Math. Soc. (3)* 10 (1960) 24–30.
- [15] Petra E. Holmes, Robert A. Wilson, $PSL_2(59)$ is a subgroup of the Monster, *J. Lond. Math. Soc. (2)* 69 (2004) 141–152.
- [16] Petra E. Holmes, Robert A. Wilson, A new maximal subgroup of the Monster, *J. Algebra* 251 (2008) 435–447.
- [17] Peter Kleidman, Martin Liebeck, *The Subgroup Structure of the Finite Classical Groups*, London Mathematical Society Lecture Note Series, vol. 129, Cambridge University Press, 1990.
- [18] Martin W. Liebeck, Jan Saxl, On point stabilizers in primitive permutation groups, *Commun. Algebra* 19 (1991) 2777–2786.
- [19] Martin W. Liebeck, Gary M. Seitz, A survey of maximal subgroups of exceptional groups of Lie type, in: *Groups, Combinatorics & Geometry*, Durham, 2001, World Sci. Publ., River Edge, NJ, 2003.
- [20] G. Malle, D. Testerman, *Linear Algebraic Groups and Finite Groups of Lie Type*, Graduate Studies in Advanced Mathematics, vol. 133, Cambridge University Press, 2011.
- [21] Donald McCarthy, Sylow’s theorem is a sharp partial converse to Lagrange’s theorem, *Math. Z.* 113 (1970) 383–384.
- [22] Donatella Merlini, Renzo Sprugnoli, M. Cecilia Verri, The Cauchy numbers, *Discrete Math.* 306 (2006) 1906–1920.
- [23] N.D. Podufalov, Finite simple groups without elements of sixth order, *Algebra Log.* 16 (1977) 133–135.
- [24] The on-line encyclopedia of integer sequences, <https://oeis.org>.
- [25] Charles C. Sims, Enumerating p -groups, *Proc. Lond. Math. Soc. (3)* 15 (1965) 151–166.
- [26] M. Suzuki, *Group Theory I*, Grundlehren der Math. Wissenschaften, vol. 247, Springer-Verlag, Berlin, 2014.