

Levels and trends of PBDEs and HBCDs in the global environment

Law, R.J.; Covaci, A.; Harrad, S.; Herzke, D.; Abdallah, M.A.-E.; Fernie, K.; Toms, L.M.L.; Takigami, H.

DOI:

[10.1016/j.envint.2014.01.006](https://doi.org/10.1016/j.envint.2014.01.006)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Law, RJ, Covaci, A, Harrad, S, Herzke, D, Abdallah, MA-E, Fernie, K, Toms, LML & Takigami, H 2014, 'Levels and trends of PBDEs and HBCDs in the global environment: Status at the end of 2012', *Environment International*, vol. 65, pp. 147-158. <https://doi.org/10.1016/j.envint.2014.01.006>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Document subject to a Creative Commons Attribution Non-Commercial No Derivatives license

Validated December 2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Levels and trends of PBDEs and HBCDs in the global environment: status at the end of 2012

Robin J. Law ^{a,*}, Adrian Covaci ^b, Stuart Harrad ^c, Dorte Herzke ^d, Mohamed A.-E. Abdallah ^e, Kim Fernie ^f,
Leisa-Maree L. Toms ^g, Hidetaka Takigami ^h

^a *The Centre for Environment, Fisheries and Aquaculture Science, Cefas Lowestoft Laboratory, Pakefield Road, Lowestoft, Suffolk NR33 0HT, UK*

^b *Toxicological Centre, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk-Antwerp, Belgium*

^c *School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK*

^d *Norwegian Institute for Air Research, FRAM – High North Research Centre for Climate and the Environment, H. Johansens gate 14, 9296 Tromsø, Norway*

^e *Department of Analytical Chemistry, Faculty of Pharmacy, Assiut University, 71526 Assiut, Egypt*

^f *Ecotoxicology and Wildlife Health Division, Wildlife and Landscape Science Directorate, Science and Technology Branch, Environment Canada, 867 Lakeshore Road, Burlington L7R 4A6, Ontario, Canada*

^g *School of Clinical Sciences and Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Australia*

^h *Centre for Material cycles and Waste Management Research, National Institute for Environmental Studies (NIES), 16-2 Onogawa, Tsukuba, 305-8506, Ibaraki, Japan*

Abstract

In this paper, we have compiled and reviewed the most recent literature, published from January 2010 to December 2012, relating to the human exposure, environmental distribution, behaviour, fate and concentration time trends of polybrominated diphenyl ether (PBDE) and hexabromocyclododecane (HBCD) flame retardants, in order to establish their current trends and priorities for future study. Due to the large volume of literature included, we have provided full detail of the reviewed studies as Electronic Supplementary Information and here summarise the most relevant findings. Decreasing time trends for penta-mix PBDE congeners were seen for soils in northern Europe, sewage sludge in Sweden and the USA, carp from a US river, trout from three of the Great Lakes and in Arctic and UK marine mammals and many birds, but increasing time trends continue in polar bears and some birds at high trophic levels in northern Europe. This may be partially a result of the time delay inherent in long-range atmospheric transport processes. In general, concentrations of BDE209 (the major component of the deca-mix PBDE product) are continuing to increase. Of major concern is the possible/likely debromination of the large reservoir of BDE209 in soils and sediments worldwide, to yield lower brominated congeners which are both more mobile and more toxic, and we have compiled the most recent evidence for the occurrence of this degradation process. Numerous studies reported here reinforce the importance of this future concern. Time trends for HBCDs are mixed, with both increases and decreases evident in different matrices and locations and, notably, with increasing occurrence in birds of prey.

Keywords: PBDEs, HBCDs, distribution, temporal trends, BDE209 debromination

1. Introduction

Since brominated flame retardants (BFRs) were first detected in environmental samples in Sweden in 1979-1981 (Andersson and Blomkvist, 1981) interest in their provenance, environmental distribution and behaviour, fate and effects has grown enormously. A large number of reviews have attempted to document this and to identify directions for future research in order to resolve unanswered questions relating to the significance of the polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecanes (HBCDs). The first to review this topic was de Wit (2002); amongst the most recent reviews describing environmental levels are de Wit et al. (2010), Dominguez et al. (2012) and Law and Herzke (2012), and analytical aspects have even more recently been reviewed by Dirtu et al. (2013) and Xu et al. (2013). Inputs due to e-waste recycling activities in Asia and their potential effects on the health of local populations have also become of increasing concern over this period (Schnoor, 2012; Zhang et al., 2012a) and it is clear that the international trade in e-waste continues despite regulation (Anon., 2012; BBC, 2010).

In this paper, we aim to compile and assess the most recent evidence (published from January 2010 to December 2012) in relation to the human exposure, environmental distribution, behaviour, fate and concentration time trends of the PBDEs and HBCDs, with a view to identifying research directions and priorities in relation to these well-established BFRs. In view of the sheer volume of the literature reviewed (over 200 papers), we have adopted a novel approach in which this paper provides a summary of the main data reviewed within each topic and details are provided as electronic supplementary information available on the journal website, in order to keep the main paper to a reasonable length. This paper is timely as its data summary bridges the divide between the heyday of the first tranche of flame retardants to be used extensively (the PBDEs and HBCDs) and the rise of their successors, the novel BFRs, which are themselves beginning to be addressed broadly in the literature currently (Covaci et al., 2011; Papachlimitzou et al., 2012; Law et al., 2013).

2. Human exposure

2.1 *Indoor air and dust*

The low vapour pressures of PBDEs and HBCDs mean that, in indoor environments, they partition preferentially to dust. Consequently, over the period of this review, the majority of studies of indoor contamination address human exposure via ingestion of settled dust, rather than via inhalation of indoor air. Two papers of note have addressed specifically the presence of PBDEs and HBCDs in indoor air over the period of this review. The first reports on the development and validation of a passive air sampler that by providing measurement of both vapour *and* particulate phase PBDEs and HBCDs in air, facilitates future monitoring of inhalation exposure to primarily particulate-associated BFRs, like BDE209 and other alternative FRs with similar physico-chemical properties (Abdallah and Harrad, 2010). The second study (Björklund et al., 2012) reported concentrations of BDEs (including BDE209) in air inside 33 buildings to be indistinguishable from those measured simultaneously in air ventilating the same buildings. The authors concluded that contaminated indoor air is an important source of BDEs to

outdoor air. A further study documented concentrations of tri- to hexa-BDEs in both indoor air and dust from 20 indoor locations (homes, offices and laboratories) in Toronto, Canada in 2006 (Zhang et al, 2011a). Although the difference was not significant, the concentrations of Σ BDE₇ in office air (average = 0.79 ng m⁻³) exceeded those in homes (average = 0.49 ng m⁻³). These indoor concentrations exceeded substantially those reported for Toronto outdoor air, also suggesting ventilation of indoor air to be a source to the external environment. For PBDEs with 5 or less bromine atoms, significant correlation between air and dust concentrations was observed, confirming the hypothesis that PBDEs in dust arise from gaseous deposition from air. The lack of such correlation for higher brominated BDEs suggests this process to be less important, with other processes such as abrasion of PBDE-containing materials possibly playing a greater role.

Several studies have emerged recently that report levels of PBDEs in dust from countries for which such information has been hitherto unavailable. These studies include reports of concentrations in dust from homes and cars in the Czech Republic (Kalachova et al., 2012), homes and computer suites in Poland (Król et al., 2012), homes in Romania (Dirtu and Covaci, 2010), and offices in South Africa (Kefeni and Okonkwo, 2012). Concentrations of PBDEs in dust from the Eastern European studies were 84 – 5,900 μ g Σ tri-deca BDEs kg⁻¹, 331-3,100 μ g Σ tri-hepta BDEs kg⁻¹ and 495 μ g Σ tri-deca BDEs kg⁻¹ (median) in Czech, Polish and Romanian homes respectively; thereby providing demonstrable evidence of dust ingestion as an important contributor to the exposure of the population in these countries. HBCDs were also found in Czech and Romanian house dust, at Σ HBCD concentrations of < 0.3 – 950 μ g kg⁻¹ (Kalachova et al., 2012) and 190 μ g kg⁻¹ (median) Dirtu and Covaci, 2010), respectively. These concentrations are slightly lower than those reported for countries like the UK and the USA, for which median Σ HBCD concentrations of 730 and 390 μ g kg⁻¹, respectively, have been reported previously (Abdallah et al., 2008). The study of Kefeni and Okwonko (2012), revealed median concentrations of BDE47 and BDE99 in office dust from Pretoria, South Africa to be – at 44 and 77 μ g kg⁻¹, respectively - in line with those detected in the UK (median values 23 and 65 μ g kg⁻¹ for BDE47 and BDE99 respectively; Harrad et al., 2008) and Japan (median values of 31 and 38 μ g kg⁻¹ for BDE47 and BDE99, respectively; Suzuki et al., 2006), but an order of magnitude lower than in the USA (median values of 978 and 1,760 for BDE47 and BDE99, respectively; Batterman et al., 2010). In contrast, levels of BDE209 in South African office dust – at an average of 53 μ g kg⁻¹ – were substantially lower than reported in other countries (average concentrations of 6,930, 30,000 and 2,400 μ g kg⁻¹ in office dust from the USA, UK and Japan, respectively). These results imply use of penta-mix PBDE but not deca-mix PBDE formulations in South Africa. Overall, recent data reinforce the position that concentrations in indoor dust (especially of the penta-PBDE congeners) are at least an order of magnitude higher than elsewhere. Likewise, recent evidence underlines that – probably due to the more stringent fire safety regulations in the UK – concentrations of BDE209 in indoor dust in the UK are on a par with those in North America and exceed substantially those elsewhere. Moreover, while regional and national variability exists and concentrations in developing and transition economies are often lower than those in developed regions, the burgeoning database serves to emphasise the ubiquitous indoor presence of BDEs and HBCDs.

2.2 Outdoor air

The growing realisation of the importance of indoor contamination as a source of human exposure has arguably shifted attention away from outdoor air. Notwithstanding this, there remains a strong imperative for studies of the levels of PBDEs and HBCDs in outdoor air, in order to understand better how these BFRs distribute through the environment and enter the food chain. A key concern about PBDEs (a characteristic of POPs generally) is their ability to achieve global distribution, even in locations remote from sources, such as the sparsely populated Tibetan Plateau in two recent studies (Wang et al., 2010; Xiao et al., 2012) and Antarctica (Dickhut et al., 2012).

New data on dust contamination with BDEs has also emerged that augments the global database with respect to levels in hitherto uncharacterised or poorly characterised microenvironment categories. These include: cars, offices, and classrooms in child daycare centres and primary schools. Another area for which new information has been generated is the spatial and temporal variability in BDE and HBCD concentrations in dust from the same room. Knowledge of such variability provides important insights into the role of putative sources, and of the uncertainty associated with exposure assessments based on single spot samples of dust taken at one point in time and space. A further potential influence on the concentrations of BDEs in indoor dust samples is the particle size fraction sampled and analysed. It is likely that this issue will form a focus for future research over the coming years, given factors such as the likely dependence of BDEs concentration on particle size and the relative bioavailability of BDEs associated with different particle size fractions. A further area likely to attract increasing attention in coming years is the potential for social inequity in exposure to BDEs and related BFRs. This stems from the hypothesis that lower-income households will contain a disproportionate number of older, deteriorating, or poorly-manufactured PBDE-treated furniture. One study reports worldwide maximum concentrations of BDE47 and BDE99 in dust from low-income homes in California (126,000 and 220,000 $\mu\text{g kg}^{-1}$, respectively) (Quirós-Alcalá et al, 2011). Also in California, Rose et al (2010) determined concentrations of BDEs in blood plasma from 100 children aged 24-60 months, and reported lower maternal education to be significantly associated with higher concentrations in children. Notwithstanding this evidence, much work remains if this hypothesis of social inequity in exposure is to be tested rigorously. For example, higher concentrations in children in the latter study were associated significantly with factors more likely related to higher socioeconomic status; viz higher consumption of pork and poultry (for penta-mix BDE congeners) and the presence in the home of new furniture and mattresses (for nona- and deca-BDE congeners).

The period covered by this review has seen two notable studies that have examined the relationship between measures of indoor contamination of BDEs and body burdens in individuals in contact with such contamination. In Denmark, Vorkamp et al (2011) studied 43 pregnant women reporting concentrations of some penta-mix BDE congeners in dust to be significantly correlated with those in both matched air and placenta samples, indicating indoor exposures may be an important pathway of exposure for these BDEs, but not BDE209. In Massachusetts, USA, Johnson et al (2010) reported similar correlation for penta-mix BDEs in house dust and blood serum of 12 male-female couples; with the low detection frequency of BDE209 in serum preventing evaluation of correlation between dust and serum

levels for this congener. Combined, these studies add weight to the hypothesis that indoor contamination can be an important driver of human body burdens of BFRs.

2.3 *Dietary exposure*

A number of food basket studies have been performed recently and the exposure of the general population to PBDEs and HBCDs has been estimated. In Sweden, fish and dairy products were the major contributors to PBDE intake (Tornkvist et al., 2011). In the same study, HBCD intake was primarily from fish (65%). In the USA, meat products are a major contributor to PBDE intake (Huwe and West, 2011). HBCD levels were low, consistent with the lesser use of this FR, with the highest levels in canned fish. Eggs are another group of food items which can have a high contribution to the daily intake of PBDEs. Levels of PBDEs and HBCD were high in home-produced eggs in the vicinity of e-waste recycling sites in China (Zheng et al., 2012). HBCD levels were lower than those of BDEs in this study. In comparison with a previous study of home-produced eggs from free range chickens in Belgium, levels of BDEs were greater than those of HBCD and BDE209 was the dominant congener (Covaci et al., 2009). Lower levels were also found in eggs from Canada (Rawn et al., 2011).

Most of the studies have indicated a rather diverse contribution of food items to the dietary exposure, ranging from meat, fish, canned products, and eggs. It also appears that diet becomes an important exposure route only when dust levels of BFRs and thus the contribution of dust to total exposure are low. The importance of each exposure route might be country-specific.

It is important to mention that a number of issues have only very recently been investigated, such as bioaccessibility of ingested food, changes in concentrations and patterns of BFRs during cooking processes (Domingo et al., 2011) and modeling of BFR dietary exposure (Trudell et al., 2011a,b). It is expected that they will continue to grow in importance in the next years.

2.4 *Levels in human milk and blood*

Overall, 7 papers were identified to report PBDEs and/or HBCD temporal trends in human matrices published since January 2010. Years of collection ranged from 1993 to 2009 with sampling matrices including human blood serum and human breast milk. Results were variable. For PBDEs, decreasing temporal trends or a peak followed by plateau were identified in Australia (Toms et al., 2012), Italy (Alivernini et al., 2011), Germany in Lower Saxony (Hoopmann et al., 2012) and Baden-Württemberg (Link et al., 2012), and in the Great Lakes Basin of the USA (Turyk et al., 2010). In contrast, PBDE concentrations increased over time in Guinea-Bissau (West Africa) (Linderholm et al., 2010) and Ghana (Asante et al., 2011) (Figure 1). HBCD concentrations, where reported, were also variable and no studies showed a temporal trend (Asante et al., 2011; Toms et al., 2012).

Points of interest in these studies were an increase in BDE209, a congener rarely reported in these papers, in children from Baden-Württemberg, Germany from 2002-2003 to 2008-2009 (Link et al., 2012); and a shift in congener distribution with an increase in the proportion of BDE153 seen in human samples compared to the previously ubiquitously dominant BDE47 (Linderholm et al., 2010; Turyk et al., 2010; Hoopmann et al., 2012). Reasons for this shift are suggested to be either differences in metabolism of BDE congeners, with BDE153 being more persistent, or to be related to changes in exposure due to the phase out of the penta-mix PBDE product, but not the deca-mix PBDE formulation, during the study period (Turyk et al., 2010; Hoopmann et al., 2012).

Other papers published within this timeframe investigating BFR concentrations in human matrices, but not temporal trends, showed some interesting results. Concentrations of BDEs in serum from children in the U.S. have been positively associated with duration of breast feeding and socio-economic status (Bradman et al., 2012; Stapleton et al., 2012) as well as maternal serum during pregnancy and having no safe places to play (Bradman et al., 2012). In a recent study of umbilical and maternal serum as well as human milk, the congener profile changed over the first month of lactation, however, maternal serum PBDE concentrations did not change significantly (Jakobsson et al., 2012).

3. Sewage sludge and wastewater

In sewage sludge samples from China (Yang et al., 2011), Korea (Hwang et al., 2012) and Italy (Cincinelli et al., 2012), BDE209 dominated the profile. In Chicago, USA, from 1975-2008, penta-PBDE concentrations increased initially and leveled-off around 2000 (production of the penta-mix PBDE product ceased in the USA in 2004) (Hale et al., 2012). Concentrations of BDE209 in biosolids rose from 1995-2008, doubling every 5 years, another indication of the size of the environmental reservoir of BDE209 which has been created.

Sewage sludge and wastewater samples will also be important matrices to examine in the future in order to establish which alternative FR compounds are being used as replacements for PBDEs and HBCD, as they concentrate inputs from diffuse industrial and domestic sources into a point source for release to the environment, which can then readily be sampled.

4. Water and suspended particulates

Few data have been reported recently in these matrices, which is unsurprising as their partitioning and accumulation characteristics make other matrices more attractive for study as seawater concentrations are very low. However, a study of novel brominated and phosphorus flame retardants (PFRs) in the River Aire, UK, has highlighted a type of passive sampling device, the ceramic dosimeter, which may be more widely applicable in the future for flame retardant studies (Cristale et al., 2013).

5. Sediments, plants and soils

5.1 *Sediments*

Many of the studies reported add to concerns relating to debromination and possible consequent toxic effects from the now considerable reservoir and predominance of BDE209 in sediments across the world due to continued and extensive use of the deca-mix PBDE formulation. Evidence for debromination is outlined in a later section of this paper.

In Taihu Lake in China, Zhou et al. (2012) estimated the inventory of BDE209 to be 26.3 tonnes, a substantial amount, although similar estimates have not been made for other lakes. This gives additional weight to the potential significance of BDE209 debromination in the future. BDE209 dominated the PBDE congener profiles in Taiwan (Jiang et al., 2011) Korea (Hong et al., 2010; Lee et al., 2012; Moon et al., 2007), Indonesia (Ilyas et al., 2011), in the Clyde estuary UK (Vane et al., 2010), Canada (Grant et al., 2011) and in the Scheldt estuary (Van Ael et al., 2012). Significant concentrations of HBCDs were observed in sediments from Tianjin, China (Zhang et al., 2013), Korea (Hong et al., 2010) and Indonesia (Ilyas et al., 2011).

Polychaete worms have been shown to be able to remobilise BFRs buried at 50 cm depth to the surface sediments (Josefsson et al., 2010; Dale and Miller, 2008), so contaminants buried at earlier times can re-enter the environmental cycle by this mechanism. This may also prove to have an impact on the remobilisation of BDE209 degradation products in the future.

5.2 *Plants and soils*

Plants have been shown to be able to translocate and remobilise PBDEs from soils, although at a lower level than for chlorobiphenyls (CBs) (e.g. Zhao et al., 2012). PBDEs were also more susceptible to metabolism than CBs of the same level of halogen substitution. Wu et al. (2012a) demonstrated isomer-specific HBCD accumulation in maize, and consequent oxidative stress and DNA damage. Li et al. (2011) showed that uptake of HBCDs into plants, and so into the human diet, was limited by adsorption to the soil matrix.

6. **Biota**

6.1 *Terrestrial and aquatic-based mammals*

Reflecting the paucity of research characterizing the exposure of biota to environmental contaminants in these ecosystems, only four studies have been published between 2010 and 2012. Novel research was presented on the exposure of bats to PBDEs (Kannan et al., 2010). PBDE-exposure of two species of otters in the UK and North America was also characterized.

PBDEs have now been determined in insectivorous brown bats from the USA (Kannan et al., 2010). BDE99 (41%) and BDE47 (39%) dominated adipose Σ BDE concentrations (118-8,290 $\mu\text{g kg}^{-1}$ lipid weight (lw) and BDE28 showed a greater proportion of the BDE burden in their brains than in adipose tissue.

In England and Wales, concentrations of Σ PBDE₂₆ in liver of 30 Eurasian otters were 92 – 19,900 $\mu\text{g kg}^{-1}$ lw. BDE47 dominated the profile (78%) of these otters followed by BDE153 and BDE100 (Walker et al., 2012). In the north-eastern US, the hepatic Σ PBDE₇ concentrations (15 – 8,480 mg kg^{-1} lw) of a different species of otter, river otters, were also dominated by BDE47 (62%) and did not show any age- or sex-related differences in concentrations (Stansley et al., 2010). On Vancouver Island, Canada, fecal Σ PBDE₁₂ concentrations of river otters were low but significantly higher in otters from Victoria Harbour (0.04 – 2.7 mg kg^{-1} lw) than from Esquimalt Harbour (0.01 – 0.44 mg kg^{-1} lw) or outside of the harbours (0.01 – 1.5 mg kg^{-1} lw) (Guertin et al., 2010).

6.2 *Birds' tissues and eggs*

There have been approximately 46 studies published between 2010 and 2012 addressing concentrations, temporal trends and spatial patterns, in PBDEs and/or HBCDs in birds. Several studies have reported on the isomeric HBCD pattern in bird tissues, with the dominance of α -HBCD apparent in many but not all birds (e.g., Guerra et al. 2012). The highest concentrations of HBCDs have now been observed in birds of prey, notably in a peregrine falcon egg from Montreal, Canada collected in 2007 (Σ HBCDs 14,600 $\mu\text{g kg}^{-1}$ lw; Guerra et al., 2012), with α -HBCD dominating, and in the muscle of a sparrowhawk collected in 1995 from an undeclared location in the UK (Σ HBCDs 19,000 $\mu\text{g kg}^{-1}$ lw; Leslie et al., 2011), with γ -HBCD dominating. Several important review papers describing concentrations and trends of these particular BFRs in bird tissues have been published during this time period, notably an overall review of concentrations in bird tissues by Chen and Hale (2010), a review of PBDEs, HBCD and novel FRs in birds and other wildlife from China (Wu et al., 2012b) and the Arctic (de Wit et al., 2010; Letcher et al., 2010), and in specific bird species (Henny et al., 2010).

Several studies have concluded that the geographical patterns and trends of PBDEs and HBCDs measured in the eggs of peregrine falcons (Guerra et al., 2012; de Wit et al., 2010; Chen et al., 2012), ospreys (Henny et al., 2011), herring gulls (Fliedner et al., 2012; Sørmo et al., 2011; Carlsson et al., 2011), and other bird species (Custer et al., 2010; Sun et al., 2012a&b), reflect continental usage patterns, human population densities, and/or introduced legislation governing the manufacturing and usage of these BFRs. Other studies have concluded that sources of exposure to PBDEs as reflected in bird eggs have changed over time (e.g., Crosse et al., 2012).

Global comparisons were made of median and maximal PBDE concentrations in bird tissues (liver, eggs; 2005 – 2010) published between 2010 and 2012. A detailed discussion of the methods used in this analysis is provided in the SI. Median PBDE concentrations of bird tissues were similar regardless of diet, ecosystem or continent, contrasting with continental differences in aquatic birds reported over a longer time period by Chen and Hale (2010). However, a similar significant continental pattern was observed

with maximal PBDE levels of freshwater birds ($P = 0.02$), all of which were piscivorous or omnivorous with one exception (Morales et al., 2012). Freshwater birds in North America had higher PBDE concentrations ($N = 6$; median: $6775 \mu\text{g kg}^{-1} \text{lw}$) than those in Asia ($N = 12$; median: $2100 \mu\text{g kg}^{-1} \text{lw}$) or Europe ($N = 2$; median: $560 \mu\text{g kg}^{-1} \text{lw}$) (Figure 2). These continental differences remained evident ($P = 0.5$) following the removal of the particularly high maximal PBDE concentration ($70,000 \mu\text{g kg}^{-1} \text{lw}$) in osprey (Henny et al., 2011). The continental pattern for freshwater birds likely reflects the greater historical usage of PBDEs in North America (Chen and Hale 2010 and references therein). Globally, the analysis also revealed that terrestrial birds had higher maximal PBDE concentrations ($N = 8$; median: $6600 \mu\text{g kg}^{-1} \text{lw}$) than freshwater ($N = 20$; median: $3250 \mu\text{g kg}^{-1} \text{lw}$) or marine birds ($N = 8$; median: $318 \mu\text{g kg}^{-1} \text{lw}$) ($P = 0.02$), possibly reflecting long-range transport patterns of PBDEs.

6.3 Fish, shellfish and benthic invertebrates

Data were reported for a number of areas/species, including a number of fish species from the Rhone River, the Scheldt estuary, the Bizerte lagoon in Tunisia, rivers in Germany, Norwegian waters and from Brazil. In fish from the Gila River in Arizona, USA, high concentrations of ΣBDE_{50} were reported (up to 12.7 mg kg^{-1} wet weight), amongst the highest in the USA (Echols et al., 2013). Congeners with 2,3,4- or 2,4,5- substitution on one ring were seen to be preferentially metabolised.

Despite some evidence of decreasing trends in some species and locations, elevated concentrations of BDEs in fish are still reported. Shanmuganathan et al. (2011) reported ΣBDE_{11} concentrations of $1.0 - 45 \mu\text{g kg}^{-1}$ fresh weight in edible fish products from markets in Adelaide, Australia. BDE209 was the dominant congener, followed by BDE47, BDE99 and BDE100.

In China, relatively low levels of HBCD were determined in 12 edible fish species from South China. The concentrations of ΣHBCDs ranged from non-detectable to $0.19 \mu\text{g kg}^{-1}$ wet weight, with a detection frequency of 70%. This was at the lower end of the concentration range observed globally. Moreover, ΣHBCDs concentrations were higher in both freshwater and seawater farmed fish than in wild marine fish, indicating that human activities probably represent an important input source of HBCD in aquaculture (Meng et al., 2012), although whether via discharges to water or direct usage in their activities is unclear.

Similar low HBCD levels in China were reported by Xia et al, 2011, reporting HBCD levels in two species of marine fish from nine Chinese coastal cities. HBCDs were detected in all samples analyzed, indicating ubiquitous contamination of these compounds in the Chinese coastal environment. The average ΣHBCDs concentration was $3.7 \mu\text{g kg}^{-1} \text{lw}$ (range: $0.57-10 \mu\text{g kg}^{-1} \text{lw}$), which is lower than seen in many other regions of the world, especially Europe, where HBCDs have been used more intensively than elsewhere. Geographically, the highest HBCD level present in fish was found in Dalian, in northern China, and the lowest occurred in Wenzhou (Xia et al., 2011).

In Japan, similarly low levels of ΣHBCDs were observed in 54 wild and 11 farmed seafood samples from four regions. For the fish classified as *Anguilliformes*, *Perciformes*, *Clupeiformes* and farmed *Salmoniformes*, the median concentrations of ΣHBCDs were 2.1, 0.75, 0.12 and $1.3 \mu\text{g kg}^{-1} \text{ww}$,

respectively. However, HBCDs were not detected in samples classified as *Crustacea*, *Mollusca*, *Pleuronectiformes* and *Scorpaeniformes* or, if detected, the levels were very low (Nakagawa et al, 2010).

6.4 Turtles

Significant knowledge about the PBDE exposure of sea turtles, many of which are threatened or endangered, was gained between 2010 and 2012 (Swarthout et al., 2010; Alava et al., 2011; Malarvannan et al., 2011; Ragland et al., 2011; Stewart et al., 2011). As with birds, maternal transfer of PBDEs to eggs and hatchling sea turtles occurs (Stewart et al., 2011), and maternal diet likely influences PBDE egg concentrations (Alava et al., 2011). BDE47 was usually the dominant PBDE congener (Swarthout et al. 2010), but for several species, PBDE concentrations and profiles differed between migratory and residential turtles (e.g., Ragland et al. 2011). A geographical gradient of increasing Σ BDE egg yolk concentrations for loggerhead turtles occurred from around Florida ($< 0.1 - 7.8 \text{ ng g}^{-1} \text{ lw}$) to North Carolina ($0.4 - 37 \text{ ng g}^{-1} \text{ lw}$) (Alava et al., 2011).

6.5 Marine mammals

Concentrations of PBDEs were determined in a number of cetacean species from Brazilian waters, of particular interest as the majority of data available hitherto are for animals from the Northern Hemisphere (Yogui et al., 2011; Leonel et al., 2012; Dorneles et al., 2010). In 10 cetacean species, concentrations in liver ranged from $3 - 5,960 \mu\text{g kg}^{-1} \text{ lw}$, similar to concentrations seen in Northern Hemisphere dolphins. The highest concentration was seen in a false killer whale which feeds over the continental shelf and, in general, higher concentrations were seen in species feeding in estuarine areas or over the continental shelf than in oceanic species.

Mongillo et al. (2012) used an individual-based modelling approach to predict accumulation of ten PBDE congeners in specific individuals within the southern resident killer whale populations off the west coasts of Canada and the USA. Model predictions for current concentrations agreed well with Σ BDE₁₀ concentrations measured in biopsy samples from known individuals. The predicted concentrations over the life-span of individual killer whales were consistent with a doubling time of ca. 3 – 4 years.

In adipose tissue of polar bears from western Hudson Bay, sampled at intervals between 1991 and 2007, concentrations of PBDEs showed significant increases of 13% annually between 1991 and 2007 (McKinney et al., 2010). During the 1990s, α -HBCD was not detected initially, but, following 2000, was detected at concentrations 5- to 10-fold lower than those of the PBDEs, and with a maximum concentration observed in 2003. There were differences in the relative proportions of BDE47 and BDE153, with adult males having lower proportions of BDE47 and higher proportions of BDE153 than adult females and sub-adult bears, which significant statistical correlations suggest reflect dietary differences and the modifying role of lactation on dietary uptake by females and sub-adults (McKinney et al., 2010).

6.6 Food chain studies

Vuorinen et al. (2012) studied biomagnification of PBDEs in Atlantic salmon from its main prey species (sprat and herring) in three areas of the Baltic Sea. BDE accumulation seemed to be highly dependent on both age and fat content, illustrating both the lipophilicity of these contaminants and past accumulation in tissues.

Pectoral muscle of kingfishers (*Alcedo atthis*) and their prey fish species from an electronic recycling site in south China were assessed for the concentrations and potential for biomagnification of PBDEs (17 congeners), DBDPE and BTBPE (Mo et al., 2012); here, we report on the PBDE findings only. In the kingfishers, ΣBDE_{17} levels ranged from 2,030 to 26,400 $\mu\text{g kg}^{-1}$ lw (median: 8,760 $\mu\text{g kg}^{-1}$ lw) which is several times higher than those in the muscles of 7 waterbird species (37-2,300 $\mu\text{g kg}^{-1}$ lw) and 3 terrestrial species (1,000-5,200 $\mu\text{g kg}^{-1}$ lw) from the same e-waste recycling site (Zhang et al., 2011b; Sun et al., 2012a). BDE47 was the predominant congener (~39% contribution) followed by BDE154, BDE153, BDE100 and BDE99 in descending order and contributing ~50% to the ΣBDE_{17} concentrations. The biomagnification factors (BMFs) for all PBDE congeners, except BDE28, were greater than unity, with mean BMFs of 1.4 to 4.5 for the ΣBDE_{17} depending on the fish species, and the authors conclude that these were within the range of most BMFs calculated in other bird studies involving different tissues (Mo et al., 2012).

7. Summary of time trends

7.1 Europe

Schuster et al. (2011) collected background soils from 70 locations along a latitudinal transect between the UK and Norway in 2008, 10 years after their first analysis in 1998 (for a map see Law et al. (2008)). Comparisons between the datasets showed a general decline in PBDE concentrations over time. In sewage sludges from Sweden, Olofsson et al. (2012) reported a decrease (ca. 20% each year) in concentrations of BDE154 and BDE183 (representative of the penta-mix and octa-mix PBDE products, respectively) during 2004-2010, while concentrations of BDE209 (the major component of the deca-mix PBDE product) increased by 16% over the same period. The downward trend for BDE154 seemed to be linear, whilst the upward trend for BDE209 seemed to be increasingly steep.

In a study undertaken in Germany (Esslinger et al., 2011), pooled herring gull eggs from the North and Baltic Sea coasts were analysed for HBCDs. During 1998-2008, all egg pools were dominated by α -HBCD, and temporal trends showed increasing concentrations until 2000 before beginning to decline.

7.2 North America

In archived sewage sludge (biosolids) samples from Chicago, USA, from 1975-2008, penta-PBDE concentrations increased and levelled-off around 2000, while deca-PBDE concentrations rose from 1995-2008, doubling approximately every 5 years (Hale et al., 2012). Congener patterns in contemporary biosolids supported the contention that BDE209 can be debrominated to yield lesser brominated PBDE congeners (BDE206, BDE207 and BDE208, in particular, in this case).

In trout from the Great Lakes sampled during 1980-2009, Crimmins et al. (2012) recorded temporal trends for ΣBDE_5 . In Lakes Huron, Michigan and Ontario, decreases in concentration were observed after 2000-2001. In Lakes Erie and Superior, concentrations seem to have stabilised, but not begun to decline significantly.

In general, penta-BDE concentrations in a range of environmental media (air, sediment, wastewater effluent, landfill effluent, aquatic biota and birds) reported from Canada increased until approximately 2000, when levelling off or decreasing trends were observed, consistent with the introduction of control measures (Backus et al., 2010).

Chen et al. (2011) found elevated concentrations of HBCDs in fish from the USA. Carp from the Hyco River, Virginia, exhibited contrasting time trends for PBDEs and HBCD. Between 1999-2002 and 2006-2007, mean ΣHBCDs concentrations rose from 13 to 4,640 $\mu\text{g kg}^{-1}$ lw, whilst mean ΣBDE_9 concentrations fell from 40,700 to 9,140 $\mu\text{g kg}^{-1}$ lw over the same period. This suggests that HBCD usage in the USA may have risen following the termination of PBDE manufacture in the USA in 2004.

7.3 *The Arctic and Antarctic*

In western Hudson Bay, ΣBDE_7 concentrations in the adipose tissue of polar bears showed significant increases of 13% annually between 1991 and 2007 (McKinney et al., 2010).

Rotander et al. (2012) reported time trends of PBDEs in seven species of marine mammals (long-finned pilot whales, minke whales, fin whales, harbour porpoise, Atlantic white-sided dolphins, ringed and hooded seals) from the Arctic and the North Atlantic taken during 1986-2009. The highest levels were found in samples from the late 1990s-2000. In blubber of Weddell seals from the Antarctic taken in 2006, out of 7 PBDE congeners analyzed, BDE47 was detected in only two animals (Trumble et al., 2012).

Muir and de Wit (2010) reported some evidence that environmental levels of penta-PBDE related congeners were leveling off or beginning to decline in Arctic samples. They examined studies of Arctic char, burbot, lake trout, ringed seals, northern fulmar, thick-billed murre, Brünnich's guillemots, ringed seals and beluga and concluded that most studies indicated declining concentrations of BDE47 and BDE99. Most surprising was the predominance of BDE209 in air samples, all of it in the particle phase.

8. **E-waste recycling**

Over the period covered by this review, concern has grown considerably about the potential for environmental contamination arising from the rudimentary treatment of e-waste (Schnoor, 2012). A

particular focus has been on such activities in China (Chi et al, 2011), with a notable number of studies reporting relatively high PBDE and/or HBCD concentrations in birds from e-waste recycling areas in China (Figure 2) (Liu et al. 2010, Yu et al. 2011, Zhang et al. 2011b, Sun et al. 2012a, b). although recent Chinese government actions to limit the importation of e-waste has resulted in attention switching to other regions such as Africa, Thailand, and Vietnam (Chi et al, 2011; Muenhor et al, 2010; Tue et al, 2010). The beneficial impacts of this Chinese government action are beginning to become evident in that country, with decreasing concentrations reported in some studies (e.g. Fu et al., 2012; Zhang et al., 2012b). Future research in this aspect of the environmental impacts of BFRs is likely to focus on a better characterisation of the impacts on human exposure and health, and on filling the substantial gap in knowledge of the extent of such informal e-waste treatment activities in hitherto under-studied regions such as Africa. Also, further study of the formation and release of brominated and mixed brominated/chlorinated dioxins and furans during these activities and their possible health effects is warranted (Leung et al. 2011; Zhang et al. 2012c&d; Nguyen et al. 2013).

9. Environmental and laboratory evidence for the debromination of BDE209 and transformation of HBCDs

Of major concern is the possible debromination of the large reservoir of BDE209 in soils and sediments worldwide deriving from use of the deca-PBDE product, with the potential to yield large quantities of lower-brominated congeners which are both more mobile and more toxic over a considerable time period. This debromination has been shown to occur in sediments from Lake Thun, Switzerland (Bogdal et al., 2010), for example. In sediment cores from the Clyde estuary, UK, the proportion of nona-BDE congeners (BDE206, BDE207 and BDE208) was greater than reported for the deca-BDE product, and this may also represent debromination of BDE209 (Vane et al., 2010). Similarly, the large contribution of both octa- and nona-BDEs to Σ BDE was indicated as being due to BDE209 debromination. The presence of lower brominated BDE congeners in biota from Korea may also point in this direction, as the penta- and octa-PBDE products were little used in that country (Mo et al., 2012). Li et al (2011) studied BDE209 uptake and degradation in ryegrass, reporting microbial degradation of BDE209 with a higher proportion of di- to hepta-BDE congeners in plant tissues than in soil. Huang et al. (2013) studied degradation of four BDE congeners (BDE28, BDE47, BDE99 and BDE209) in *in vitro* root crude enzyme extracts from maize, ryegrass and pumpkin. Degradation of BDE209 was observed in all three extracts, yielding congeners BDE206, BDE207, BDE208 and BDE183 and lower brominated congeners down to BDE7 (which has a low enthalpy of formation; Zeng et al. (2008)), the major proportion of the transformation occurring within 24h of the start of the experiment. A stepwise loss of bromine was observed for all congeners studied, although the patterns of debrominated products produced differed between plant species. Given the large amount of the deca-mix PBDE product in sediments, it would be very useful to establish whether marine plants, from plankton to seaweeds, have a similar capability to degrade BDE209, and at what rates, and this is a topic for future study.

Wei et al. (2013) examined the photolytic debromination pathways of 13 BDE congeners, including BDE209, BDE208, BDE207, BDE206 and BDE183. The congeners were exposed to sunlight in hexane for

up to 64h. Photolytic degradation of BDE209 yielded 45 congeners, with from nona- to tri-BDE substitution. The environmental significance of this process, given the more limited light penetration in many terrestrial and aquatic environments, is unknown.

As described above, Lee et al. (2012) noted the large contributions of nona- and octa-BDE congeners to the total concentrations in riverine sediments of Korea. The relatively large contribution of tri- to hepta-BDE congeners relative to the limited use of the penta- and octa-mix PBDE products in Korea, and the high concentrations of BDE202 and BDE207, may reflect debromination of BDE209 (Mo et al., 2012). Hale et al. (2012) ascribed higher than expected concentrations of BDE206, BDE207 and BDE208 in US biosolids to debromination of BDE209.

Du et al. (2012) exposed zebrafish to the three individual HBCD isomers. Bioaccumulation was concentration dependent, and accumulation efficiencies, biomagnification factors, and half-lives for α -HBCD were the highest of those for the three HBCD isomers, suggesting that these factors may underlie the predominance of α -HBCD in biota samples. Munschy et al. (2011) exposed juvenile common sole to food spiked with six BDE congeners (BDE28, BDE47, BDE99, BDE100, BDE153, BDE209) over a 3 month period, followed by 5 months of depuration. Certain BDE congeners not present in the food were detected in fish tissues: BDE49 and another tetra-BDE congener; a penta-BDE congener; BDE154; BDE183 and another hepta-BDE congener; and BDE202. Sole were able to metabolise BDE209 to yield lower brominated congeners and preferential removal of *meta*-substituted bromines suggested that the deiodinase enzyme systems may play a role in the debromination process.

Echols et al. (2013) determined BDEs in fish (common carp, channel catfish and largemouth bass) taken from the Gila River, USA, in 2003. The presence of BDE155, BDE188 and BDE202 all indicated debromination of BDE209 in these fish. Additionally, congeners with 2,3,4- or 2,4,5- substitution were shown to be vulnerable to selective metabolism in carp.

McKinney et al. (2011) studied biotransformation of BDE209 using an in vitro system based on liver microsomes from various Arctic-feeding marine mammals. Significant depletion of BDE209 was observed, although no evidence of simply debrominated metabolites was observed and the biotransformation products could not be identified.

In harbour porpoises during 1992-2008, Law et al. (2010) noted increases in the proportions of BDE153, BDE154 and BDE100. The authors acknowledged that this may have arisen from changes in sources, but concluded that it was more likely to reflect debromination of higher congeners, such as BDE209, in the environment, either by fish or in bottom-living organisms exposed via sediments and then entering the food chain.

10. Knowledge gaps and recommendations for future research

Following earlier steep rises in concentrations of both PBDEs and HBCD, controls on the use of the penta- and octa-mix PBDE products are beginning to yield declines in concentration in environmental

samples. Major concerns still remain in relation to potential debromination of BDE209 and exposure to both people and animals around e-waste recycling sites. Much knowledge has been gained regarding concentrations of BDEs and HBCDs in birds and other wildlife, with concentrations reported from remote locations globally (e.g., Indian Ocean islands).

There is still a need to assess the time trends of PBDEs and HBCDs in human samples for North America (and Europe, at least for HBCDs). PBDE concentrations have decreased in Europe, but time trends for HBCDs are not so clear. PBDE and HBCD trends in Asia are also unclear.

Exposure of children to BFRs would be better assessed by direct measurements of PBDE and HBCD concentrations in their serum, rather than indirectly through estimation of dietary intake and dust ingestion. Some ethical issues relating to this have yet to be overcome, and the potential utility of non- or minimally invasive measures of body burden, such as hair and nails, should be investigated.

While evidence continues to mount that indoor contamination is an important influence on human body burdens, larger studies that evaluate the role of indoor exposures versus diet would be welcome. Likewise, while indoor dust contamination appears to correlate with body burdens for some individuals; studies that clarify the exact mechanism(s) via which this occurs (e.g. ingestion, dermal contact) are required.

Due to the ongoing restrictions on PBDEs and the impending inclusion of HBCDs in the Stockholm Convention on POPs, there has been a shift in the use of FRs, with the recent increasing consumption of novel BFRs and OPFRs. Investigation of occurrence and time trends in various environmental media, biota and human samples for these alternative FRs is strongly recommended.

There is a gap in knowledge concerning the metabolism/debromination of BDE209 and HBCD in birds, and concerning the potential effects of these FRs and PBDEs in birds and mammals. Also, there is minimal information regarding the exposure and possible effects of these compounds in amphibians and reptiles.

E-waste recycling and dumping sites are a continuing topic of interest for research, especially in Asia and Africa. Further monitoring of human exposure and environmental contamination in these regions are warranted. In particular, the formation and release of brominated and mixed brominated/chlorinated dioxins and furans during these activities and their possible health effects should be addressed.

Given the large amount of the deca-mix PBDE product in sediments, it would be very useful to establish whether marine plants, from plankton to seaweeds, have a similar capability to degrade BDE209 to that of the terrestrial plants mentioned above (see section 9), and at what rates, and this is a topic for future study.

Acknowledgements

At Cefas, work on flame retardants has been funded by the (now defunct) Department of the Environment, the Department for Environment, Food and Rural Affairs and the Bromine Science Environmental Forum. AC is funded by the University of Antwerp and the Flanders Funds for Scientific Research (FWO). The contribution of SH and AC has received funding from the European Union Seventh Framework Programme [FP7/2007-2013] within the INTERFLAME grant agreement no. 295138. KF is funded by Environment Canada, and LT is funded by ARC DECRA 120100161. This review was originally prepared for and presented at BFR2013, the 6th International Symposium on Flame Retardants, held in San Francisco CA, USA, on 7 – 10 April 2013.

References

- Abdallah MA-E, Harrad S. Modification and calibration of a passive air sampler for monitoring vapor and particulate phase brominated flame retardants in indoor air: application to car interiors. *Environ Sci Technol* 2010; 44: 3059-3065.
- Abdallah MA-E, Harrad S, Ibarra C, Diamond M, Melymuk L, Robson M, Covaci A. Hexabromocyclododecanes In Indoor dust From Canada, the United Kingdom, and the United States. *Environ Sci Technol* 2008; 42: 459–464.
- Alava JJ, Keller JM, Wyneken J, Crowder L, Scott G, Kucklick JR. Geographical variation of persistent organic pollutants in eggs of threatened loggerhead sea turtles (*Caretta caretta*) from southeastern United States. *Environ Toxicol Chem* 2011; 30: 1677-1688.
- Alivernini S, Battistelli CL, Turrio-Baldassarri L. Human milk as a vector and an indicator of exposure to PCBs and PBDEs: temporal trend of samples collected in Rome. *Bull Environ Contam Toxicol* 2011; 87: 21-25.
- Andersson Ö, Blomkvist G. Polybrominated aromatic pollutants found in fish from Sweden. *Chemosphere* 1981; 10: 1051-1060.
- Anon. Illegal WEEE exporters to pay out £220,000. *Ends Report* 2012; 455: 40-41. Environmental Data Services, London.
- Asante KA, Adu-Kumi S, Nakahiro K, Takahashi S, Isobe T, Sudaryanto A, Devanathan G, Clarke E, Ansa-Asare OD, Dapaah-Siakwan S, Tanabe S. Human exposure to PCBs, PBDEs and HBCDs in Ghana: temporal variation, sources of exposure and estimation of daily intakes by infants. *Environ Int* 2011; 37: 921-928.
- Backus SM, Martin P, Lewis M, Banic C, Spry D, Marvin C, Pelletier M, Smyth SA, Berthiaume A, Fraser S, Shen J, Harris J. Integrating multi-media monitoring of PBDEs in the Canadian environment: current status and trends. *Proceedings of BFR2010, Kyoto, Japan.* #90156.
- Batterman S, Godwin C, Chernyak S, Jia S, Charles S. Brominated flame retardants in offices in Michigan, U.S.A. *Environ Int* 2010; 36: 548-556.
- BBC. Europe still exporting electronic waste. 2010. www.bbc.co.uk/news/world-europe-10846395 accessed 4th August 2010.
- BJörklund JA, Thuresson K, Palm Cousins A, Sellström U, Emenius G, de Wit C. Indoor air is a significant source of tri-decabrominated diphenyl ethers to outdoor air via ventilation systems. *Environ Sci Technol* 2012; 46: 5876-5884.
- Bogdal C, Scheringer M, Schmid P, Bläuenstein M, Kohler M, Hungerbühler K. Levels, fluxes and time trends of persistent organic pollutants in Lake Thun, Switzerland: combining trace analysis and multimedia modelling. *Sci Total Environ* 2010; 408: 3654-3663.

- Bradman A, Castorina R, Sjödin A, Fenster L, Jones RS, Harley KG, Chevrier J., Holland NT, Eskenazi B. Factors associated with serum polybrominated diphenyl ether (PBDE) levels among school-age children in the CHAMACOS cohort. *Environ Sci Technol* 2012; 46: 7373-7381.
- Carlsson P, Herzke D, Wedborg M, Gabrielsen GW. Environmental pollutants in the Swedish marine ecosystem, with special emphasis on polybrominated diphenyl ethers (PBDE). *Chemosphere* 2011; 82: 1286-1292.
- Chen D, Hale RC. A global review of polybrominated diphenyl ether flame retardant contamination in birds. *Environ Int* 2010; 36: 800-811.
- Chen D, La Guardia MJ, Luellen DR, Harvey E, Mainor TM, Hale RC. Do temporal and geographic patterns of HBCD and PBDE flame retardants in US fish reflect evolving industrial usage? *Environ Sci Technol* 2011; 45: 8254-8261.
- Chen D, Letcher RJ, Burgess NM, Champoux L, Elliott JE, Hebert CE, Martin P, Wayland M, Weseloh DVC, L Wilson. Flame retardants in eggs of four gulls species (Laridae) from breeding sites spanning Atlantic to Pacific Canada. *Environ Pollut* 2012; 168: 1-9.
- Chi A, Streicher-Porte M, Wang MYL, Reuter MA. Informal electronic waste recycling: a sector review with special focus on China. *Waste Management* 2011; 31: 731-742.
- Cincinelli A, Martellini T, Misuri L, Lanciotti E, Sweetman A, Laschi S, Palchetti I. PBDEs in Italian sewage sludge and environmental risk of using sewage sludge for land application. *Environ Pollut* 2012; 161: 229-234.
- Covaci A, Harrad S, Abdallah MA-E, Ali N, Law RJ, Herzke D, de Wit CA. Novel brominated flame retardants: a review of their analysis, environmental fate and behaviour. *Environ Int* 2011; 37: 532-556.
- Covaci A, Roosens L, Dirtu AC, Waegeneers N, Van Overmeire I, Neels H, Goeyens L. Brominated flame retardants in Belgian home-produced eggs: levels and contamination sources. *Sci Total Environ* 2009; 407: 4387-4396.
- Crimmins BS, Pagano JJ, Xia X, Hopke PK, Milligan MS, Holsen TM. Polybrominated diphenyl ethers (PBDEs): turning the corner in Great Lakes trout 1980-2009. *Environ Sci Technol* 2012; 46: 9890-9897.
- Cristale J, Katsoyiannis A, Chen C, Jones KC, Lacorte S. Assessment of flame retardants in river water using a ceramic dosimeter passive sampler. *Environ Pollut* 2013; 172: 163-169.
- Crosse JD, Shore RF, Jones KC, Pereira MG. Long term trends in PBDE concentrations in gannet (*Morus bassanus*) eggs from two UK colonies. *Environ Pollut* 2012; 161: 93-100.
- Custer TW, Kannan K, Tao L, Yun S.H., Trowbridge A. Perfluorinated compounds and polybrominated diphenyl ethers in great blue heron eggs from three colonies on the Mississippi River, Minnesota. *Waterbirds* 2010; 33: 86-95.
- Dale RK, Miller DC. Hydrologic interactions of infaunal polychaetes and intertidal groundwater discharge. *Mar Ecol Prog Ser* 2008; 363: 205-215.
- de Wit CA. An overview of brominated flame retardants in the environment. *Chemosphere* 2002; 46: 583-624.
- de Wit CA, Herzke D, Vorkamp K. Brominated flame retardants in the Arctic environment - trends and new candidates. *Sci Total Environ* 2010; 408: 2885-2918.
- Dickhut RM, Cincinelli A, Cochran M, Kylin H. Aerosol-mediated transport and deposition of brominated diphenyl ethers to Antarctica. *Environ Sci Technol* 2012; 46: 3135-3140.
- Dirtu AC, Abdallah M, Covaci A. Advances in the sample preparation of brominated flame retardants and other brominated compounds. *TrAC* 2013; 43: 189-203.
- Dirtu AC, Covaci A. Estimation of daily intake of organohalogenated contaminants from food consumption and indoor dust ingestion in Romania. *Environ Sci Technol* 2010; 44: 6297-6304.

- Domingo, JL. Influence of cooking processes on the concentrations of toxic metals and various organic environmental pollutants in food: A review of the published literature. *Crit Rev Food Sci Nutr* 2011; 51: 29-37.
- Domínguez AA, Law RJ, Herzke D, de Boer J. Bioaccumulation of brominated flame retardants. In: Eljarrat, E. and Barceló, D. (eds.) *Brominated Flame Retardants*. 2011. *Hdb Env Chem* 16: 141-185. doi: 10.1007/698_2010_95. Springer-Verlag Berlin Heidelberg, Germany.
- Dorneles PR, Lailson-Brito J, Dirtu AC, Weijs L, Azevedo AF, Torres JPM, Malm O, Neels H, Blust R, Das K, Covaci A. Anthropogenic and naturally-produced organobrominated compounds in marine mammals from Brazil. *Environ Int* 2010; 36: 60-67.
- Du M, Lin L, Yan C, Zhang X. Diastereoisomer- and enantiomer-specific accumulation, depuration, and bioisomerization of hexabromocyclododecanes in zebrafish (*Danio rerio*). *Environ Sci Technol* 2012; 46: 11040-11046.
- Echols KR, Peterman PH, Hinck JE, Orazio CE. Polybrominated diphenyl ether metabolism in field collected fish from the Gila River, Arizona, USA – levels, possible sources, and patterns. *Chemosphere* 2013; 2013: 20-27.
- Esslinger S, Becker R, Jung C, Schröter-Kermani C, Bremser W, Nehls I. Temporal trend (1988-2008) of hexabromocyclododecane enantiomers in herring gull eggs from the German coastal region. *Chemosphere* 2011; 83: 161-167.
- Fliedner A, Rüdell H, Jürling H, Müller J, Neugebauer F, Schröter-Kermani C. Levels and trends of industrial chemicals (PCBs, PFCs, PBDEs) in archived herring gull eggs from German coastal regions. *Environ Sci Europe* 2012; 24: 1-15.
- Fu J, Wang T, Wang P, Qu G, Wang Y, Zhang Q, Zhang A. Temporal trends (2005-2009) of PCDD/Fs, PCBs, PBDEs in rice hulls from an e-waste dismantling area after stricter environmental regulations. *Chemosphere* 2012; 88: 330-335.
- Gentes M-L, Letcher RJ, Caron-Beaudoin E, Verreault J. Novel flame retardants in urban-feeding ring-billed gulls from the St Lawrence River, Canada. *Environ Sci Technol* 2012; 46: 9735-9744.
- Grant PBC, Johannessen SC, MacDonald RW, Yunker MB, Sanborn M, Dangerfield N, Wright C, Ross PS. Environmental fractionation of PCBs and PBDEs during particle transport as recorded by sediments in coastal waters. *Environ Toxicol Chem* 2011; 30: 1522-1532.
- Guerra P, Alae M, Jiménez B, Pacepavicius G, Marvin C, MacInnis G, Eljarrat E, Barceló D, Champoux L, Fernie K. Emerging and historical brominated flame retardants in peregrine falcon (*Falco peregrinus*) eggs from Canada and Spain. *Environ Int* 2012; 40: 179-186.
- Guertin DA, Harestad AS, Ben-David M, Drouillard KG, Elliott JE. Fecal genotyping and contaminant analyses reveal variation in individual river otter exposure to localized persistent contaminants. *Environ Toxicol Chem* 2010; 29: 275-284.
- Hale RC, La Guardia MJ, Harvey E, Chen D, Mainor TM, Luellen DR, Hundal LS. Polybrominated diphenyl ethers in U.S. sewage sludges and biosolids: temporal and geographical trends and uptake by corn following land application. *Environ Sci Technol* 2012; 46: 2055-2063.
- Harrad S, Ibarra C, Abdallah A-E, Boon R, Neels H, Covaci A. Concentrations of brominated flame retardants in dust from United Kingdom cars, homes, and offices: causes of variability and implications for human exposure. *Environ Int* 2008; 34: 1170–1175.
- Henny CJ, Grove RA, Kaiser JL, Johnson BL. North American osprey populations and contaminants: Historic and contemporary perspectives. *J Toxicol Environ Health – Part B. Crit Rev* 2010; 13: 579-603.
- Henny CJ, Grove RA, Kaiser JL, Johnson BL, Furl CV, Letcher RJ. 2011. Wastewater dilution index partially explains observed polybrominated diphenyl ether flame retardant concentrations in osprey eggs from Columbia River Basin, 2008-2009. *Ecotoxicol* 2011; 20: 682-697.

- Hong SH, Kannan N, Jin Y, Won JH, Han GM, Shim WJ. Temporal trend, spatial distribution, and terrestrial sources of PBDEs and PCBs in Masan Bay, Korea. *Mar Pollut Bull* 2010; 60: 1836-1841.
- Hoopmann M, Albrecht UV, Gierden E, Huppmann R, Suchenwirth R. Time trends and individual characteristics associated with polybrominated diphenyl ethers in breast milk samples 2006-2009 in Lower Saxony, Germany. *Int J Hyg Environ Health* 2012; 215: 352-359.
- Huang H, Zhang S, Wang S, Lv J. In vitro biotransformation of PBDEs by root crude enzyme extracts: potential role of nitrate reductase (NaR) and glutathione S-transferase (GST). *Chemosphere* 2013; 90: 1885-1892.
- Huwe JK, West M. Polybrominated diphenyl ethers in US meat and poultry from two statistically designed surveys showing trends and levels from 2002 to 2008. *J Agric Food Chem* 2011; 59: 5428-5434.
- Hwang I-K, Kang H-H, Lee I-S, Oh J-E. Assessment of characteristic distribution of PCDD/Fs and BFRs in sludge generated at municipal and industrial wastewater treatment plants. *Chemosphere* 2012; 88: 888-894.
- Ilyas M, Sudaryanto A, Setiawan IE, Riyadi AS, Isobe T, Takahashi S, Tanabe S. Characterization of polychlorinated biphenyls and brominated flame retardants in sediments from riverine and coastal waters of Surabaya, Indonesia. *Mar Pollut Bull* 2011; 62: 89-98.
- Jakobsson K., Fang J., Athanasiadou M., Rignell-Hydbom A., Bergman, Å. Polybrominated diphenyl ethers in maternal serum, umbilical cord serum, colostrum and mature breast milk. Insights from a pilot study and the literature. *Environ Int* 2012; 47: 121-130.
- Jiang J-J, Lee C-L, Fang M-D, Ko F-C, Baker JE. Polybrominated diphenyl ethers and polychlorinated biphenyls in sediments of southwest Taiwan: regional characteristics and potential sources. *Mar Pollut Bull* 2011; 62: 815-823.
- Johnson PI, Stapleton HM, Sjödin A, Meeker JD. Relationships between polybrominated diphenyl ether concentrations in house dust and serum. *Environ Sci Technol* 2010; 44: 5627-5632.
- Josefsson S, Leonardsson K, Gunnarsson JS, Wiberg K. Bioturbation-driven release of buried PCBs and PBDEs from different depths in contaminated sediments. *Environ Sci Technol* 2010; 44: 7456-7464.
- Kalachova K, Hradkova P, Lankova D, Hajslova J., Pulkrabova, J. Occurrence of brominated flame retardants in household and car dust from the Czech Republic. *Sci Total Environ* 2012; 441: 182-193.
- Kannan K, Yun SH, Rudd RJ, Behr M. High concentrations of persistent organic pollutants including PCBs, DDT, PBDEs and PFOS in little brown bats with white-nose syndrome in New York, USA. *Chemosphere* 2010; 80: 613-618
- Kefeni KK, Okonkwo JO. Analysis of major congeners of polybromobiphenyls and polybromodiphenyl ethers in office dust using high resolution gas chromatography-mass spectrometry. *Chemosphere* 2012; 87: 1070-1075.
- Klosterhaus SL, Stapleton HM, La Guardia MJ, Greig DJ. Brominated and chlorinated flame retardants in San Francisco Bay sediments and wildlife. *Environ Int* 2012; 47: 56-65.
- Król S, Zabliegala B, Namiesnik J. Determination of polybrominated diphenyl ethers in house dust using standard addition method and gas chromatography with electron capture and mass spectrometric detection. *J Chromatog A* 2012; 1249: 201-214.
- Law RJ, Barry J, Bersuder P, Barber JL, Deaville R, Reid RJ, Jepson PD. Levels and trends of brominated diphenyl ethers in blubber of harbor porpoises (*Phocoena phocoena*) from the U.K., 1992-2008. *Environ Sci Technol* 2010; 44: 4447-4451.
- Law RJ, Herzke D. Current levels and trends of brominated flame retardants in the environment. In: Eljarrat, E. and Barceló, D. (eds.) *Brominated Flame Retardants*. 2011. *Hdb Env Chem* 16: 123-140. doi: 10.1007/698_2010_82. Springer-Verlag Berlin Heidelberg, Germany.

- Law RJ, Herzke D, Harrad S, Morris S, Bersuder P, Allchin CR. Levels and trends of HBCD and BDEs in the European and Asian environments, with some information for other BFRs. *Chemosphere*; 2008, **73**, 223-241.
- Law RJ, Losada S, Barber J, Bersuder P, Deaville R, Brownlow A, Penrose R, Jepson PD. Alternative flame retardants, Dechlorane Plus and BDEs in the blubber of harbour porpoises (*Phocoena phocoena*) stranded or bycaught in the UK during 2008. *Environ Int* 2013; 60: 81-88.
- Lee I-S, Kim K-S, Kim S-J, Yoon JH, Choi KH, Choi S-D, Oh J-E. Evaluation of mono- to deca-brominated diphenyl ethers in riverine sediment of Korea with special reference to the debromination of DeBDE209. *Sci Total Environ* 2012; 432: 128-134.
- Leonel J, Taniguchi S, Sasaki DK, Cascaes MJ, Dias PS, Botta S, de O Santos MC, Montone RC. Contamination by chlorinated pesticides, PCBs and PBDEs in Atlantic spotted dolphin (*Stenella frontalis*) in western South Atlantic. *Chemosphere* 2012; 86: 741-746.
- Leslie HA, Leonards PEG, Shore RF, Walker LA, Bersuder PRC, Morris S, Allchin CR, de Boer J. Decabromodiphenylether and hexabromocyclododecane in wild birds from the United Kingdom, Sweden and The Netherlands: screening and time trends. *Chemosphere* 2011; 82: 88-95.
- Letcher RJ, Bustnes JO, Dietz R, Jenssen BM, Jørgensen EH, Sonne C, Verreault J, Vijayan MM, Gabrielsen GW. Exposure and effects assessment of persistent organohalogen contaminants in arctic wildlife and fish. *Sci Total Environ* 2010; 408: 2995-3043.
- Leung AOW, Zheng J, Yu CK, Liu WK, Wong CKC, Cai ZW, Wong MH. Polybrominated Diphenyl Ethers and Polychlorinated Dibenzo-p-dioxins and Dibenzofurans in Surface Dust at an E-Waste Processing Site in Southeast China. *Environ Sci Technol* 2011; 45: 5775-5782.
- Li Y, Zhou Q, Wang Y, Xie X. Fate of tetrabromobisphenol A and hexabromocyclododecane brominated flame retardants in soil and uptake by plants. *Chemosphere* 2011; 82: 204-209.
- Linderholm L, Biague A, Mansson F, Norrgren H, Bergman Å, Jakobsson K. Human exposure to persistent organic pollutants in West Africa - a temporal trend study from Guinea-Bissau. *Environ Int* 2010; 36: 675-682.
- Link B, Gabrio T, Mann V, Schilling B, Maisner V, König M, Flicker-Klein A, Zollner I, Fischer G. Polybrominated diphenyl ethers (PBDE) in blood of children in Baden-Württemberg between 2002/03 and 2008/09. *Int J Hyg Environ Health* 2012; 215: 224-228.
- Liu J, Luo X-J, Yu L-H, He M-J, Chen S-J, Mai B-X. polybrominated diphenyl ethers (PBDEs) polychlorinated biphenyls (PCBs), hydroxylated and methoxylated-PBDEs and methylsulfonyl-PCBs in bird serum from south China. *Arch Environ Contam Toxicol* 2010; 59: 492-501.
- Malarvannan G, Isobe T, Covaci A, Prudente M, Tanabe S. Accumulation of brominated flame retardants and polychlorinated biphenyls in human breast milk and scalp hair from the Philippines: levels, distribution and profiles. *Sci Total Environ* 2013; 442: 366-379.
- McKinney MA, Dietz R, Sonne C, De Guise S, Skirnisson K, Karlsson K, Steingrímsson E, Letcher RJ. Comparative hepatic microsomal biotransformation of selected PBDEs, including decabromodiphenyl ether, and decabromodiphenyl ethane flame retardants in Arctic marine-feeding mammals. *Environ Toxicol Chem* 2011; 30: 1506-1514.
- McKinney MA, Stirling I, Lunn NJ, Peacock E, Letcher RJ. The role of diet on long-term concentration and pattern trends of brominated and chlorinated contaminants in western Hudson Bay polar bears, 1991-2007. *Sci Total Environ* 2010; 408: 6210-6222.
- Meng XZ, Xiang N, Duan YP, Chen L, Zeng EY. Hexabromocyclododecane in consumer fish from South China: implications for human exposure via dietary intake. *Environ Toxicol Chem* 2012 ; 31 : 1424-1430.
- Mo L, Wu J-P, Luo X-J, Zou F-S, Mai B-X. Bioaccumulation of polybrominated diphenyl ethers, decabromodiphenyl ethane, and 1,2-bis(2,4,6-tribromophenoxy) ethane flame retardants in

- kingfishers (*Alcedo atthis*) from an electronic waste-recycling site in South China. *Environ Toxicol Chem* 2012; 31: 2153-2158.
- Mongillo TM, Holmes EE, Noren DP, VanBlaricom GR, Punt AE, O'Neill SM, Ylitalo GM, Hanson MB, Ross PS. Predicted polybrominated diphenyl ether (PBDE) and polychlorinated biphenyl (PCB) accumulation in southern resident killer whales. *Mar Ecol Prog Ser* 2012; 453: 263-277.
- Moon HB, Kannan K, Lee SJ, Choi M. Polybrominated diphenyl ethers (PBDEs) in sediment and bivalves from Korean coastal waters. *Chemosphere* 2007; 66: 243-251.
- Mora MA, Sericano JL, Baxter C. Swallows as indicators of environmental pollution of the Rio Grande/Rio Bravo basin: are persistent organic pollutants a concern? *Arch Environ Contam, Toxicol* 2012; 62: 512-518.
- Morales L, Martrat MG, Olmos J, Parera J, Vicente J, Bertolero A, Ábalos JM, Lacorte S, Santos FJ, Abad E. Persistent organic pollutants in gull eggs of two species (*Larus michahellis* and *Larus audouinii*) from the Ebro delta Natural Park. *Chemosphere* 2012; 88: 1306-1316.
- Muenhor D, Harrad S, Ali, N, Covaci A. Brominated flame retardants (BFRs) in air and dust from electronic waste storage facilities in Thailand. *Environ Int* 2010; 36: 690-698.
- Muir DCG, de Wit CA Trends of legacy and new persistent organic pollutants in the circumpolar Arctic: overview, conclusions and recommendations. *Sci Total Environ*; 2010: 3044-3051.
- Munsch C, Héas-Moisan K, Tixier C, Olivier N, Gastineau O, Le Bayon N, Buchet V. Dietary exposure of juvenile common sole (*Solea solea* L.) to polybrominated diphenyl etherds (PBDEs): part 1. Bioaccumulation and elimination kinetics of individual congeners and their debrominated metabolites. *Environ Pollut* 2011; 159: 229-237.
- Nakagawa R, Murata S, Ashikuza Y, Shintani Y, Hori T, Tsutsumi T. Hexabromocyclododecane determination in seafood samples collected from Japanese coastal areas. *Chemosphere* 2010; 81: 445-452.
- Nguyen MT, Takahashi S, Subramanian A, Sakai S, Tanabe S. Environmental contamination and human exposure to dioxin-related compounds in e-waste recycling sites of developing countries. *Environ Sci Proc Imp* 2013; 15:1326-1331.
- Olofsson U, Bignert A, Haglund P. Time-trends of metals and organic contaminants in sewage sludge. *Water Res.* 2012; 46: 4841-4851.
- Papachlimitzou A, Barber JL, Losada S, Bersuder P, Law RJ. A review of the analysis of novel brominated flame retardants. *J Chromatog A* 2012; 1219: 15-28.
- Ragland JM, Arendt MD, Kucklick JR, Keller JM. Persistent organic pollutants in blood plasma of satellite-tracked adult male loggerhead sea turtles (*Caretta caretta*). *Environ Toxicol Chem* 2011; 30: 1549-1556.
- Quirós-Alcalá L, Bradman A, Nishioka M, Harnly ME, Hubbard A, McKone TE, Eskenazi B. Concentrations and loadings of polybrominated diphenyl ethers in dust from low-income households in California. *Environ Int* 2011; 37 : 592-596.
- Rawn DFK, Sadler A, Quade SC, Sun WF, Lau BPY, Kosarac I, Hayward S, Ryan JJ. Brominated flame retardants in Canadian chicken egg yolks. *Food Addit Contam Part A* 2011; 28: 807-815.
- Rose M, Bennett D, Bergman Å, Fångström B, Pessah IN, Hertz-Picciotto I. PBDEs in 2-5 year old children from California and associations with diet and indoor environment. *Environ Sci Technol* 2010; 44: 2648-2653.
- Rotander A, van Bavel B, Polder A, Rigét F, Auðunsson GA, Gabrielsen GW, Víkingsson G, Bloch D, Dam M. Polybrominated diphenyl ethers (PBDEs) in marine mammals from Arctic and North Atlantic regions, 1986-2009. *Environ Int* 2012; 40: 102-109.
- Schnoor J. Extended producer responsibility for E-waste. *Environ Sci Technol* 2012; 46: 7927.

- Schechter A, Colacino J, Patel K, Kannan K, Yun SH, Haffner D, Harris TR, Birnbaum L. Polybrominated diphenyl ether levels in food stuffs collected from three locations from the United States. *Toxicol Appl Pharmacol* 2010; 243: 217-234.
- Schuster JK, Gioia R, Moeckel C, Agarwal T, Bucheli TD, Breivik K, Steinnes E, Jones KC. Has the burden and distribution of PCBs and PBDEs changed in European background soils between 1998 and 2008 ? Implications for sources and processes. *Environ Sci Technol* 2011; 45: 7291-7297.
- Shanmuganathan D, Megharaj M, Chen ZL, Naidu R. Polybrominated diphenyl ethers (PBDEs) in marine foodstuffs in Australia: residue levels and contamination status of PBDEs. *Mar Pollut Bull* 2011; 63: 154-159.
- Sørmo EG, Lie E, Ruus A, Gaustad H, Skaare JU, Jenssen BM. Trophic level determines levels of brominated flame-retardants in coastal herring gulls. *Ecotox Environ Safety* 2011; 74: 2091-2098.
- Stansley W, Velinsky D, Thomas R. Mercury and halogenated organic contaminants in river otters (*Lontra canadensis*) in New Jersey, USA. *Environ Toxicol Chem* 2010; 29: 2235-2242.
- Stapleton HM, Eagle S, Sjödin A, Webster TF. Serum PBDEs in a North Carolina toddler cohort: Associations with handwipes, house dust, and socioeconomic variables. *Environ Health Perspect* 2012; 120: 1049-1054.
- Stewart KR, Keller JM, Templeton R, Kucklick JR, Johnson C. Monitoring persistent organic pollutants in leatherback turtles (*Dermochelys coriacea*) confirms maternal transfer. *Marine Pollut Bull* 2011; 62: 1396-1409.
- Sun Y-X, Luo X-J, Mo L, He M-J, Zhang Q, Chen S-J, Zou F-S, Mai B-X. Hexabromocyclododecane in terrestrial passerine birds from e-waste, urban and rural locations in the Pearl River Delta, South China: levels, biomagnification, diastereoisomer- and enantiomer-specific accumulation. *Environ Pollut* 2012b; 171: 191-198.
- Sun YX, Luo XJ, Mo L, Zhang Q, Wu JP, Chen SJ, Zou FS, Mai BX. Brominated flame retardants in three terrestrial passerine birds from south China: Geographical pattern and implication for potential sources. *Environ Pollut* 2012a; 162: 381-388.
- Suzuki G, Nose K, Takigami H, Takahashi S, Sakai S, PBDEs and PBDD/Fs in house and office dust from Japan. *Organohalogen Compd.* 2006; 68: 1843-1846.
- Swarthout RF, Keller JM, Peden-Adams M, Landry AM, Fair PA, Kucklick JR. Organohalogen contaminants in blood of Kemp's ridley (*Lepidochelys kempii*) and green sea turtles (*Chelonia mydas*) from the Gulf of Mexico. *Chemosphere* 2010; 78: 731-741.
- Toms LML, Guerra P, Eljarrat E, Barceló D, Harden F, Hobson P, Sjödin A, et al., 2012. Brominated flame retardants in the Australian population: 1993-2009. *Chemosphere* 2012; 89: 398-403.
- Tornkvist A, Glynn A, Aune M, Darnerud PO, Ankarberg EH. PCDD/F, PCB, PBDE, HBCD and chlorinated pesticides in a Swedish market basket from 2005 - Levels and dietary intake estimations. *Chemosphere* 2011; 83: 193-199.
- Trudel D, Tlustos C, Von Goetz N, Scheringer M, Hungerbuehler K. PBDE exposure from food in Ireland: Optimising data exploitation in probabilistic exposure modeling. *J Exp Sci Environ Epidemiol* 2011a; 21 ; 565-575.
- Trudel D, Scheringer M, von Goetz N, Hungerbuehler K. Total consumer exposure to polybrominated diphenyl ethers in North America and Europe. *Environ Sci Technol* 2011b; 45: 2391-2397.
- Trumble SJ, Robinson EM, Noren SR, Usenko S, Davis J, Kanatous SB. Assessment of legacy and emerging persistent organic pollutants in Weddell seal tissue (*Leptonychotes weddellii*) near McMurdo Sound, Antarctica. *Sci Total Environ* 2012; 439: 275-283.
- Tue NM, Sudaryanto A, Minh TB, Isobe T, Takahashi S, Viet PH, Tanabe S. Accumulation of

- polychlorinated biphenyls and brominated flame retardants in breast milk from women living in Vietnamese e-waste recycling sites. *Sci Total Environ* 2010; 408: 2155-2162.
- Turyk ME, Anderson HA, Steenport D, Buelow C, Imm P, Knobeloch L. Longitudinal biomonitoring for polybrominated diphenyl ethers (PBDEs) in residents of the Great Lakes basin. *Chemosphere* 2010; 81: 517-522.
- Van Ael E, Covaci A, Blust R, Bervoets L. Persistent organic pollutants in the Scheldt estuary: environmental distribution and bioaccumulation. *Environ Int* 2012; 48: 17-27.
- Vane CH, Ma Y-J, Chen S-J, Mai B-X. Increasing polybrominated diphenyl ether (PBDE) contamination in sediment cores from the inner Clyde Estuary, UK. *Environ Geochem Health* 2010; 32: 13-21.
- Vorkamp K, Thomsen M, Frederiksen M, Pedersen M, Knudsen LE. Polybrominated diphenyl ethers (PBDEs) in the indoor environment and associations with prenatal exposure. *Environ Int* 2011; 37: 1-10.
- Vuorinen PJ, Keinänen M, Kiviranta H, Koistinen J, Kiljunen M, Myllylä T, Pönni J, Peltonen H, Verta M, Karjalainen J. Biomagnification of organohalogens in Atlantic salmon (*Salmo salar*) from its main prey species in three areas of the Baltic Sea. *Sci Total Environ* 2012; 421-422: 129-143.
- Walker LA, Moeckel C, Pereira MG, Beith SJ, Potter ED, Chadwick EA, Shore RF. Polybrominated diphenyl ethers (PBDEs) in Eurasian otters (*Lutra lutra*) collected from Britain in 2010: a Predatory Bird Monitoring Scheme (PBMS) Report. 2012. Centre for Ecology and Hydrology, Lancaster, UK. 14 pp.
- Wang X-P; Gong P; Yao T-D; Jones KC Passive air sampling of organochlorine pesticides, polychlorinated biphenyls, and polybrominated diphenyl ethers across the Tibetan Plateau. *Environ Sci Technol*. 2010a; 44: 2988-2993.
- Wang Y, Lam JCW, So MK, Yeung LWY, Cai Z, Hung CLH, Lam PKS. Polychlorinated dibenzo-*p*-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), dioxin-like polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in waterbird eggs of Hong Kong, China. *Chemosphere* 2012 ; 86 : 242-247.
- Wei H, Zou Y, Li A, Christensen ER, Rockne KJ. Photolytic debromination pathway of polybrominated diphenyl ethers in hexane by sunlight. *Environ Pollut* 2013; 174: 194-200.
- Wu J, Zhang Y, Luo X, She Y, Yu L, Chen S, Mai B. A review of polybrominated diphenyl ethers and alternative brominated flame retardants in wildlife from China: Levels, trends, and bioaccumulation characteristics. *J Environ Sci* 2012b; 24: 183-194.
- Wu T, Wang S, Huang HL, Zhang SZ. Diastereomer-specific uptake, translocation, and toxicity of hexabromocyclododecane diastereoisomers to maize. *J Agric Food Chem* 2012a; 60: 8528-8534.
- Xia CH, Lam JCW, Wu XG, Sun LG, Xie ZQ, Lam PKS. Hexabromocyclododecanes (HBCDs) in marine fishes along the Chinese coastline. *Chemosphere* 2011; 82: 1662-1668.
- Xiao H, Shen L, Su Y, Barresi E, Dejong M, Hung H, Lei Y-D, Wania F, Reiner E J, Sverko E, Kang S-C. Atmospheric concentrations of halogenated flame retardants at two remote locations: The Canadian High Arctic and the Tibetan Plateau. *Environ Pollut* 2012, 161: 154-161.
- Xu W, Wang X, Cai Z. Analytical chemistry of the persistent organic pollutants identified in the Stockholm Convention : a review. *Anal Chim Acta* 2013 ; 790 : 1-13.
- Yang C, Meng X-Z, Chen L, Xia S. Polybrominated diphenyl ethers in sewage sludge from Shanghai, China: possible ecological risk applied to agricultural land. *Chemosphere* 2011; 85: 418-423.
- Yogui GT, Santos MCO, Bertozzi CP, Sericano JL, Montone RC. PBDEs in the blubber of marine mammals from coastal areas of São Paulo, Brazil, southwestern Atlantic. *Mar Pollut Bull* 2011; 62: 2666-2670.
- Yu YX, Huang NB, Zhang XY, Li JL, Yu ZQ, Han SY, Lu M, Van der Wiele T, Wu MH, Sheng GY, Fu JM. Polybrominated diphenyl ethers in food and associated human daily intake assessment considering bioaccessibility measured by simulated gastrointestinal digestion. *Chemosphere* 2011; 83: 152-160.

- Zeng X, Simonich SLM, Robrock KR, Korytár P, Alvarez-Cohen L, Barofsky DF. Development and validation of a congener-specific photodegradation model for polybrominated diphenyl ethers. *Environ Toxicol Chem* 2008; 27: 2427-2435.
- Zhang B-Z, Zhang K, Li S-M, Wong CS, Zeng EY. Size-dependent dry deposition of airborne polybrominated diphenyl ethers in urban Guangzhou, China. *Environ Sci Technol* 2012b; 46: 7207-7214.
- Zhang K, Schnoor JL, Zeng EY. E-waste recycling: where does it go from here? *Environ Sci Technol* 2012a; 46: 10861-10867.
- Zhang T, Huang YR, Chen SJ, Liu AM, Xu PJ, Li N, Qi L, Ren Y, Zhou ZG, Mai BX. PCDD/Fs, PBDD/Fs, and PBDEs in the air of an e-waste recycling area (Taizhou) in China: current levels, composition profiles, and potential cancer risks. *J Environ Monit* 2012c; 14:3156-3163.
- Zhang W-H, Wu Y-X, Simonnot MO. Soil contamination due to e-waste disposal and recycling activities: a review with special focus on China. *Pedosphere* 2012d; 22: 434-455.
- Zhang X, Diamond ML, Robson M, Harrad S. Sources, emissions, and fate of polybrominated diphenyl ethers and polychlorinated biphenyls indoors in Toronto, Canada. *Environ Sci Technol* 2011a; 45: 3268-3274.
- Zhang XL, Luo XJ, Liu HY, Yu LH, Chen SJ, Mai BX. Bioaccumulation of several brominated flame retardants and Dechlorane Plus in waterbirds from an e-waste recycling region in south China: associated with trophic level and diet sources. *Environ Sci Technol* 2011b; 45: 400-405.
- Zhang Y, Ruan Y, Sun H, Zhao L, Gan Z. Hexabromocyclododecanes in surface sediments and a sediment core from rivers and harbor in the northern Chinese city of Tianjin. *Chemosphere* 2013; 90: 1610-1616.
- Zhao MM, Zhang SZ, Wang S, Huang HL. Uptake, translocation, and debromination of polybrominated diphenyl ethers in maize. *J Environ Sci* 2012; 24: 402-409
- Zheng XB, Wu JP, Luo X-J, Zeng Y-H, She Y-Z, Mai B-X. Halogenated flame retardants in home-produced eggs from an electronic waste recycling region in South China: levels, composition profiles, and human dietary exposure assessment. *Environ Int* 2012; 45: 122-128.
- Zhou P, Lin K, Zhou X, Zhang W, Huang K, Liu L, Guo J. Distribution of polybrominated diphenyl ethers in the surface sediments of the Taihu Lake, China. *Chemosphere* 2012; 88: 1375-1382.

FIGURE CAPTIONS

Figure 1. Concentration of BDE47 ($\mu\text{g kg}^{-1}\text{ lw}$) in human milk by year and country of collection (*results from human blood serum)

Figure 2. Significant continental differences were evident in maximal ΣPBDE concentrations ($\mu\text{g kg}^{-1}\text{ lw}$) measured in tissues (liver, eggs) of birds inhabiting freshwater ecosystems (Kruskal Wallis; $P = 0.02$). Concentrations were higher in freshwater birds from North America than those in birds from Asia or China, even after the removal of the highest levels reported by Henny and his colleagues (2011). Tissues were collected between 2005 and 2010. Only minor inconsistencies are expected to have occurred in the statistical results as a result of the varying number of BDE congeners (11 to 18) used to calculate the ΣBDE levels. Letters below each bar refer to the original study: A – Henny et al., 2011; B – Klosterhaus et al., 2012; C – Mora et al., 2012; D – Chen et al., 2012; E – Gentes et al., 2012; F – Morales et al., 2012; G – Mo et al., 2012; H – Zhang et al., 2011b; I – Wang et al., 2012.