

Horizontally acquired AT-rich genes in *Escherichia coli* cause toxicity by sequestering RNA polymerase

Lamberte, Lisa E.; Baniulyte, Gabriele; Singh, Shivani S.; Stringer, Anne M; Bonocora, Richard P.; Stracy, Mathew; Kapanidis, Achillefs N.; Wade, Joseph T.; Grainger, David C.

DOI:

[10.1038/nmicrobiol.2016.249](https://doi.org/10.1038/nmicrobiol.2016.249)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Lamberte, LE, Baniulyte, G, Singh, SS, Stringer, AM, Bonocora, RP, Stracy, M, Kapanidis, AN, Wade, JT & Grainger, DC 2017, 'Horizontally acquired AT-rich genes in *Escherichia coli* cause toxicity by sequestering RNA polymerase', *Nature Microbiology*, vol. 2, 16249. <https://doi.org/10.1038/nmicrobiol.2016.249>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 03/08/2017

Lamberte, Lisa E., et al. "Horizontally acquired AT-rich genes in *Escherichia coli* cause toxicity by sequestering RNA polymerase." *Nature microbiology* 2.3 (2017): nmicrobiol2016249.

doi:10.1038/nmicrobiol.2016.249

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1
2
3 **Horizontally acquired AT-rich genes in *Escherichia coli***
4 **cause toxicity by sequestering RNA polymerase**
5
6
7

8 Lisa E. Lamberte¹, Gabriele Baniulyte^{1†}, Shivani S. Singh¹, Anne M. Stringer², Richard P. Bonocora²,
9 Mathew Stracy³, Achillefs N. Kapanidis³, Joseph T. Wade^{2,4} and David C. Grainger^{1*}

10
11
12
13 *for correspondence

14 Email: d.grainger@bham.ac.uk Tel: +44 (0)121 4145437
15
16

17 ¹ Institute of Microbiology and Infection, School of Biosciences, University of Birmingham,
18 Edgbaston, Birmingham, B15 2TT, UK

19 ² Wadsworth Center, New York State Department of Health, Albany, NY, 12208, USA

20 ³ Biological Physics Research Group, Clarendon Laboratory, Department of Physics, University of
21 Oxford, Oxford OX1 3PU, UK

22 ⁴ Department of Biomedical Sciences, School of Public Health, University at Albany, SUNY, Albany,
23 NY, 12201, USA

24 [†] Present address: Department of Biomedical Sciences, School of Public Health, University at Albany,
25 SUNY, Albany, NY, 12201, USA
26
27
28
29

30 **ABSTRACT**

31 **Horizontal gene transfer permits rapid dissemination of genetic elements between individuals in**
32 **bacterial populations. Transmitted DNA sequences may encode favourable traits. However, if**
33 **acquired DNA has an atypical base composition, it can reduce host fitness. Consequently,**
34 **bacteria have evolved strategies to minimise the harmful effects of foreign genes. Most notably,**
35 **xenogeneic silencing proteins bind incoming DNA that has a higher AT-content than the host**
36 **genome. An enduring question has been to understand why such sequences are deleterious.**
37 **Here, we show that the toxicity of AT-rich DNA in *Escherichia coli* frequently results from**
38 **constitutive transcription initiation within the coding regions of genes. Left unchecked, this**
39 **causes titration of RNA polymerase and a global downshift in host gene expression.**
40 **Accordingly, a mutation in RNA polymerase that diminishes the impact of AT-rich DNA on**
41 **host fitness, reduces transcription from constitutive, but not activator-dependent, promoters.**

42

43 **INTRODUCTION**

44 Bacteria obtain DNA from their environment by direct uptake (transformation), the action of viruses
45 (transduction), or the acquisition of transmissible plasmids (conjugation)¹. Thus, “horizontal” DNA
46 transfer allows phenotypes to spread through bacterial populations. Transferred traits can be
47 beneficial¹. However, acquired sequences with a high AT-content may reduce host fitness²⁻⁶. Bacteria
48 have developed mechanisms to diminish the toxicity of AT-rich genes. One approach involves a
49 family of xenogeneic DNA binding proteins^{7,8}. These proteins target DNA that has a higher AT-
50 content than the host genome (“AT-rich” DNA), and silence transcription⁹⁻¹². However, it is unclear
51 why AT-rich DNA is toxic and why transcriptional silencing is beneficial.

52

53 The Histone-like Nucleoid Structuring (H-NS) protein of γ -proteobacteria is the best-characterised
54 xenogeneic silencing protein^{9,13}. To reduce transcription, H-NS oligomerises across AT-rich genes⁹⁻¹².
55 This process is triggered by nucleation at sequences containing a T:A step¹⁴⁻¹⁹. Intriguingly, promoter
56 -10 hexamers (consensus 5'-TATAAT-3', recognised by the RNA polymerase σ^{70} subunit) are
57 excellent inducers of H-NS nucleation^{15,20}. Hence, H-NS can exclude RNA polymerase from, or trap
58 RNA polymerase at, promoter DNA^{8,9,21,22}.

59

60 A challenge has been to understand why horizontally acquired AT-rich DNA is toxic. As we noted
61 previously, such regions contain a disproportionately high number of promoter-like sequences²³⁻²⁶.
62 Furthermore, we showed that H-NS suppresses transcription primarily from promoters that are
63 intragenic and/or far from gene starts²⁵. Here, we test the hypothesis that transcription from intragenic
64 promoters is a major cause of toxicity in cells lacking H-NS. Working with a sub-set of AT-rich
65 genes, we demonstrate dramatic effects of H-NS on intragenic transcription. When this transcription is
66 disabled, associated fitness costs decrease. Genome-wide, if derepressed, intragenic transcription
67 sequesters RNA polymerase. This causes a global downshift in canonical gene expression.

68 Accordingly, a mutation in RNA polymerase that reduces transcription from constitutive promoters,
69 but not those that are activator-dependent, compensates for loss of H-NS.

70

71 RESULTS

72 *Identification of the canonical yccE promoter*

73 We chose *yccE* as a model gene to understand xenogeneic silencing. We selected *yccE* because it is
74 AT-rich (66% A/T), silenced and bound by H-NS²⁷, and contains at least two intragenic
75 promoters^{25,28}. Importantly, *yccE* can be studied in isolation because the adjacent genes are not H-NS
76 bound (Figure 1a). We sought to understand the source of *yccE* transcription in cells lacking H-NS.
77 As a starting point, we identified the canonical *yccE* promoter. We were aided by previous
78 observations that *yccE* is a target for σ^{32} (heat shock σ factor)²⁹. Segments of the *yccE* intergenic
79 region were fused to *lacZ* in reporter plasmid pRW50 (Figure S1a). Promoter activity located to a 55
80 base pair (bp) DNA fragment immediately upstream of *yccE* (Figure S1b). Figure 1b shows the
81 sequence of this DNA fragment, named *yccE* Δ 200. A σ^{32} -dependent promoter sequence is apparent.
82 We monitored transcripts produced from this promoter (*PyccE*) *in vitro*. To do this, fragments from
83 the *yccE* intergenic region were cloned upstream of the *loop* terminator in plasmid pSR. Transcripts
84 initiating at *PyccE*, and terminating at *loop*, should be ~106 nucleotides (nt) in length. Coincidentally,
85 the RNA-I transcript, derived from the pSR replication origin, is 108/107 nt long. To avoid confusing
86 transcripts, we first monitored RNAs produced using pSR without a *PyccE* insert. The data show that
87 σ^{32} dependent synthesis of the RNA-I transcript is inefficient (Figure 1c, compare lanes 1 and 2).
88 When the entire *yccE* intergenic region was cloned in pSR σ^{32} dependent *yccE* transcripts were
89 identified (Figure 1c, lane 3). The same transcripts were observed upon cloning the truncated
90 *yccE* Δ 200 fragment (Figure 1c, lane 4). A 10 bp truncation at the 5' end of *yccE* Δ 200 abolished
91 transcription (Figure 1c, lane 5). To confirm correct identification of *PyccE*, DNA recognition
92 elements for σ^{32} were mutated (detailed in Figure 1b). All mutations greatly reduced promoter activity
93 (Figure 1d).

94

95 *Increased transcription of yccE in Δ hns cells does not require PyccE*

96 Convention dictates that *PyccE* should cause increased *yccE* transcription in cells lacking H-NS. We
97 considered this unlikely given the low activity, and σ^{32} dependence, of *PyccE*. Hence, we generated a
98 series of *yccE::lacZ* fusions to investigate the contribution of *PyccE*. The different constructs, labelled
99 *i* through *iv*, are illustrated below the graph in Figure 1e. We first measured *lacZ* expression in wild-
100 type cells (Figure 1e, grey bars). When *yccE* was in the forward orientation, low-level *lacZ* expression
101 was apparent. This was abolished upon deletion of *PyccE* (compare constructs *ii* and *iii*). In the
102 reverse orientation, *yccE* stimulated higher *lacZ* expression (construct *iv*). This is consistent with our
103 previous identification of a strong antisense promoter at the 5' end of *yccE* (adjacent to *lacZ* in this

104 assay)²⁸. Remarkably, in the absence of H-NS, *lacZ* expression increased in all scenarios (Figure 1e,
105 white bars). This transcription must be due to intragenic promoters.

106 *The yccE coding sequence is enriched for promoters*

107 A search identified 21 possible promoters within *yccE*. To test for function, we isolated promoters on
108 56 bp DNA fragments and fused them to *lacZ* (Figure 2a). Note that such sequences are too short to
109 be subject to repression by H-NS^{25,30}. Eleven DNA fragments stimulated β -galactosidase expression
110 twofold or more above background (labelled A-K in Figure 2a). Under the conditions of our
111 experiment, 7 of the 11 intragenic promoters were more efficient at driving transcription than *P_{yccE}*
112 (Figure 2a, black bar). Note that fragment “A” contains the strong antisense promoter described
113 previously²⁸.

114 *Transcription initiation within yccE is repressed by H-NS in vitro*

115 In the context of the full *yccE* gene, intragenic promoters should be repressed by H-NS. We tested this
116 *in vitro* using the pSR system. As noted above, with empty pSR, the 108/107 nt RNA-I transcript is
117 produced. In addition, larger transcripts are generated from genes native to the plasmid. The complete
118 set of transcripts produced from pSR is shown in Lane 1 of Figure 2b. In the figure, large (>1000 nt)
119 pSR derived transcripts are highlighted by a black box and RNA-I is highlighted by a grey box. We
120 introduced *yccE* into pSR, in either the forward or reverse orientation, upstream of the *loop*
121 terminator. Since *yccE* is 1257 bp in length, most intragenic transcripts should be separable from pSR
122 derived transcripts. As expected, numerous transcripts between 100 and 1000 nt in length were
123 detected for both *yccE* containing plasmids (compare Lanes 1, 2 and 6 in Figure 2b). For the plasmid
124 containing *yccE* in the reverse orientation an additional small transcript was detected (highlighted by
125 lower blue box in Figure 2b). This was expected, the transcript generated from the strong antisense
126 promoter at the 5' end of *yccE*, has a size of 90 nt in this assay. Regardless of *yccE* orientation,
127 addition of H-NS inhibited synthesis of most *yccE* derived transcripts (Lanes 3-5 and 6-9). Synthesis
128 of the RNA-I transcript was enhanced by H-NS suggesting that RNA polymerase is titrated by
129 promoters within *yccE*.

130 *Intragenic promoters are the source of increased yccE transcription in cells lacking H-NS*

131 To confirm that H-NS repressed intragenic *yccE* promoters *in vivo*, we made derivatives of our
132 *yccE::lacZ* fusions. The derivatives carry mutations in intragenic -10 elements. Schematics are below
133 the graph in Figure 2c. The open arrows show *yccE* lacking internal promoters. Full details are in
134 Figure S2. In the absence of H-NS, the ability of mutated *yccE* alleles to stimulate *lacZ* expression
135 was reduced (Figure 2c). Thus, both biochemical and genetic inspection show silencing of intragenic
136 transcription by H-NS. We refer to this as “pseudo-regulation” that occurs independently of, and may
137 be mistaken for, the control of mRNA synthesis. Thus, supposed gene regulatory effects of H-NS may
138 often be due to intragenic promoters.

139 *The fitness cost of yccE is a consequence of intragenic transcription*

140 We predicted a link between intragenic transcription and reduced fitness associated with loss of
141 H-NS. Hence, *E. coli* M182, and the *hns::kan* variant, were transformed with pSR carrying *yccE* with
142 or without internal promoters. Importantly, *yccE* mRNA cannot be expressed in this scenario; no
143 upstream promoter is present. We monitored cultures inoculated with these strains (Figure 2d,e). In all
144 cases, cells lacking *hns* had reduced fitness compared to the parent. However, this fitness defect was
145 smaller for the *yccE* derivative lacking internal promoters (Figure 2e). Complete elimination of the
146 fitness defect was not expected; AT-rich genes present on the *E. coli* chromosome also contribute.

147

148 *Repression of intragenic transcription by H-NS reduces the fitness cost of many AT-rich genes*

149 We identified other solitary genes targeted by H-NS: *yfdF*, *ykgH*, *yjgN* and *yjgL*. We cloned these, and
150 derivatives lacking intragenic promoters, upstream of the λ oop terminator in pSR. The *fepE* gene,
151 which has an AT-content of 55%, was included as a control. The constructs are illustrated above the
152 gel image in Figure 3a. Full gene sequences are in Figure S3. Figure 3a shows results of *in vitro*
153 transcription experiments. Whilst RNA polymerase did not initiate transcription within *fepE* (Lane 1),
154 intragenic transcription was observed for *yfdF*, *ykgH*, *yjgN* and *yjgL* (Lanes 2, 4, 6 and 8). Mutation of
155 intragenic promoters reduced this transcription (Lanes 3, 5, 7 and 9). The coding regions described
156 above were also cloned upstream of *lacZ* in pRW50 (Figure 3b). Expression of *lacZ* was measured in
157 M182 or the *hns::kan* derivative (Figure 3b). Upon deletion of *hns*, expression of *lacZ* downstream of
158 *yfdF*, *ykgH*, *yjgN* or *yjgL*, but not *fepE*, increased (solid arrows). In contrast, no such increase
159 occurred when intragenic promoters were mutated (open arrows). These analyses are consistent with
160 silencing of intragenic promoters by H-NS at all loci tested. Hence, we measured the fitness cost of
161 multicopy *yfdF*, *ykgH*, *yjgN* and *yjgL* in cells with and without H-NS. In all cases, the fitness deficit
162 between wild-type and *hns::kan* cells decreased upon mutation of intragenic promoters (Figure 3c,d).
163 Furthermore, changes in fitness and *lacZ* expression were significantly correlated (Figure S4a).

164 *AT-rich genes titrate RNA polymerase and cause a global downshift in housekeeping transcription*

165 RNA polymerase levels are limited in *E. coli*³¹. Consequently, multiple copies of a strong promoter
166 can hinder growth and titrate transcription of the *lac* operon (Figure S4b). Interestingly, RNA
167 polymerase levels do not increase in cells lacking H-NS (Figure S5). Therefore, competition for the
168 enzyme must increase; more promoters compete for a limited supply of RNA polymerase. Logically,
169 migration of RNA polymerase to spurious promoters should cause a global downshift in canonical
170 transcription. However, despite many studies of the H-NS controlled transcriptome, a universal
171 downshift has never been reported^{9,10,27,32,33}. We reasoned that this might result from data
172 normalisation approaches used previously. Briefly, transcriptome analysis compares RNA levels in
173 two strains. Comparison requires a point of reference believed to be consistent between the strains.
174 For example, it may be assumed that housekeeping genes are similarly transcribed or that averaged

175 transcription across all genes will be equivalent. Problematically, these approaches cannot
176 differentiate between technical variation and genuine shifts in global transcription. To circumvent this
177 problem, spiked-in RNA standards can be used³⁴. We adapted this tactic to quantify global effects of
178 H-NS on transcription. Briefly, we grew *E. coli* MG1655, and the *hns::kan* variant, to mid-log phase.
179 We then counted colony-forming units for each *E. coli* culture. The same was done for a single culture
180 of *Salmonella* Typhimurium. Accordingly, we were able to mix a defined number of clonal
181 *S. Typhimurium* cells with each *E. coli* strain. The cell mixtures were subject to transcriptome
182 analysis. For identically grown *S. Typhimurium* cells, differences in RNA abundance must result from
183 processing variation. Hence, at the end of the procedure, transcripts mapping uniquely to
184 *S. Typhimurium* were used to normalise the data. Figure 4a shows a post-normalisation plot of read
185 depth for each *E. coli* gene in each strain. The diagonal blue line shows the expected position of data
186 points if transcription is unchanged between strains. There is a downshift in transcription of genes not
187 bound by H-NS (black). Conversely, H-NS bound genes (red) are transcribed more frequently. This
188 behaviour is exemplified in Figure 4b.

189 *RNA polymerase titration can be visualised directly*

190 Genes bound by H-NS are overrepresented near the chromosome replication terminus (Ter)^{35,36}.
191 Conversely, most RNA polymerase binds near the origin of replication (Ori)³⁷. These loci occupy
192 distinct intracellular territories; Ter typically frequents mid-cell whilst Ori migrates to the poles³⁷⁻³⁹.
193 Consequently, it should be possible to visualise titration of RNA polymerase at the cellular level.
194 Previously, we used super resolution microscopy to track individual RNA polymerase molecules in
195 live *E. coli*⁴⁰. Here, we repeated this analysis to examine the effect of deleting *hns*. In wild-type cells,
196 RNA polymerase clusters near the quarter cell positions (Figure 4c, top). However, in cells lacking
197 H-NS, RNA polymerase is redistributed and mid-cell is occupied (Figure 4c, middle). Consistent with
198 unaltered RNA polymerase abundance, occupancy of mid-cell corresponds to reduced RNA
199 polymerase abundance elsewhere (Figure 4c).

200

201 *RNA polymerase mutation can compensate for loss of H-NS by favouring regulated transcription*

202 An aspartic acid substitution in the RNA polymerase σ^{70} subunit (G424D in *E. coli*) can compensate
203 for loss of H-NS⁴¹. This side chain could clash with the promoter -10 element during transcription
204 initiation⁴¹. According to our model, promoters within genes are constitutive; they rely solely on their
205 DNA sequence to bind RNA polymerase. Conversely, canonical promoters are regulated; a complex
206 array of transcriptional activators stabilise RNA polymerase binding and DNA unwinding⁴². We
207 reasoned that transcriptional defects, due to the G424D mutation, might be more pronounced at
208 constitutive promoters. To test this, we purified RNA polymerase containing σ^{70} or the G424D
209 derivative. We then investigated the ability of the RNA polymerase derivatives to stimulate
210 unwinding of a promoter -10 element using KMnO₄ footprinting. The semi-synthetic NM501

211 promoter was used because it has constitutive activity, by virtue of a consensus -10 element, but can
212 be upregulated by the transcriptional activator CRP⁴³. The G424D mutation completely abolished
213 DNA opening by RNA polymerase in the absence of CRP (Figure 4d, compare lanes 2-5 and 6-9). In
214 contrast, when CRP was present, the σ^{70} G424D mutation had little effect (compare lanes 10-13 and
215 14-17). Similarly, *in vivo*, the σ^{70} G424D mutation caused transcriptional defects at 7 of 8 constitutive
216 promoters (Figure 4e). However, σ^{70} G424D functioned as well as, or better than, wild type RNA
217 polymerase at activator dependent promoters (Figure 4f).

218

219 DISCUSSION

220 The AT-rich genes examined here impose a fitness cost due to intragenic promoters. This
221 phenomenon is likely to be widespread in bacteria; functional homologues of H-NS are apparent in
222 diverse species⁸. Furthermore, the DNA binding properties of RNA polymerase are highly
223 conserved⁴². We suggest that misappropriation of cellular resources underlies the *hns* phenotype.
224 Redeployment of the finite RNA polymerase pool causes uniform suppression of canonical
225 transcription. Whilst this general effect is likely to be pervasive, we do not exclude organism-specific
226 complications. For example, in *Pseudomonas aeruginosa*, loss of H-NS-like proteins causes prophage
227 induction and cell death^{44,45}. Interestingly, linear H-NS filaments do not pose a barrier to transit of
228 RNA polymerase⁴⁶. Accordingly, transcription of an mRNA, and silencing of intragenic promoters,
229 could occur simultaneously. For example, in this work, *PycC* was active both in isolation and
230 upstream of H-NS bound *yccE*. We speculate that gene silencing by H-NS may have evolved to
231 discriminate between canonical and spurious RNA synthesis.

232

233 MATERIALS AND METHODS

234 *Strains, plasmids and general methods*

235 *E. coli* JCB387 $\Delta nir \Delta lac$ and MG1655 have been described previously^{47,48}. M182*hns::kan* and
236 KF26*hns::kan* were constructed by P1 transduction of *hns::kan* from MG1655 into the respective
237 parent strains. The MG1655 *hns::kan* strain was provided by Ding Jin. M182*rpoD::kan* was generated
238 using gene doctoring according to the protocol of Lee *et al*⁴⁹ using the plasmids and oligonucleotides
239 listed in Table S1. Note that, prior to gene doctoring, M182 strains were transformed with plasmid
240 pVR σ that encodes *rpoD*⁵⁰. Quickchange mutagenesis was used to introduce the G424D mutation into
241 pVR σ encoded *rpoD* (Table S1). Fortuitously, chromosomal and plasmid encoded σ^{70} were produced
242 at indistinguishable levels (Figure S6). Sample sizes for all experiments were selected to ensure
243 reproducibility in line with our previous work.

244 *DNA fragments and gene expression assays*

245 Promoter::*lacZ* fusions were made by cloning DNA fragments upstream of *lacZ* in the low copy
246 number plasmid pRW50⁵¹. The nested deletions in the *yccE* intergenic region were generated by PCR

247 and oligonucleotides shown in Table S1. The various *yccE*, *yfdF*, *ykgH*, *yjgN* and *yjgL* alleles were
248 synthesised by Invitrogen and some contain silent mutations to remove *EcoRI* or *HindIII* restriction
249 sites to facilitate cloning (Figures S2a and S3). Oligonucleotides used to amplify the different alleles
250 for cloning into pSR and pRW50 are shown in Table S1. The 56 bp intragenic *yccE* fragments were
251 generated with overlapping oligonucleotides (Table S1). The resulting single stranded overhangs were
252 filled with DNA polymerase before cloning. β -galactosidase assays were done using the protocol of
253 Miller⁵². All assay values are the mean of three biological replicates and the error bars show standard
254 deviation from the mean. Experiments were done at least twice. Cells were grown aerobically at 37°C,
255 to mid-log phase, in LB media.

256 *Bioinformatic analysis of genes and design of new coding regions*

257 Our stringent search criteria selected putative σ^{70} dependent promoters as described previously²⁵.
258 Thus, sequences were selected that matched the motifs 5'-TAnAAT-3', 5'-TATnAT-3' or 5'-TATAnT-
259 3'. The relaxed search selected the sequence 5'-TAnnnT-3'. To inactivate promoter -10 elements the
260 initial 5'-TA-3' was replaced with 5'-GG-3'

261 *Proteins, KMnO₄ footprinting and in vitro transcription assays*

262 H-NS and RNA polymerase were prepared as described previously⁵³. DNA fragments for KMnO₄
263 footprinting experiments were derived from Qiagen maxi-preparations of plasmid pSR. Thus,
264 promoter DNA fragments were excised from pSR by sequential digestion with *HindIII* and then *AatII*.
265 After digestion fragments were labelled at the *HindIII* end using [γ -³²P]-ATP and polynucleotide
266 kinase. Footprints were done as described by Grainger *et al.*⁵⁴. The *in vitro* transcription experiments
267 were done as described by Savery *et al.*⁵⁵ using the system of Kolb *et al.*⁵⁶. A Qiagen maxiprep kit
268 was used to purify supercoiled pSR plasmid carrying the different promoter inserts. This template (16
269 $\mu\text{g ml}^{-1}$) was pre-incubated with purified H-NS in buffer containing 20 mM Tris pH 7.9, 5 mM
270 MgCl₂, 500 μM DTT, 50 mM KCl, 100 $\mu\text{g ml}^{-1}$ BSA, 200 μM ATP, 200 μM GTP, 200 μM CTP, 10
271 μM UTP with 5 μCi [α -³²P]-UTP. The reaction was started by adding purified *E. coli* RNA
272 polymerase. Labelled RNA products were analysed on a denaturing polyacrylamide gel. All *in vitro*
273 assays were repeated at least three times in their entirety.

274 *Growth assays*

275 Cells lacking H-NS rapidly acquire compensatory mutations⁴¹. Consequently, reproducible changes in
276 growth were only obtained when precautions were taken to minimise this phenomenon. The primary
277 precaution was to reduce the number of division cycles that strains passed through during experimental
278 setup. Thus, M182 and the *hns::kan* derivative were taken directly from long-term -80°C storage and
279 used immediately to inoculate LB medium. After incubation for several hours at 37°C cells were
280 harvested and competency was induced using ice cold CaCl₂. The cells were then transformed with
281 desired plasmids and transformants were isolated on selective agar plates. A colony from each plate
282 was suspended in LB medium and aliquots of this were used immediately to inoculate fresh media so

283 that growth could be monitored. Cells were grown either in LB medium at 37°C or in M9 minimal
284 medium at 30°C. Values shown are from three biological replicates and the experiments were done on
285 two separate occasions.

286 *Western blotting*

287 To determine relative protein levels in different strains cells were grown in LB media at 37°C.
288 Aliquots of the culture were harvested at indicated time points and CFUs determined. Following this
289 quantification the same number of cells from each aliquot were resuspended in SDS-PAGE gel
290 loading buffer. After heating to 90°C for two minutes we separated proteins present in each lysate
291 using SDS-PAGE. The proteins were then transferred to a Hybond-ECL nitrocellulose membrane
292 using an Invitrogen XCell II Blot Module. Mouse anti-sera against the σ^{70} and β subunits of RNA
293 polymerase (Neoclone) and H-NS (a gift from Jay Hinton) was used to detect the relevant proteins.
294 Primary antibody binding was probed with horseradish peroxidase-linked rabbit anti-mouse antisera
295 (Sigma-Aldrich A9044). The experiments were done on two separate occasions.

296 *Standard RNA-seq*

297 RNA-seq experiments were done in duplicate. *E. coli* MG1655 and *E. coli* MG1655 Δhns were grown
298 in LB medium at 30°C to an OD₆₀₀ of 0.5-0.7. RNA was harvested, RNA-seq libraries were
299 constructed and libraries were sequenced as described previously⁵⁷.

300 *RNA-seq with a spiked in control*

301 Transcriptome analysis experiments shown were performed in duplicate using *E. coli* MG1655 and *E.*
302 *coli* MG1655 Δhns . We also ran the same experiment, again in duplicate, using *E. coli* M182 and the
303 *hns::kan* derivative. The results were near identical. For each replicate three mid-log phase bacterial
304 cultures were prepared. The three cultures were *E. coli* MG1655, *E. coli* MG1655 Δhns or *Salmonella*
305 Typhimurium 14028s in LB medium. Before harvesting RNA the number of colony forming units
306 (CFUs) per unit volume was determined for each culture by plating dilutions of each culture on
307 nutrient agar and counting bacterial colonies. Aliquots of the two *E. coli* cultures were mixed *S.*
308 Typhimurium cells. The volume of *Salmonella* cells was normalised to the CFUs for the
309 corresponding *E. coli* culture. This step is crucial because it allows processing artefacts, due to
310 differences in lysis efficiency, RNA recovery or cDNA synthesis, to be removed. Thus, in the final
311 transcriptome analysis, the number of sequencing reads corresponding to the *S. Typhimurium* genome
312 should be identical for each sample; they were derived from the same number of clonal *S.*
313 Typhimurium cells. Hence, differences can only result from downstream processing. RNA was
314 harvested, and RNA-seq libraries were prepared, as described previously⁵⁷. RNA-seq libraries were
315 sequenced on an Illumina Hi-Seq 2000 instrument (University at Buffalo, SUNY, Buffalo, NY,
316 USA).

317 *Data normalisation and transcriptome analysis*

318 Our normalisation procedure is based on the addition of a proportional number of *S. Typhimurium*
319 cells to each sample of *E. coli* cells immediately before harvesting RNA. All reads were first mapped
320 to the *E. coli* MG1655 genome using CLC Genomics Workbench (Version 8.0; default parameters
321 except that a perfect match was required). Unmapped reads were mapped to the *S. Typhimurium*
322 14028s genome (same parameters as above). The number of mapped *S. Typhimurium* reads, for each
323 *E. coli* sample, was used to determine a correction factor for each sample. For example, if Sample A
324 has twice as many *S. Typhimurium* reads as Sample B, the correction factor for Sample A will be
325 twice that for Sample B. Having calculated a correction factor for each sample, we remapped all
326 sequence reads to the *S. Typhimurium* 14028s genome using CLC Genomics Workbench (same
327 parameters as above). Unmapped reads were then mapped to the *E. coli* MG1655 genome using CLC
328 Genomics Workbench (same parameters as above). Total read coverage per gene was calculated using
329 a custom Python script. These values were normalised to the length of the gene, and further
330 normalised using the correction factor (described above). Raw data are available from the
331 ArrayExpress database using accession number E-MTAB-4751.

332 *Super resolution microscopy of RNA polymerase*

333 Cell preparation, microscopy, and analysis were performed as described previously^{40,58}. In brief, we
334 used an endogenous fusion of photoactivatable fluorescent protein PAmCherry, with the β' subunit of
335 RNA polymerase, encoded by *E. coli* strain KF26⁵⁸. Glycerol stocks of KF26, and the *hns::kan*
336 derivative, were used to inoculate fresh Rich Defined Media (RDM, Teknova). When the culture
337 attained an OD₆₅₀ value of ~0.2 cells were collected by centrifugation and resuspended. Cells in 1 μ l
338 of the suspension were placed on an RDM agarose pad and imaged for 300000 frames at 15 ms
339 exposure on a custom built single-molecule TIRF microscope. For each strain, 9 fields of view were
340 imaged. The experiment was done on 3 separate occasions. Molecules were imaged by
341 photoactivating and localising fluorophores, and joining localisations over multiple frames to obtain
342 trajectories of individual molecules. To measure RNA polymerase mobility, we calculated an
343 apparent diffusion coefficient from the mean squared displacement of trajectories of individual
344 molecules, and used a threshold to distinguish transcriptionally active and promoter bound molecules
345 from the rest of the population. Cell outlines were determined from the brightfield image, and the
346 average intracellular location of RNA polymerase was established from a 2D histogram of the mean
347 trajectory positions relative to the cell outline for all DNA-bound molecules in at least 100 cells. Ter
348 positioning was determined using similar analysis of strain PZ111 carrying a *tetO* array inserted at
349 Ter, and expressing a TetR-mYpet fusion. The strain was constructed by P1 transduction of *tetR*-
350 mYPet kan, into an AB1157 strain with an array of 240 *tetO* sequences 50 kb clockwise of *dif* (*ter3*)⁵⁹.

351 *Data availability*

352 The data that support these findings are available from the corresponding author upon request. The
353 raw data for RNA-seq experiments are available from the ArrayExpress database using accession
354 number E-MTAB-4751. Original gel images are shown in Figure S7.

355 **ACKNOWLEDGEMENTS**

356 This work was funded by Leverhulme Trust project grant (RPG-2013-147) and Wellcome Trust
357 Career Development Fellowship (WT085092MA) awarded to D.C.G. Support for J.T.W. was a
358 National Institutes of Health Director's New Innovator Award (1DP2OD007188). A.N.K. and M.S.
359 were supported by BBSRC grant (BB/N018656/1; to ANK and MS), and a Wellcome Trust
360 Investigatorship (110164/Z/15/Z; to ANK). We thank Jay Hinton for the gift of anti-H-NS anti-sera.

361 **CONTRIBUTIONS**

362 D.C.G. and J.T.W. designed the study and wrote the manuscript. L.E.L., G.B., S.S.S., A.M.S., R.P.B.,
363 and M.S. generated the data and prepared it for publication. M.S. and A.N.K. provided new analytical
364 tools and critically discussed the manuscript with D.C.G. and J.T.W. All authors contributed to data
365 analysis and interpretation.

366 **REFERENCES**

- 367 1. Soucy SM, Huang J, Gogarten JP. 2015. Horizontal gene transfer: building the web of life. *Nat Rev*
368 *Genet* **16**: 472-482.
- 369 2. Doyle M, Fookes M, Ivens A, Mangan MW, Wain J, Dorman CJ. 2007. An H-NS-like stealth
370 protein aids horizontal DNA transmission in bacteria. *Science* **315**: 251-252.
- 371 3. Popa O, Hazkani-Covo E, Landan G, Martin W, Dagan T. 2011. Directed networks reveal genomic
372 barriers and DNA repair bypasses to lateral gene transfer among prokaryotes. *Genome Res* **21**: 599-
373 609.
- 374 4. Popa O, Dagan T. 2011. Trends and barriers to lateral gene transfer in prokaryotes. *Curr Opin*
375 *Microbiol* **145**: 615-623
- 376 5. Raghavan R, Kelkar YD, Ochman H. 2012. A selective force favoring increased G+C content in
377 bacterial genes. *Proc Natl Acad Sci* **109**: 14504-14507.
- 378 6. Baltrus DA. 2013. Exploring the costs of horizontal gene transfer. *Trends Ecol Evol* **8**: 489-495.
- 379 7. Dorman CJ. 2007. H-NS, the genome sentinel. *Nat Rev Microbiol.* **5**: 157-61.
- 380 8. Singh K, Milstein JN, Navarre WW. 2016. Xenogeneic Silencing and Its Impact on Bacterial
381 Genomes. *Annu Rev Microbiol.* **70**: 199-213.
- 382 9. Navarre WW, Porwollik S, Wang Y, McClelland M, Rosen H, Libby SJ, Fang FC. 2006. Selective
383 silencing of foreign DNA with low GC content by the H-NS protein in Salmonella. *Science* **313**: 236-
384 238.

- 385 10. Lucchini S, Rowley G, Goldberg MD, Hurd D, Harrison M, Hinton JC. 2006. H-NS mediates the
386 silencing of laterally acquired genes in bacteria. *PLoS Pathog* **28**: e81.
- 387 11. Smits WK, Grossman AD. 2010. The transcriptional regulator Rok binds A+T-rich DNA and is
388 involved in repression of a mobile genetic element in *Bacillus subtilis*. *PLoS Genet* **6**: e1001207
- 389 12. Gordon BR, Li Y, Wang L, Sintsova A, van Bakel H, Tian S, Navarre WW, Xia B, Liu J. 2010.
390 Lsr2 is a nucleoid-associated protein that targets AT-rich sequences and virulence genes in
391 *Mycobacterium tuberculosis*. *Proc Natl Acad Sci* **107**: 5154-5159.
- 392 13. Dorman CJ. 2014. H-NS-like nucleoid-associated proteins, mobile genetic elements and
393 horizontal gene transfer in bacteria. *Plasmid* **75**: 1-11.
- 394 14. Bouffartigues E, Buckle M, Badaut C, Travers A, Rimsky S. 2007. H-NS cooperative binding to
395 high-affinity sites in a regulatory element results in transcriptional silencing. *Nat Struct Mol Biol* **14**:
396 441-418.
- 397 15. Gordon BR, Li Y, Cote A, Weirauch MT, Ding P, Hughes TR, Navarre WW, Xia B, Liu J. 2011.
398 Structural basis for recognition of AT-rich DNA by unrelated xenogeneic silencing proteins. *Proc*
399 *Natl Acad Sci* **108**: 10690-10695.
- 400 16. Arold ST, Leonard PG, Parkinson GN, Ladbury JE. 2010. H-NS forms a superhelical protein
401 scaffold for DNA condensation. *Proc Natl Acad Sci* **107**: 15728-15732.
- 402 17. Amit R, Oppenheim AB, Stavans J. 2003. Increased bending rigidity of single DNA molecules by
403 H-NS, a temperature and osmolarity sensor. *Biophys J* **84**: 2467-73.
- 404 18. Dame RT, Noom MC, Wuite GJ. 2006. Bacterial chromatin organization by H-NS protein
405 unravelled using dual DNA manipulation. *Nature*. **444**: 387-390.
- 406 19. Liu Y, Chen H, Kenney LJ, Yan J. 2010. A divalent switch drives H-NS/DNA-binding
407 conformations between stiffening and bridging modes. *Genes Dev*. **24**:339-44.
- 408 20. Landick R, Wade JT, Grainger DC. 2015. H-NS and RNA polymerase: a love-hate relationship?
409 *Curr Opin Microbiol* **24**: 53-59.
- 410 21. Winardhi RS, Yan J, Kenney LJ. 2015. H-NS Regulates Gene Expression and Compacts the
411 Nucleoid: Insights from Single-Molecule Experiments. *Biophys J*. **109**: 1321-1329.
- 412 22. Dame RT, Wyman C, Wurm R, Wagner R, Goosen N. 2002. *J Biol Chem* **277**: 2146-2150.
- 413 23. Huang Q, Cheng X, Cheung MK, Kiselev SS, Ozoline ON, Kwan HS. 2012. High-density
414 transcriptional initiation signals underline genomic islands in bacteria. *PLoS One* **7**: e33759.
- 415 24. Singh SS and Grainger DC. 2013. H-NS can facilitate specific DNA-binding by RNA polymerase
416 in AT-rich gene regulatory regions. *PLoS Genet* **9**: e1003589.
- 417 25. Singh SS, Singh N, Bonocora RP, Fitzgerald DM, Wade JT, Grainger DC. 2014. Widespread
418 suppression of intragenic transcription initiation by H NS *Genes Dev* **28**: 214-219.
- 419 26. Lam KN and Charles TC. 2015. Strong spurious transcription likely contributes to DNA insert
420 bias in typical metagenomic clone libraries. *Microbiome* **3**: 22.

- 421 27. Kahramanoglou C, Seshasayee AS, Prieto AI, Ibberson D, Schmidt S, Zimmermann J, Benes V,
422 Fraser GM, Luscombe NM. 2011. Direct and indirect effects of H-NS and Fis on global gene
423 expression control in *Escherichia coli*. *Nucleic Acids Res* **39**: 2073-91.
- 424 28. Chintakayala K, Singh SS, Rossiter AE, Shahapure R, Dame RT, Grainger DC. 2013. *E. coli* Fis
425 protein insulates the *cbpA* gene from uncontrolled transcription. *PLoS Genet* **9**: e1003152.
- 426 29. Wade JT, Castro Roa D, Grainger DC, Hurd D, Busby SJ, Struhl K, Nudler E. 2006. Extensive
427 functional overlap between sigma factors in *Escherichia coli*. *Nat Struct Mol Biol*. **13**: 806-814.
- 428 30. Haycocks JR, Sharma P, Stringer AM, Wade JT, Grainger DC. 2015. The molecular basis for
429 control of ETEC enterotoxin expression in response to environment and host. *PLoS Pathog*
430 **11**:e1004605.
- 431 31. Piper SE, Mitchell JE, Lee DJ, Busby SJ. 2009. A global view of *Escherichia coli* Rsd protein and
432 its interactions. *Mol Biosyst* **5**: 1943-1947.
- 433 32. Srinivasan R, Scolari VF, Lagomarsino MC, Seshasayee AS. 2015. The genome-scale interplay
434 amongst xenogene silencing, stress response and chromosome architecture in *Escherichia coli*.
435 *Nucleic Acids Res*. **43**: 295-308.
- 436 33. Oshima T, Ishikawa S, Kurokawa K, Aiba H, Ogasawara N. 2006. *Escherichia coli* histone-like
437 protein H-NS preferentially binds to horizontally acquired DNA in association with RNA polymerase.
438 *DNA Res* **13**: 141-53.
- 439 34. Lovén J, Orlando DA, Sigova AA, Lin CY, Rahl PB, Burge CB, Levens DL, Lee TI, Young RA.
440 2012. Revisiting global gene expression analysis. *Cell* **151**: 476-482.
- 441 35. Lawrence JG, Ochman H. 1998. Molecular archaeology of the *Escherichia coli* genome. *Proc.*
442 *Natl. Acad. Sci. USA*. **95**: 9413-9417.
- 443 36. Zarei M, Sclavi B, Cosentino Lagomarsino M. 2013. Gene silencing and large-scale domain
444 structure of the *E. coli* genome. *Mol Biosyst*. **9**: 758-767.
- 445 37. Dame RT, Kalmykova OJ, Grainger DC. 2011. Chromosomal macrodomains and associated
446 proteins: implications for DNA organization and replication in Gram negative bacteria. *PLoS Genet*.
447 **7**: e1002123.
- 448 38. Junier I, Boccard F, Espéli O. 2014. Polymer modeling of the *E. coli* genome reveals the
449 involvement of locus positioning and macrodomain structuring for the control of chromosome
450 conformation and segregation. *Nucleic Acids Res*. **42**: 1461-1473.
- 451 39. Youngren B, Nielsen HJ, Jun S, Austin S. 2014. The multifork *Escherichia coli* chromosome is a
452 self-duplicating and self-segregating thermodynamic ring polymer. *Genes Dev*. **28**: 71-84.
- 453 40. Stracy M, Lesterlin C, Garza de Leon F, Uphoff S, Zawadzki P, Kapanidis AN. 2015. Live-cell
454 superresolution microscopy reveals the organization of RNA polymerase in the bacterial nucleoid.
455 *Proc Natl Acad Sci USA*. **112**: E4390-9.
- 456 41. Ali SS, Soo J, Rao C, Leung AS, Ngai DH, Ensminger AW, Navarre WW. 2014. Silencing by H-
457 NS potentiated the evolution of *Salmonella*. *PLoS Pathog* **10**: e1004500.

- 458 42. Lee DJ, Minchin SD, Busby SJ. 2012. Activating transcription in bacteria. *Annu Rev Microbiol.*
459 **66**: 125-152.
- 460 43. Miroslavova NS and Busby SJ. 2006. Investigations of the modular structure of bacterial
461 promoters. *Biochem Soc Symp* **73**: 1-10.
- 462 44. Castang S, McManus HR, Turner, KH, Dove, SL. 2008. H-NS family members function
463 coordinately in an opportunistic pathogen. *Proc. Natl. Acad. Sci. USA* **105**: 18947-18952.
- 464 45. Li C, Wally H, Miller SJ, Lu CD. 2009. The Multifaceted Proteins MvaT and MvaU, Members of
465 the H-NS Family, Control Arginine Metabolism, Pyocyanin Synthesis, and Prophage Activation in
466 *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **191**: 6211-6218.
- 467 46. Kotlajich MV, Hron DR, Boudreau BA, Sun Z, Lyubchenko YL, Landick R. 2015. Bridged
468 filaments of histone-like nucleoid structuring protein pause RNA polymerase and aid termination in
469 bacteria. *Elife* **4**: e04970.
- 470 47. Page L, Griffiths L, Cole JA. 1990. Different physiological roles of two independent pathways for
471 nitrite reduction to ammonia by enteric bacteria. *Arch Microbiol* **154**: 349-354.
- 472 48. Keseler IM, Mackie A, Peralta-Gil M, Santos-Zavaleta A, Gama-Castro S, Bonavides-Martínez C,
473 Fulcher C, Huerta AM, Kothari A, Krummenacker M, *et al.* 2013. EcoCyc: fusing model organism
474 databases with systems biology. *Nucleic Acids Res* **41**: D605-12.
- 475 49. Lee, DJ, Bingle LE, Heurlier K, Pallen MJ, Penn CW, Busby SJ, Hobman JL. 2009. Gene
476 doctoring: a method for recombineering in laboratory and pathogenic *Escherichia coli* strains. *BMC*
477 *Microbiol* **9**: 252.
- 478 50. Rhodius VA, Busby SJ. 2000. Interactions between activating region 3 of the *Escherichia coli*
479 cyclic AMP receptor protein and region 4 of the RNA polymerase sigma70 subunit: application of
480 suppression genetics. *J Mol Biol* **299**: 311-324.
- 481 51. Lodge J, Fear J, Busby S, Gunasekaran P, Kamini NR. 1992. Broad host range plasmids carrying
482 the *Escherichia coli* lactose and galactose operons. *FEMS Microbiol Lett.* **74**: 271-276.
- 483 52. Miller J 1972. Experiments in Molecular Genetics. Cold Spring Harbor, NY: Cold Spring Harbor
484 Laboratory Press.
- 485 53. Grainger DC, Goldberg MD, Lee DJ, Busby SJ. 2008. Selective repression by Fis and H-NS at the
486 *Escherichia coli* *dps* promoter. *Mol Microbiol* **68**: 1366-1377.
- 487 54. Grainger DC, Belyaeva TA, Lee DJ, Hyde EI, Busby SJ. 2004. Transcription activation at the
488 *Escherichia coli* *melAB* promoter: interactions of MelR with the C-terminal domain of the RNA
489 polymerase alpha subunit. *Mol Microbiol* **51**: 1311-1320.
- 490 55. Savery NJ, Lloyd GS, Kainz M, Gaal T, Ross W, Ebright RH, Gourse RL, Busby SJ. *et al.* 1998.
491 Transcription activation at Class II CRP-dependent promoters: identification of determinants in the C-
492 terminal domain of the RNA polymerase alpha subunit. *EMBO J* **17**: 3439-3447.
- 493 56. Kolb A, Kotlarz D, Kusano S, Ishihama A 1995. Selectivity of the *Escherichia coli* RNA
494 polymerase E sigma 38 for overlapping promoters and ability to support CRP activation. *Nucleic*
495 *Acids Res* **23**: 819-826.

- 496 57. Stringer AM, Currenti S, Bonocora RP, Baranowski C, Petrone BL, Palumbo MJ, Reilly AA,
497 Zhang Z, Erill I, Wade JT. 2014. Genome-scale analyses of *Escherichia coli* and *Salmonella enterica*
498 AraC reveal noncanonical targets and an expanded core regulon. *J Bacteriol* **196**: 660-671.
- 499 58. Endesfelder U, Finan K, Holden SJ, Cook PR, Kapanidis AN, Heilemann M. 2013. Multiscale
500 spatial organization of RNA polymerase in *Escherichia coli*. *Biophys J*. **105**: 172-181.
- 501 59. Zawadzki P, Stracy M, Ginda K, Zawadzka K, Lesterlin C, Kapanidis AN, Sherratt DJ. 2015. The
502 Localization and Action of Topoisomerase IV in *Escherichia coli* Chromosome Segregation Is
503 Coordinated by the SMC Complex, MukBEF. *Cell Rep*. **13**: 2587-2596.

504

505 FIGURES

506 **Figure 1: Characterisation of the *yccE* locus.**

507 **a) Genomic context of *yccE* and its promoter.** The panel shows *yccE*, and surrounding genes,
508 alongside data describing H-NS binding (ChIP-seq²⁷) and RNA abundance (standard RNA-seq; this
509 work, done in duplicate). Data are representative.

510 **b) Sequence of the *yccE*Δ200 DNA fragment containing *PyccE*.** The *PyccE* -10 and -35 elements
511 are in bold and underlined. A consensus σ^{32} promoter sequence is in grey for comparison. Mutations
512 made to disrupt sequence elements are in red. Transcription can initiate at adjacent nucleotides. These
513 are labelled (+1) and are highlighted by a bent arrow.

514 **c) Analysis of transcripts generated from *PyccE* *in vitro*.** The gel image shows transcripts
515 generated by RNA polymerase, associated with either σ^{70} or σ^{32} , from pSR plasmid DNA templates.
516 The schematic diagrams above the gel image represent the native cloning site of pSR (left hand side)
517 or derivatives containing a *PyccE* insert (right hand side). The different *PyccE* containing DNA
518 fragments inserted are indicated below the gel image in parenthesis. *PyccE* derived transcripts
519 manifest as a 107/106 nucleotide (nt) doublet. The 108/107 nt RNA-I transcript is derived from the
520 plasmid replication origin. The experiment was done three times. Data are representative.

521 **d) Effect of *PyccE* mutation *in vivo*.** The graph shows LacZ activity data obtained from *E. coli*
522 JCB387 cells carrying different *yccE*Δ200 derivatives cloned in pRW50. Assays were done in
523 triplicate and error bars show standard deviation from the mean.

524 **e) Induction of *yccE* transcription in the absence of H-NS does not require *PyccE* *in vivo*.** The
525 panel illustrates a series of *yccE::lacZ* fusions labelled *i-iv*. Genes are shown as block arrows and
526 *PyccE* is shown as a bent line arrow. The β -galactosidase activity was measured in lysates of M182
527 (grey bars) and M182*hns::kan* cells (open bars). Assays were done in triplicate and error bars show
528 standard deviation from the mean.

529

530 **Figure 2: H-NS represses intragenic *yccE* transcription and associated fitness costs**

531 **a) Identification of intragenic *yccE* promoters.** The data are β -galactosidase activities driven by
532 short intragenic *yccE* DNA fragments in strain JCB387. The bars align with the location of the DNA
533 fragment relative to *yccE* and show sense (upper) and antisense (lower) transcription. Bars labelled
534 “a”-“k” have at least 2-fold over background activity (empty pRW50; shown by dashed line). The
535 black bar represents the canonical promoter *PyccE*. Note that, two DNA fragments resisted cloning.
536 Hence, 19 of the 21 potential promoters were tested. Assays were done in triplicate. Error bars show
537 standard deviation from the mean.

538 **b) Transcription can initiate at multiple sites within *yccE* *in vitro*.** Gel image showing RNA
539 generated *in vitro* separated by denaturing gel electrophoresis. DNA templates, with the *yccE* gene
540 cloned upstream of the *loop* terminator in plasmid pSR, are illustrated above the gel. Transcripts
541 generated by RNA polymerase (400 nM) with empty pSR plasmid (lane 1) are highlighted by a black
542 dashed line. Transcripts initiating within *yccE* in the forward (lanes 2-5) or reverse (lanes 6-9)
543 orientation are highlighted by a blue dashed line. The control RNA-I transcripts are highlighted by a
544 grey dashed line. H-NS was added at concentrations of 0.8, 1.5, or 3.0 μ M. The experiment, done
545 three times, is representative.

546 **c) Mutation of promoters within *yccE* prevents induction in Δhns cells.** The lower illustrations
547 show *yccE* (blue arrows) cloned upstream of *lacZ* (red arrow). A solid blue arrow represents wild type
548 *yccE* whereas open arrows indicate *yccE* with mutated intragenic promoter -10 elements. For each
549 *lacZ* fusion, β -galactosidase activity was measured in lysates of M182 (grey bars) and M182*hns::kan*
550 cells (open bars). Assays were done in triplicate and error bars show standard deviation from the
551 mean.

552 **d,e) The fitness cost of *yccE* is reduced when intragenic promoters are mutated.** The figure
553 illustrates changes in culture OD₆₅₀ following inoculation of LB medium. The inoculum was M182
554 (solid line) or M182*hns::kan* (dashed line) transformed with the pSR plasmid carrying d) wild type
555 *yccE* or e) *yccE* with internal promoter -10 elements mutated. Cells were grown at 37°C. The
556 experiment was done in triplicate. Error bars show standard deviation from the mean.

557

558 **Figure 3: H-NS represses intragenic transcription and associated fitness costs at many loci.**

559 **a) Transcription initiation within the coding regions of H-NS target genes *in vitro*.** An *in vitro*
560 transcription assay using different AT-rich genes, cloned upstream of the *loop* terminator in plasmid
561 pSR, as a template. The different DNA constructs are illustrated above the gel. The cloned genes have
562 an AT-content of 65% (*yfdF*), 63% (*ykgH*), 63% (*yjgN*) and 68% (*yjgL*). For each cloned gene a solid
563 arrow represents the wild type DNA sequence whereas an open arrow is a derivative where intragenic
564 promoter -10 elements have point mutations. Note that the *fepA* gene was used as a control and has an
565 AT-content of 55%. The positions of transcripts generated by RNA polymerase (400 nM) from the
566 *fepE* control (lane 1) or the other cloned genes (lanes 2-9) are labelled. The *in vitro* transcription
567 assays were run on three separate occasions. Data are representative.

568 **b) Increased transcription in cells lacking H-NS frequently requires intragenic promoters *in vivo*.**
569 The panel illustrates a series of DNA constructs where different gene coding regions have been
570 cloned upstream of *lacZ* (red arrow). For each cloned gene a solid arrow represents the wild type
571 DNA sequence whereas an open arrow is a derivative where intragenic promoter -10 elements have
572 point mutations. The cloned genes have an AT-content of 65% (*yfdF*), 63% (*ykgH*), 63% (*yjgN*) and
573 68% (*yjgL*). Note that the *fepA* gene is used as a control and has an AT-content of 55%. For each *lacZ*
574 fusion β -galactosidase activity was measured in lysates of M182 (grey bars) and M182*Δhns* (open
575 bars) cells. Assays were done in triplicate and error bars show standard deviation from the mean.

576 **c,d) The toxicity of many AT-rich genes is a consequence of spurious intragenic transcription.**
577 The figure illustrates growth of M182 (solid line) or M182*hns::kan* (dashed line) cells transformed
578 with the pSR plasmid carrying different AT-rich genes. Panel c) shows wild type gene derivatives and
579 d) shows derivatives with internal promoter -10 elements mutated (open arrows). Experiments were
580 done using M9 minimal media at 30°C. The experiment was done in triplicate and error bars show
581 standard deviation from the mean.

582

583 **Figure 4: Most transcription is uniformly downregulated in cells lacking H-NS.**

584 **a,b) Most transcription is uniformly downregulated in cells lacking H-NS.** a) the plot illustrates
585 changes in global transcription caused by loss of H-NS. Data points represent H-NS bound (red) and
586 unbound (black) genes. Genes with unaltered transcription should fall on the diagonal blue line. Data
587 are from duplicate RNA-seq experiments. b) The basally expressed *fad* genes whose transcription is
588 reduced in the absence of H-NS. Genes bound by H-NS are in red and other genes are black. Graphs
589 show H-NS binding²⁷ and RNA abundance (RNA-seq with spiked in control; this work). Data are
590 representative.

591 **c) RNA polymerase is redistributed in cells lacking H-NS.** RNA polymerase distribution in wild
592 type (top) and *hns::kan* cells (middle). Each heat map shows the average position of DNA-bound (i.e.
593 transcribing or interacting with a promoter) RNA polymerase molecules within the cell. The bottom
594 panel shows the average position of Ter as determined by visualising a TetR-mYpet fusion bound at an
595 array of Ter proximal *tetO* sequences. Each distribution was generated from 100 cells between 3.5 and
596 4.5 μ m in length. Each square is 1/624 of total cell area.

597 **d) The σ^{70} G424D mutation hinders constitutive but not activator dependent NM501 promoter
598 activity *in vitro*.** KMnO₄ footprinting reactions analysed on a denaturing polyacrylamide gel. Bands
599 are indicative of open complex near the transcription start site (+1). RNA polymerase added at

600 concentrations of 200, 250, 300 or 350 nM and CRP was 1.0 μ M. The experiment, done three times,
601 is representative.

602 **e,f) The σ^{70} G424D mutation hinders constitutive but not activator dependent promoter activity**
603 ***in vivo***. Different promoter DNA fragments were cloned upstream of *lacZ* (red arrow). For each
604 promoter the location of key DNA sequence elements is represented by a box. In each case, the box is
605 coloured according to the relationship between the DNA sequence and the consensus sequence for
606 that element: perfect (dark blue), imperfect (pale blue) or completely absent (white). For each *lacZ*
607 fusion β -galactosidase activity was measured in lysates of JCB387*rpoD*::kan carrying pVR σ (white
608 bars) or pVR σ^{G424D} (black bars). The experiment was done in triplicate. Error bars show standard
609 deviation from the mean.
610

Lamberte_Figure 1







