

Dimer and trimer nanoantennas from a transformation optics perspective

Pacheco-Peña, Victor; Fernández-Domínguez, Antonio I; Beruete, Miguel; Luo, Yu; Navarro-Cia, Miguel

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Pacheco-Peña, V, Fernández-Domínguez, AI, Beruete, M, Luo, Y & Navarro-Cia, M 2017, Dimer and trimer nanoantennas from a transformation optics perspective. in S Zouhdi , J Rho & H Lee (eds), *The 8th International Conference on Metamaterials, Photonic Crystals and Plasmonics, Meta'17*. META, Gif-sur-Yvette, META'17, Incheon, Korea, Republic of, 25/07/18. <https://metaconferences.org/ocs/files/meta17_proceedings.pdf>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 23/07/2018

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Dimer and trimer nanoantennas from a transformation optics perspective

V. Pacheco-Peña¹, A.I. Fernández-Domínguez², M. Beruete^{1,3}, Y. Luo⁴, and M. Navarro-Cía^{5*}

¹Antennas Group-TERALAB, Universidad Pública de Navarra, Spain

²Departamento de Física Teórica de la Materia Condensada and Condensed Matter Physics Centre (IFIMAC),
Universidad Autónoma de Madrid, Spain

³Institute of Smart Cities, Universidad Pública de Navarra, Spain

⁴School of Electrical & Electronic Engineering, Nanyang Technological University, Singapore

⁵School of Physics and Astronomy, University of Birmingham, United Kingdom

*corresponding author: m.navarro-cia@bham.ac.uk

Abstract-Understanding and controlling the light-matter interaction is of fundamental importance for science and technology. Scenarios involving nanoemitters and nanoantennas are nowadays routinely measured in the lab. However, interpretation of the observations is not always straightforward. We will show here how conformal transformation can be used to provide an analytical description of the commonly used bowtie nanoantenna and its trimer counterpart.

Most recent effort in nanophotonics and nanoplasmonics has been directed toward fabrication advancements. Very complex structures have been realized experimentally and their properties are routinely measured [1]. In combination with simulations the community has gained significant insight on the physics at the nanoscale [2],[3]. However, further insight could be gained if analytical descriptions of practical scenarios are laid down. In addition, an analytical frame would speed up the design procedure.

Transformation optics attracted the attention of the research community when it was used as a design tool for invisibility cloaking [4]. More recently, this technique has shown great promise to understand plasmonic nanoantennas [5], bridging the void of analytical tools at the nanoscale. Motivated by this, we looked at conformal transformation to provide an analytical description of two-dimensional bowtie nanoantennas [6]. Here, we extend this work on bowtie nanoantennas and investigate the underlying physics in trimers. We restrict ourselves to two-dimensional nanoantennas operating in quasi-statics to keep the analytical character of the results.

The bowtie nanoantenna and its trimer counterpart are complex geometries to handle analytically. However, by applying the conformal transformation $z = \ln(z')$, where z and z' are the spatial coordinates in the transformed and original frame respectively, the nanoantennas are mapped into an infinite array of metal slabs. If the nanoantennas are excited by a single nanoemitter modelled as a line dipole, such line dipole is transformed into an array of line dipoles; for the bowtie, the unit cell with periodicity 2π in the transformed space will be comprised of two metal slabs, whereas for the trimer, it will include three metal slabs. By virtue of the conformal transformation and the quasi-static treatment, the power dissipated in the original and transformed frames are identical. The latter can be computed by simply evaluating the electric field at the dipole position:

$$P_{nr} = P_{abs}^{(x',y')} = P_{abs}^{(x,y)} = -\frac{1}{2} \omega \text{Im}\{p_x E_{1x} + p_y E_{1y}\} \quad (1)$$

where P_{nr} is the nonradiative power emission by the dipole, ω is the angular frequency, p_x and p_y are the components of the emitter dipole moment along x and y directions, and E_{1x} and E_{1y} are the components of the electric field along x and y in the region of the z plane where the line dipole is located.

Figure 1 shows the nonradiative Purcell enhancement (P_{nr} normalized to the 2D dipole radiation $P_0 = 1/16\omega^3\mu_0|p|^2$; μ_0 is the free-space permeability and $|p|$ the magnitude of the dipole moment) for an aluminum bowtie and its trimer counterpart as a function of the nanoemitter orientation. The extra arm of the trimer yields a significant increase of the nonradiative Purcell enhancement and makes the nanoantenna more robust to the dipole orientation [7].




Figure 1. Nonradiative Purcell enhancement spectra for an 8nm-arm aluminum dimer (left) and trimer (right) under different emitter orientation excitation.

Acknowledgements, work supported by the Spanish Government [TEC2014-51902-C2-2-R]; V.P.-P. was sponsored by the Spanish MECD [FPU AP-2012-3796]; A.I.F.-D. by EU 7th Framework Programme [FP7-PEOPLE-2013-CIG-63099] and the Spanish MINECO [FIS2015-64951-R]; M.B. by the Spanish Government [RYC-2011-08221]; Y.L. by the Singapore Ministry of Education [MOE2015-T2-1-145], NRF-CRP grant [NRF2015NRF-CRP002-008] and NTU-A*STAR Silicon Technologies Centre of Excellence [11235150003]; M.N.-C. by University of Birmingham [Birmingham Fellowship].

REFERENCES

1. Novotny, L. and B. Hecht, *Principles of Nano-Optics*, Cambridge University Press, UK, 2012.
2. Aouani, H., M. Rahmani, M. Navarro-Cía and S. A. Maier, "Third-Harmonic-Upconversion Enhancement from a Single Semiconductor Nanoparticle Coupled to a Plasmonic Antenna," *Nature Nanotech.*, Vol. 9, No. 4, 290-294, 2014.
3. Chikkaraddy, R., *et al.*, "Single-molecule strong coupling at room temperature in plasmonic nanocavities," *Nature*, Vol. 535, No. 7610, 127-130, 2016.
4. Schurig, D., J.J. Mock, B. J. Justice, S.A. Cummer, J.B. Pendry, A.F. Starr and D.R. Smith, Metamaterial electromagnetic cloak at microwave frequencies, *Science*, Vol. 314, No. 5801, 977-980, 2006.
5. Pendry, J.B., A.I. Fernández-Domínguez, Y. Luo and R. Zhao, "Capturing Photons with Transformation Optics," *Nat. Physics*, Vol. 9, No. 8, 518-522, 2013.
6. Pacheco-Peña, V., M. Beruete, A.I. Fernández-Domínguez, Y. Luo and M. Navarro-Cía, "Description of bow-tie nanoantennas excited by localized emitters using conformal transformation," *ACS Photon.*, Vol. 3, No. 7, 1223-1232, 2016.
7. Pacheco-Peña, V., A.I. Fernández-Domínguez, Y. Luo, M. Beruete and M. Navarro-Cía, "Aluminum Nanotriangles for Light-Matter Coupling Robust to Nanoemitter Orientation," *submitted*.