

## Changes in the ankle muscles co-activation pattern after 5 years following total ankle joint replacement

De la Fuente, Carlos; Martinez-Valdes, Eduardo; Cruz-Montecinos, Carlos; Guzman-Venegas, Rodrigo ; Arriagada, David; Peña y Lillo, Roberto ; Henriquez, Hugo ; Carpes, Felipe

DOI:

[10.1016/j.clinbiomech.2018.09.019](https://doi.org/10.1016/j.clinbiomech.2018.09.019)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

De la Fuente, C, Martinez-Valdes, E, Cruz-Montecinos, C, Guzman-Venegas, R, Arriagada, D, Peña y Lillo, R, Henriquez, H & Carpes, F 2018, 'Changes in the ankle muscles co-activation pattern after 5 years following total ankle joint replacement', *Clinical Biomechanics*, vol. 59, pp. 130-135.  
<https://doi.org/10.1016/j.clinbiomech.2018.09.019>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Changes in the ankle muscles co-activation pattern after 5 years following total ankle joint replacement**

2

3 Carlos De la Fuente<sup>a,f,k</sup> (delafuente@gmail.com), Eduardo Martinez-Valdes<sup>b,c</sup>  
4 (e.a.martinezvaldes@bham.ac.uk), Carlos Cruz-Montecinos<sup>d,e</sup> (ccmkine@gmail.com), Rodrigo Guzman-  
5 Venegas<sup>f</sup> (rguzman@uandes.cl), David Arriagada<sup>g</sup> (david.arriagada.t@gmail.com), Roberto Peña y Lillo<sup>g,h</sup>  
6 (robertopyl@gmail.com), Hugo Henríquez<sup>h,i</sup> (hhenrquez@yahoo.com.ar), Felipe P Carpes<sup>j</sup> ✉  
7 (carpes@unipampa.edu.br).

8

9 <sup>a</sup> Carrera de Kinesiología, Departamento de Cs. de la Salud, Facultad de Medicina, Pontificia Universidad  
10 Católica, #7820436 Santiago, Chile.

11 <sup>b</sup> School of Sport, Exercise and Rehabilitation Sciences, Centre of Precision Rehabilitation for Spinal Pain (CPR  
12 Spine), College of Life and Environmental Sciences, University of Birmingham, # Edgbaston B15 2TT  
13 Birmingham, United Kingdom.

14 <sup>c</sup> Centro de Investigación en Fisiología del Ejercicio (CIFE), Universidad Mayor, Santiago, Chile

15 <sup>d</sup> Laboratory of Biomechanics and Kinesiology, Hospital San José, #8380419 Santiago, Chile.

16 <sup>e</sup> Department of Physical Therapy, Faculty of Medicine, Universidad de Chile, #8380453 Santiago, Chile.

17 <sup>f</sup> Laboratorio LIBFE. Escuela de Kinesiología. Universidad de los Andes, #7620086 Santiago, Chile.

18 <sup>g</sup> Kinesiología, Universidad de Santiago de Chile, #9170124 Santiago, Chile.

19 <sup>h</sup> Servicio de Tobillo y Pie, Instituto Traumatológico, #8340220 Santiago, Chile.

20 <sup>i</sup> Facultad de Medicina, Universidad de Chile, #8380419 Santiago, Chile.

21 <sup>j</sup> Laboratory of Neuromechanics, Universidade Federal do Pampa, Campus Uruguaiana, #97508000 Uruguaiana,  
22 Brazil.

23 <sup>k</sup> Centro de Salud Deportivo, Clinica Santa Maria, #8340518, Santiago, Chile.

24

25 \* Corresponding author (✉).

26 Felipe Carpes. Email address: carpes@unipampa.edu.br (F. Carpes). Address: #97508000 Uruguaiana, Brazil.

27

28 **ABSTRACT**

29 *Background:* The Hintegra® arthroplasty provides inversion-eversion stability, permits axial rotation, ankle  
30 flexion-extension, and improvements of the gait patterns are expected up to 12 months of rehabilitation.  
31 However, sensorimotor impairments are observed in ankle flexors/extensors muscles after rehabilitation, with  
32 potential negative effects on locomotion. Here we determined the timing and amplitude of co-activation of the  
33 tibialis anterior and medial gastrocnemius muscles during gait by assessing non-operated and operated legs of  
34 patients with total ankle replacement, 5 years after surgery.

35 *Methods:* Twenty-nine patients (age: 58 [5.5] years, height: 156.4 [6.5] cm, body mass: 72.9 [6.5] kg, 10 men,  
36 and 19 women) that underwent Hintegra® ankle arthroplasty were included. Inclusion criteria included 5 years  
37 prosthesis survivorship. The onset and offset of muscle activation (timing), as well as the amplitude of  
38 activation, were determined during barefoot walking at self-selected speed by surface electromyography. The  
39 timing, percentage, and index of co-activation between the tibialis anterior and medial gastrocnemius were  
40 quantified and compared between non-operated and operated legs.

41 *Findings:* The operated leg showed higher co-activation index and temporal overlapping between tibialis  
42 anterior and medial gastrocnemius during gait ( $P < .001$ ).

43 *Interpretation:* The neuromuscular changes developed during the process of degeneration do not appear to be  
44 restored 5 years following arthroplasty. The insertion of an ankle implant may restore anatomy and alignment  
45 but neuromuscular adaptations to degeneration are not corrected by 5 years following joint replacement.

46

47

48 Abstract's word count: 229/250

49 Main text's word count: 3544/4000

50

51 *Keywords:* EMG; joint stiffness; total ankle replacement; walking; gastrocnemius; tibialis anterior.

## 52 **1. Introduction**

53 Lower limb osteoarthritis is associated with pain and impaired function with negative effects on locomotion.  
54 More specifically, ankle osteoarthritis impairs sagittal plane motion (dorsiflexion and plantarflexion) and torque  
55 (Al-Mohrougi et al., 2018). In many cases the joint degeneration will reach levels that require the total joint  
56 replacement. Ankle arthroplasty is a surgical procedure in which the tibiotalar joint is replaced (Giannini et al.,  
57 2000; Caravaggi et al., 2015). Eighty percent of candidates are patients who developed ankle osteoarthritis  
58 secondary to trauma (post-traumatic etiology) (Horisberger et al., 2009; Bloch et al., 2015), with far fewer  
59 outcomes reported compared to hip and knee arthroplasty (Al-Mohrougi et al., 2018). In the past 40 years more  
60 than 30 models of ankle prosthesis have been introduced, with few outcomes against hip and knee arthroplasty  
61 (Giannini et al., 2000; Henricson et al., 2011). In this regard, Hintegra® arthroplasty is a third generation  
62 prosthesis that provides inversion-eversion stability, permits axial rotation, ankle flexion-extension being one of  
63 the few ankle prosthesis showing good results concerning activities of daily living (Michael et al., 2008;  
64 Hintermann, 2005). Previous studies showed that the Hintegra prosthesis is built with stable components, with  
65 low rate of complications (Michael et al., 2008; Hintermann, 2005; Barg et al., 2011), allowing an adequate  
66 range of motion (Valderrabano et al., 2003; Michael et al., 2008; Hintermann, 2005). However, the lack of  
67 restoration of gait symmetry and joint function remains a problem after ankle arthroplasty (Caravaggi et al.,  
68 2015).

69 During the first year after Hintegra® arthroplasty, there are adaptations in the first three months of  
70 rehabilitation that negatively affects the gait biomechanics. These include lower maximal plantar-flexion  
71 moment, total adduction moment, medial ground reaction force, and higher anterior ground reaction force  
72 during gait by increasing mechanical loading on knee joint (Valderrabano et al., 2007). This adaptive period was  
73 previously reported as negative phase of rehabilitation (De la Fuente et al., 2014). These parameters are  
74 expected to improve towards the end of the first year after surgery (Valderrabano et al., 2007), with significant  
75 improvement of gait pattern (Valderrabano et al., 2007; Aidi et al., 2013) and quality of life (Esparragoza et al.,  
76 2011). It has been suggested that 5 years following total ankle joint replacement there is an increase in the  
77 American Orthopaedic Foot and Ankle Score (Zaidi et al., 2013). However, there is little information  
78 concerning the effect of ankle replacement on neuromuscular activity of ankle muscles (tibialis anterior and  
79 medial gastrocnemius) (Doets et al., 2007). Abnormal muscle co-activation could result in increased and  
80 abnormal kinetics on the prosthesis predisposing to loosening. Presumably the prosthesis is designed to replicate

81 normal anatomy of the ankle joint. If however the muscular co-activation does not change the forces will remain  
82 high.

83 An adequate control of tibialis anterior and medial gastrocnemius muscles depends both on the timing of  
84 agonist and antagonist activation (Hubley-Kozey et al., 2010; Rosa et al., 2010), and the intensity of their  
85 contractions, i.e. weight coefficients (Cappelini et al., 2006). These muscles sometimes present co-activation, an  
86 involuntary and concurrent activation of the antagonist, in opposition to the contraction of the agonist  
87 (Duchateau et al., 2014). Changes in co-activation occur during different physical activities e.g. running  
88 compared to walking (Cappelini et al., 2006), footwear selection (Alkjær et al., 2012) and in pathological  
89 processes e.g. cerebellar ataxia (Mari et al., 2014), foot and knee motor functions (Di Nardo et al., 2015), and  
90 ankle osteoarthritis (Von Tscharnner & Valderrabano, 2010). Furthermore, co-activation may affect gait pattern  
91 and the integrity of prostheses due to abnormal joint force development (De la Fuente et al., 2014).

92 After the Hintegra<sup>®</sup> arthroplasty placement, the gait pattern is expected to be restored, and co-activation  
93 between tibialis anterior and medial gastrocnemius should not occur at medium term after the first three months  
94 of rehabilitation (Michael et al., 2008; Hintermann, 2005). However, there is a lack of evidence concerning  
95 patterns of co-activation after ankle replacement when rehabilitation is completed. Therefore, here we determine  
96 the relative measures of amplitude and timing of co-activation of the tibialis anterior and medial gastrocnemius  
97 muscles during gait, by comparing the electromyographic activity of the non-operated and operated leg of  
98 patients with total ankle replacement, 5 years after surgery. Our null-hypotheses were: i) total ankle replacement  
99 after 5 years does not recover the timing of co-activation to the level of the non-operated leg during gait; ii) total  
100 ankle replacement after 5 years does not recover the temporal percentage of co-activation to the level of the non-  
101 operated leg during gait, and ii) total ankle replacement after 5 years does not recover the index of co-activation  
102 to the level of the non-operated leg during gait.

103

## 104 **2. Methods**

### 105 2.1. Study design

106 This is a cross-sectional, observational, analytical study design. The sample included twenty-nine patients (10  
107 men and 19 women) that underwent unilateral total ankle replacement 5 years before the measurement session  
108 in which the operated and non-operated leg were compared during gait. The inclusion criteria were: i) unilateral

109 total ankle replacement with Hintegra<sup>®</sup> arthroplasty (Newdeal SA, Vienne, France); ii) at least 5 years following  
110 replacement; iii) rehabilitation treatment of one year; iv) posttraumatic arthritis; v) age between 40 and 70 years  
111 old; vi) passive dorsal range of motion greater than 5° at the sagittal plane (De la Fuente et al., 2014); vi)  
112 radiological stability (Horisberger et al., 2009; Bai et al., 2010; Guyer and Richardson, 2008); and v) have been  
113 rehabilitated at the *Instituto Traumatológico* (Santiago, Chile) by the same surgical and rehabilitation team. The  
114 exclusion criteria were: i) need for assisted locomotion; ii) major limitations in performance of daily life  
115 activities; iii) major periarticular tissue impairment; iv) ipsilateral or contralateral hip/knee osteoarthritis; v)  
116 inflammatory diseases; vi) neurological pathology; vii) active infection; and viii) cognitive impairment (von  
117 Tschärner and Valderrabano, 2010). This study was approved by the institutional review board of the *Instituto*  
118 *Traumatológico* (Santiago, Chile) according to the principles of the Declaration of Helsinki. All participants  
119 signed a consent term agreeing to participate in this study.

120

## 121 2.2. Sample size

122 The sample size was estimated “a priori” with a pilot experiment that included 6 patients [mean (standard  
123 deviation) 56.5 (2.1) years-old, 30.1 (1.8) kg/m<sup>2</sup>, 3 men and 3 women] that fulfilled the inclusion and exclusion  
124 criteria. A sample size of 23 patients was estimated considering a difference between two dependent means  
125 (matched pairs), using two-tailed *t*-test with alpha error of 5% and statistical power of 80% for an estimated  
126 effect size of 0.62. Six additional patients were included to the sample to anticipate possible attrition (20% of  
127 estimation). The total patients assessed were 35 patients; 6 patients from the “a priori” determination of sample  
128 size and additional 29 patients. The statistical calculus was performed by G\*Power software version 3.1.9.2.  
129 (Kiel University, Germany).

130

## 131 2.3. Surgery and physical therapy procedures

132 All participants had a Hintegra<sup>®</sup> prosthesis (Hintermann, 2005) (Newdeal SA, Vienne, France). In general terms,  
133 surgery involved an anterior longitudinal incision of 10 to 12 cm performed to dissect the retinaculum.  
134 Moreover, the soft tissue and periosteum from the bone were dissected. Resection of the talus and tibia were  
135 performed to insert arthroplasty components using Hintermann<sup>®</sup> distractor and oscillating saw. Osteophytes on  
136 the talar neck and anterior aspect of medial malleolus were also removed. After that, the tibial and talar

137 components were inserted. The last inserted component was the polyethylene component. The tissues were  
138 sutured, and the procedure was finalized by fitting a short leg cast. The patients were monitored for the next  
139 three weeks after surgery, once per week, and during the first 4 weeks after surgery patients were immobilized  
140 with a short leg cast and instructed to unload the operated leg and rest.

141           The physiotherapy intervention was performed from week 4 until week 52 (Ingrosso et al., 2009).  
142 From the week 4 to 12, the patients attended the rehabilitation service, where they used a walking boot. In this  
143 period, partial weight bearing was permitted using the assistance of canes. Furthermore, stretching to improve  
144 the dorsal flexion range of motion, strengthening of ankle, knee and hip muscles, and pain relief with physical  
145 agents were activities performed. The re-education of gait without assistance, bipedal heel rise, balance  
146 exercises, and pain relief management were performed until the end of week 52 (Martin et al., 2007).  
147 Afterwards, patients returned to the to the foot and ankle hospital unit every six months to be assessed in the  
148 follow-up period.

149

#### 150 2.4. Data acquisition and processing

151 The data acquisition consisted of two stages: clinical assessment and surface electromyography recordings. The  
152 patients attended an interview, and the clinical assessment was performed at the foot and ankle service of the  
153 *Instituto Traumatológico* (Santiago, Chile). The age, body mass, height, AOFAS score (Kitaoka et al., 1994),  
154 passive dorsiflexion range of movement, difference in calf whilst standing (Saxena et al., 2011; Valdebarrano et  
155 al., 2006), and intensity of pain at rest and during walking assessed with a numerical verbal scale (0 no pain, 10  
156 maximum possible pain) (Hintermann, 2005) and were part of the clinical assessment.

157           Surface electromyography recordings were performed one week after the clinical assessment at the  
158 *Centro de Investigaciones Medicas del Instituto Traumatológico* (Santiago, Chile). After a 5-min warm-up on a  
159 cycloergometer without external load and cadence of 60 rpm, patients walked barefoot at self-selected speed  
160 (von Tscharnier and Valdebarrano, 2010; De la Fuente et al., 2014) along a 5-meter flat surface on a straight-  
161 line. Patients performed five trials for familiarization within the walking space. Although we did not measure  
162 kinematics, a common heel strike pattern was observed among the patients. The electromyography signals were  
163 acquired during walking using a Myomonitor IV electromyography amplifier (Delsys, inc., Boston, USA). Two  
164 DE-2.3 single differential surface electromyography sensors (Delsys, inc., Boston, USA) with an inter-electrode

165 distance of 10 mm were used. The data collection employed a 16-bit analog-digital converter card (National  
166 Instrument Corp., Austin, TX, USA) operated by a Matlab software (Mathworks Inc., Massachusetts, USA) at a  
167 sampling rate of 1000 Hz, band-pass filtered (20–450 Hz), and hardware amplified with a gain of 1000 V/V.  
168 The muscles assessed were tibialis anterior and medial gastrocnemius based on their lower mean  
169 electromyography frequency and intensity previously found by Valderrabano et al. (2006) in unilateral ankle  
170 osteoarthritis, and the hypothesis of Doets et al. (2007), who suggested that after ankle joint replacement, higher  
171 co-activation between these muscles could exist during gait. The surface electromyography sensors were placed  
172 according to the European recommendations for surface electromyography (Hermens et al., 2000).

173

#### 174 2.5. Signal treatment

175 The electromyography signals were filtered by a zero-lag 4<sup>th</sup> order finite impulse response Butterworth with a  
176 band pass of 20 to 450 Hz. Onset and offset times of muscle activation (Caravaggi et al., 2015) were identified  
177 using a continuous wavelet with the algorithm proposed by Merlo et al. (2003). The algorithm was performed  
178 using the Hermite-Rodriguez mother wavelet, 10% of noise power, 150 ms for the time of two detected  
179 activation intervals, and 5 ms for the spike rejection from non-rectified signals. The noise of electromyography  
180 signals was extracted when patients stood quietly before walking. Five consecutive electromyography bursts of  
181 the tibialis anterior and gastrocnemius medialis (corresponding to 5 strides) were used for the analysis.

182 All signals were full wave rectified (Figure 1). Due to the variability and non-accordance of  
183 normalization methods for analysis of electromyography signals (Rosa et al., 2010), a standardized treatment of  
184 signals by z-score method was performed. Each sample was subtracted from its expected value ( $E[X]$ ) and  
185 divided by the standard deviation. This expresses the number of standard deviations by which the sample is  
186 above the  $E[X]$ . As the whole electromyography signals showed a normal distribution, the z-score was obtained  
187 using the arithmetic mean of the data.

188

#### 189 2.6. Outcomes

190 *The timing of co-activation:* determined by the time overlap between the onsets and offsets activation (Rosa et  
191 al., 2010).



192 *Percentage of co-activation*: from the time activation onsets and offsets, the percentage of co-activation was  
193 quantified as the overlapped percentage of muscle activation (Rosa et al., 2010).

194 *Co-activation index*: determined by the overlapped amplitudes of muscle activation between the tibialis anterior  
195 and medial gastrocnemius. The co-activation index was implemented in discrete form by the trapezoidal method  
196 (Eq. 1) from the original continuous form previously defined in the literature (Rosa et al., 2010).

197

$$198 \quad Coactivation\ index = 2 \times \frac{\sum_{CoAc_{onset}}^{CoAc_{offset}} \left[ \frac{(X_{i+1} - X_i)}{2} \times \Delta t \right]}{\sum_{Agonist}^{Antagonist} \left[ \sum_{onset}^{offset} \frac{(X_{i+1} - X_i)}{2} \times \Delta t \right]_j} \times 100\% \quad (Eq. 1)$$

199

200 In the equation 1,  $CoAc_{onset/offset}$  is the co-activation time of onset or offset,  $x_i$  is the electromyography  
201 sample,  $\Delta t$  is the interval time of data acquisition given the sampling frequency, and  $j$  represents the agonist or  
202 antagonist condition. To assess the intensity of concurrent contractions, the trapezoidal areas were obtained  
203 using the “cumtrapz function” from the full-rectified signals using the Matlab software (Mathworks Inc.,  
204 Massachusetts, USA).

205

## 206 2.7. Statistical analysis

207 Data were reported as the median and interquartile range [IQ range] because the Shapiro-Wilk test revealed a  
208 non-parametric data distribution. Homoscedasticity was confirmed using the Levene’s test. To compare the  
209 timing, percentage, and index of co-activation of medial gastrocnemius and tibialis anterior between the  
210 operated and non-operated legs, a Wilcoxon Signed-Ranks test of two-tails was used with alpha error equal to  
211 5%. To assess the possible existence of co-variable and interaction with outcomes, a multiple regression respect  
212 to analysis for age, body mass, height, AOFAS score, the passive dorsal range of movement, calf circumference  
213 difference between leg during a standing posture, the intensity of pain at rest and during gait were considered at  
214  $p < 0.05$ . Data were analyzed using the Matlab software statistical toolbox (Mathworks Inc., Massachusetts,  
215 USA).

216

217 **3. Results**

218 The timing of co-activation between the tibialis anterior and medial gastrocnemius in the non-operated leg  
219 [median: 390 ms, IQ range: 80 ms] was lower than in the operated leg [median: 566 ms, IQ range: 104 ms,  
220  $p<0.001$ ]. The percentage of co-activation between the tibialis anterioris and gastrocnemius medialis in the non-  
221 operated leg [median: 0.00 %, IQ range: 1.84 %] was lower than in the operated leg [median: 100%, IQ range:  
222 0.00 % $p<0.001$ ]. The index of co-activation between the tibialis anterior and medial gastrocnemius in the non-  
223 operated leg [median: 0.00 %IQ range: 1.94 %] was lower than in the operated leg [median: 30.96%, IQ range:  
224 13.52%,  $p<0.001$ ]. The changes in co-activation are depicted in the Figure 1.

225 The demographic characteristics of the sample regarding age, body mass, height, AOFAS score,  
226 passive dorsal range of movement, calf circumference difference between legs during standing posture, intensity  
227 of pain at rest, and during gait did not co-vary or interacted with any of the electromyography outcomes  
228 ( $p>0.05$ ).

229

230 \*\*\* Table 1 around here \*\*\*

231 \*\*\* Figure 1 around here \*\*\*

232

233 **4. Discussion**

234 The most important finding of this work was that patients with the Hintegra® arthroplasty present abnormal  
235 plantar and dorsi-flexion co-activation patterns during gait after 5 years following total ankle joint replacement.  
236 Although the patients did not receive rehabilitation after 5 years following total ankle joint replacement due to  
237 the improvement observed in the AOFAS score, the present findings suggest the need for an appropriate  
238 sensorimotor rehabilitation program for patients in which problems at medium term of survival of the prosthesis  
239 still persist.

240 Although it can't be ensured (due to the lack of kinetics and kinematics assessment), the altered co-  
241 activation found in our study may suggest that both absorption and propulsion phases of walking could be  
242 committed 5 years after replacement surgery. Dorsal and plantar-flexor muscles need to be activated during  
243 different phases of gait in order to maintain the normal mechanics of the ankle joint during swing and load

244 response. This is important for the control of the anterior advance of the tibia or the generation of propulsion.  
245 Therefore, these mechanical actions need temporal muscle coordination (timing of activation). Based on the  
246 findings from Di Nardo et al. (2015) and Cappellini et al. (2006), the tibialis anterior and medial gastrocnemius  
247 muscles should present independent temporal activations during the different phases of normal gait (low speed).  
248 It contrasts with the results of patients treated with ankle prosthesis, since a temporal overlap between these  
249 muscles was found.

250         Based on the findings from Siegler et al. (2013), an altered kinetic condition can be present in patients  
251 with Hintegra<sup>®</sup> arthroplasty. Siegler et al. (2013) tri-dimensionally modeled the talus of 26 healthy adults and  
252 found that “the trochlear surface can be modeled as a skewed truncated conic saddle shape with its apex oriented  
253 laterally rather than medially”, which contrasts with the mechanical model of Inman, in which the Hintegra<sup>®</sup>  
254 arthroplasty is based (Hinterman 2005). Thus, the mechanical design of the Hintegra<sup>®</sup> arthroplasty could alter  
255 distribution or intensity of ankle joint forces during the stance phase, or both, such as the abnormal joint shear  
256 forces and delayed the first peak of vertical ground reaction force found after the negative rehabilitation phase in  
257 patients treated with an Hintegra<sup>®</sup> arthroplasty (De la Fuente et al., 2014). Our data suggests that these  
258 mechanical changes could result of sensorimotor adaptations, since an increased co-activation between dorsal  
259 and plantar flexors would reduce the velocity of the foot and tibia during stance phase, which reduces the  
260 vertical component of the ground reaction force at load response phase. However, an increased co-activation  
261 might also result of intrinsic instability of the prosthesis in the frontal plane, which could have an unstable effect  
262 over the foot based on mechanical findings from Siegler et al. (2013), further leading to the shifting of the  
263 medial gastrocnemius rather than a shift of the tibialis anterior activation, as shown here.

264         As the shifting in the medial gastrocnemius could occur, the propulsion phase might also be affected  
265 (Giannini et al., 2000; Henricson et al., 2011). This possible change is in accordance with de la Fuente et al.  
266 (2014), who proposed a gait pattern with external hip rotation that increases the activity of hip extensors (rather  
267 than plantar flexors) to generate the propulsion of the leg with Hintegra<sup>®</sup> arthroplasty, after the negative  
268 rehabilitation phase. Therefore, it is possible that patients with Hintegra<sup>®</sup> arthroplasty experience an over-  
269 stabilization of the ankle joint after at medium term, affecting the whole pattern of gait by an unstable load  
270 response and altered propulsion phase. This hypothesis still needs to be investigated.

271         On the other hand, Stubbs et al. (1998) suggested that the load over viscoelastic tissues creates a  
272 sensorimotor response called “muscle-ligament reflex” causing an altered activity of agonist and antagonist

273 muscles over time. Therefore, the surgery itself could cause the neuromuscular changes observed, since the  
274 tibialis anterior and ankle ligaments must be pulled to introduce the arthroplasty components into the ankle joint  
275 space. This is important as one wonders if neuromuscular changes are reversible with time. Our findings would  
276 suggest that at 5 years after ankle replacement they are not reversible. In making this statement perhaps  
277 prosthesis designers would be better to use an arthroplasty designed to tolerate the irreversible forces on the  
278 prosthesis. Finally, chronic pain could be another source of the neuromuscular changes observed in our study.  
279 Valderrabano et al., (2006) and von Tscharner et al. (2010) described that the time of exposure to pain in  
280 osteoarthritis is the worst stimulus to the nervous system changing the neuromuscular patterns of activations.  
281 This could induce changes in the ankle stiffness leading to pathological neuroplasticity changes, possibly  
282 present since before the arthroplasty surgery. However, in our study, the intensity of pain was not identified as a  
283 co-variable and did not show any interaction with the outcomes.

284         Due to the higher co-activation found in patients with Hintegra® arthroplasty at 5 years after ankle  
285 replacement, new rehabilitation approaches are needed to improve the locomotion in patient who shows the  
286 sensorimotor impairments described here. To our knowledge, this is the first pathological report at medium term  
287 in patients with Hintegra® arthroplasty. Future studies should identify *in vivo* whether the ankle axis of this  
288 arthroplasty is really positioned as Inman previously reported, since alterations in the prosthesis axis of rotation  
289 could lead to the sensorimotor impairments found. Also, the presence of other concomitant musculoskeletal  
290 problems (e.g., knee, hip or trunk disorders, metatarsal-cunieform, cuneiform-navicular and talo-navicular  
291 osteoarthritis) requires further examination. Finally, it is important to investigate whether previous pathological  
292 sensorimotor conditions acquired before the arthroplasty are irreversible after the surgery (i.e., chronic pain).  
293 Regarding the limitations of our study, the sample is only representative of patients with lower AOFAS score, in  
294 contrast to the results with Hintegra® (Hinterman, 2005) and other arthroplasties (Aidi et al., 2013; Caravaggi et  
295 al., 2015). Also, the measurement of pain considered a verbal pain scale (Hinterman, 2005), which may not  
296 reflect the sensorimotor impartments showed by previous studies (Ervilha et al., 2004; Lindstrøm et al., 2011).  
297 Finally, we propose that a detailed mechanical analysis of gait, in addition to the analysis of co-activation  
298 alteration, is needed to understand the possible neuro-mechanical pathological new hypotheses discussed in our  
299 study.

300

301 **5. Conclusions**

302 Altered co-activation during gait is present after at 5 years after total ankle replacement with the Hintegra®  
303 prosthesis. It may result of an attempt to compensate ankle instabilities following arthroplasty, which effects on  
304 gait dynamics require attention during rehabilitation programs.

305

306 **Conflict of interests**

307 The authors declare no conflicts of interest.

308

309 **Funding**

310 This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-  
311 profit sectors. FPC is supported by a CNPq research fellowship.

312

313 **Acknowledgements**

314 We acknowledge the assistance in data collection provided by the undergraduate students Guillermo Gonzalez,  
315 Javiera Asecio and Roberto Peña from Universidad de Chile.

316 **6. References**

- 317 Aidi R., Cro S., Gurusamy K., Siva N., Macgregor A., Henricson A., Goldberg A. 2013. The outcome of total  
318 ankle replacement: a systematic review and meta-analysis. *Bone Joint J.* 95-B(11),1500-1507.
- 319 Alkjær T., Raffalt P., Petersen N., Simonsen E. 2012. Movement behavior of high-heeled walking: how does the  
320 nervous system control the ankle joint during an unstable walking condition?. *PLoS One.*7(5),e37390.
- 321 Al-Mahrouqi M.M., MacDonald D.A., Vicenzino B., Smith M.D. 2018. Physical impairments in adults  
322 with ankle osteoarthritis: a systematic review and meta-analysis. *J Orthop Sports Phys Ther.* 7:1-43.
- 323 Bai L., Lee K., Song E., Yoon T., Seon J. 2011. Total ankle arthroplasty outcome comparison for post-traumatic  
324 and primary osteoarthritis. *Foot Ankle Int.* 31(12),1048-1058.
- 325 Barg A., Knupp M., Anderson A., Hintermann B. 2011. Total ankle replacement in obese patients: component  
326 stability, weight change, and functional outcome in 118 consecutive patients. *Foot Ankle Int.* 32(10),925-932.
- 327 Bloch B., Srinivasan S., Mangwani J. 2015. Current Concepts in the Management of Ankle Osteoarthritis: A  
328 Systematic Review. *J Foot Ankle Surg.* 54(5),932-939.
- 329 Cappellini G., Ivanenko Y., Poppele R., Lacquaniti F. 2006. Motor patterns in human walking and running *J of*  
330 *Neurophysiol.* 95(6), 3426-3437.
- 331 Caravaggi P., Lullini G., Leardini A., Berti L., Vannini F., Giannini S. 2015. Functional and clinical evaluation  
332 at 5-year follow-up of a three-component prosthesis and osteochondral allograft transplantation for total ankle  
333 replacement. *Clin Biomech (Bristol, Avon).* 30(1),59-65.
- 334 De la Fuente C., Carcuro G., Ramírez-Campillo R., Campos C., Soza F. 2015. Prosthetic wear in subjects with  
335 Hintegra total ankle joint replacement secondary to trauma in functional phase. *Fisioterapia.* 37,165-174
- 336 Di Nardo F., Mengarelli A., Maranesi E., Burattini L., Fioretti S. 2015. Assessment of the ankle muscle co-  
337 contraction during normal gait: a surface electromyography study. *J Electromyogr Kinesiol.* 25(2),347-354.
- 338 Doets H.C., van Middelkoop M., Houdijk H., Nelissen R.G., Veeger H.E. 2007. Gait analysis after successful  
339 mobile bearing total ankle replacement. *Foot Ankle Int.* 28(3),313-322.
- 340 Dyer J.O., Maupas E., de Andrade Melo S., Bourbonnais D., Nadeau S., Forget R. 2014. Changes in activation  
341 timing of knee and ankle extensors during gait are related to changes in heteronymous spinal pathways after  
342 stroke. *J Neuroeng Rehabil.*11,148.
- 343 Ervilha U.F., Arendt-Nielsen L., Duarte M., Graven-Nielsen T. 2004. The effect of muscle pain on elbow  
344 flexion and coactivation tasks. *Exp Brain Res.*156(2),174-182.

345 Esparragoza L., Vidal C., Vaquero J. 2011. Comparative study of the quality of life between arthrodesis and  
346 total arthroplasty substitution of the ankle. *J Foot Ankle Surg.* 50(4),383–387.

347 Giannini S., Leardini A., O’connor J. 2000. Total ankle replacement: Review of the designs and of the current  
348 status. *Foot AnkleSurg.* 6,77-88.

349 Guyer A., Richardson G. 2008. Current concepts review: total ankle arthroplasty. *Foot Ankle Int.* 29(2),256-  
350 264.

351 Henricson A., Nilsson J., Carlsson A. 2011. 10 year survival of total ankle arthroplasties: a report on 780 cases  
352 from the Swedish Ankle Register. *Acta Orthop.* 82(6),655-659.

353 Hermens H.J., Freriks B., Disselhorst-Klug C., Rau G. 2000. Development of recommendations for SEMG  
354 sensors and sensor placement procedures. *J Electromyogr Kinesiol.* 10(5), 361-374.

355 Hintermann B. 2005. Total ankle arthroplasty: historical overview, current concepts and future perspectives,  
356 illustrated ed. Austria: Springer.

357 Horisberger M., Hintermann B., Valderrabano V. 2009. Alterations of plantar pressure distribution in  
358 posttraumatic end-stage ankle osteoarthritis. *Clin Biomech (Bristol, Avon).* 24,303-307.

359 Hubley-Kozey C.L., Hatfield G.L., Davidson K.C. 2010. Temporal coactivation of abdominal muscles during  
360 dynamic stability exercises. *J Strength Cond Res.*24(5),1246-1255.

361 Ingrosso S., Benedetti M., Leardini A., Casanelli S., Giannini S. 2009. Gait analysis in patients operated with a  
362 novel total ankle prosthesis. *Gait Posture.* 30,132-137.

363 Kitaoka H., Alexander I., Adelar R. 1994. Clinical rating systems for the ankle, hindfoot, midfoot, hallux, and  
364 lesser toes. *Foot Ankle Int.* 15(7),349–353.

365 Lindstrøm R., Schomacher J., Farina D., Rechter L., Falla D. 2011. Association between neck muscle  
366 coactivation, pain, and strength in women with neck pain. *Man Ther.*16(1),80-86.

367 Mari S., Serrao M., Casali C., Conte C., Martino G., Ranavolo A., Coppola G., Draicchio F., Padua L., Sandrini  
368 G, Pierelli F. 2014. Lower limb antagonist muscle co-activation and its relationship with gait parameters in  
369 cerebellar ataxia. *Cerebellum.*13(2),226-236.

370 Martin R.L., Stewart G.W., Conti S.F. 2007. Posttraumatic ankle arthritis: an update on conservative and  
371 surgical management. *J Orthop Sports Phys Ther.* 37(5),253–259.

372 Merlo A., Farina D., Merletti R. 2003. A fast and reliable technique for muscle activity detection from surface  
373 EMG signals. *IEEE Trans Biomed Eng.* 50(3),316-323.

374 Michael J., Golshani A., Gargac S., Goswami T. 2008. Biomechanics of the joint clinical outcomes of total  
375 ankle replacement. *J Mech Behav Biomed Mater.* 1(4),276-294.

376 Rosa M.C., Marques A., Demain S., Metcalf C.D., Rodrigues J. 2014. Methodologies to assess muscle co-  
377 contraction during gait in people with neurological impairment - a systematic literature review. *J Electromyogr*  
378 *Kinesiol.* 24(2),179-191.

379 Saxena A., Ewen B., Mafulli N. 2011. Rehabilitation of the operated achilles tendon: parameters for predicting  
380 return to activity. *J Foot Ankle Surg.* 50(1),37-40.

381 Siegler S., Toy J., Seale D., Pedowitz D. 2014. The Clinical Biomechanics Award 2013 -- presented by the  
382 International Society of Biomechanics: new observations on the morphology of the talar dome and its  
383 relationship to ankle kinematics. *Clin Biomech (Bristol, Avon).* 29(1),1-6.

384 Stubbs M., Harris M., Solomonow M., Zhou B., Lu Y., Baratta R.V. 1998. Ligamento-muscular protective  
385 reflex in the lumbar spine of the feline. *J Electromyogr Kinesiol.* 8(4), 197-204.

386 Valderrabano V., Hintermann B., Nigg B.M., Stefanyshyn D., Stergiou P. 2003. Kinematic changes after fusion  
387 and total replacement of the ankle: part 1:Range of motion. *Foot Ankle Int.* 24(12),881-887.

388 Valderrabano V., Nigg B., von Tscherner V., Stefanyshyn D., Goepfert B., Hintermann B. 2007. Gait analysis in  
389 Ankle Osteoarthritis and Total Ankle Replacement. *Clin Biomech (Bristol, Avon).* 22,894-904.

390 Valderrabano V., von Tscherner V., Nigg B., Hintermann B., Goepfert B., Fung T., Frank C.B., Herzog W.  
391 2006. Lower leg muscle atrophy in ankle osteoarthritis. *J Orthop Res.* 24(12),2159-2169.

392 von Tscherner V., Valderrabano V. 2010. Classification of multi muscle activation patterns of osteoarthritis  
393 patients during level walking. *J Electromyogr Kinesiol.* 20(4),676-683.

394 Zaidi R., Cro S., Gurusamy K., Siva N., Macgregor A., Henricson A., Goldber A. 2013. The outcome of total  
395 ankle replacement: a systematic review and meta-analysis. *Bone Joint J.* 95-B(11):1500-1507.



396

397 **Figure caption**

398 **Figure 1. Temporal co-activation between medial gastrocnemius and tibialis anterior during gait in the**  
399 **non-operated (top) and operated (bottom) legs from patient 1.** The EMG signals of the medial  
400 gastrocnemius are shown in light gray and positive values. The tibialis anterior signals are shown in dark gray  
401 and negative values. The line shows the identified time of muscle activity as proposed by Merlo et al (2003).  
402 The upper box shows the color intensity scale of the temporal overlapping relative to the medial gastrocnemius.  
403 The bottom boxes show the intensity of each overlap found. The signals were normalized in function of z-score.