

## SLFN14 gene mutations associated with bleeding

Stapley, Rachel; Pisareva, Vera; Pisarev, Andrey; Morgan, Neil

DOI:

[10.1080/09537104.2019.1648781](https://doi.org/10.1080/09537104.2019.1648781)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Stapley, R, Pisareva, V, Pisarev, A & Morgan, N 2019, 'SLFN14 gene mutations associated with bleeding', *Platelets*, pp. 1-4. <https://doi.org/10.1080/09537104.2019.1648781>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

Rachel J. Stapley, Vera P. Pisareva, Andrey V. Pisarev & Neil V. Morgan (2019) SLFN14 gene mutations associated with bleeding, *Platelets*, DOI: 10.1080/09537104.2019.1648781

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



## *SLFN14* gene mutations associated with bleeding

Rachel J. Stapley, Vera P. Pisareva, Andrey V. Pisarev & Neil V. Morgan

To cite this article: Rachel J. Stapley, Vera P. Pisareva, Andrey V. Pisarev & Neil V. Morgan (2019): *SLFN14* gene mutations associated with bleeding, Platelets, DOI: [10.1080/09537104.2019.1648781](https://doi.org/10.1080/09537104.2019.1648781)

To link to this article: <https://doi.org/10.1080/09537104.2019.1648781>



© 2019 The Author(s). Published with license by Taylor & Francis Group, LLC.



Published online: 04 Aug 2019.



Submit your article to this journal [↗](#)




Article views: 68



View Crossmark data [↗](#)



## SLFN14 gene mutations associated with bleeding

Rachel J. Stapley<sup>1</sup>, Vera P. Pisareva<sup>2</sup>, Andrey V. Pisarev<sup>2</sup>, & Neil V. Morgan <sup>1</sup>

<sup>1</sup>Institute of Cardiovascular Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK and <sup>2</sup>Department of Cell Biology, SUNY Downstate Medical Center, Brooklyn, NY, USA

### Keywords

Bleeding, genes, inherited thrombocytopenia, mutations, platelets, SLFN14

### History

Received 7 May 2019  
Revised 18 July 2019  
Accepted 18 July 2019  
Published online 02 August 2019

### The Schlafen (SLFN) Family

The *Schlafen* (*SLFN*) family of genes, which is limited to mammals, was initially discovered in mice by Schwarz et al., investigating the T cell lineage, regulating differentiation and in some instances ablating growth [1]. In particular, overexpression of *SLFN1* resulted in a cell cycle arrest at the G0/G1 stage. Therefore, the family had been named Schlafen, which is translated from German as “to sleep” [1]. Subsequent studies have classified the *SLFN* family into three distinct subgroups based on gene/protein size and domain homology [2,3]. Ten mouse and six human *SLFN* genes have been identified to date (Figure 1) [4]. All 10 mouse *SLFN* proteins possess a core region containing a unique “slfn box” motif with an unknown function. Subgroups II and III contain an extra domain at the C-terminus of the slfn box, conserved by the flanking five amino acid signature (Ser-Trp-Ala ... -Asp-Leu) ‘SWA ... DL’ which appears to be *SLFN* specific. This was discovered in early characterization of the *SLFN* family with *SLFN3* and *SLFN4* (members 3 and 4) possessing a 200 amino acid sequence not found in those in subgroup I [1]. Adjacent to the slfn box is the ‘ATPases associated with diverse cellular activities’ (AAA) domain. Based on protein homology studies, the AAA motif is thought to function similarly to classical AAA domains with the role of ATP/GTP binding in the course of DNA and RNA metabolism and therefore playing a fundamental role in the production and function of *SLFN* proteins [5,6]. Another protein was discovered with significant similarity to those in subgroup II, extending a further 400 amino acids towards the C-terminus terminal tail. The sequence was aligned by BLAST and the first 570 amino acids were homologous to *SLFNs* 3 and 4 while the remainder was unique to named *SLFN8* leading to the

classification of the final *SLFN* subgroup, III [2]. Additional homologous genes were identified as *SLFN5*, *SLFN9*, *SLFN10* and *SLFN14*, whereby the last two coding exons correspond to this unique subgroup. NCBI conserved domain database (CDD) searches revealed significant homology of subgroup III specific extension to motifs typical in superfamily I of RNA/DNA helicases which are known to mediate DNA and RNA metabolism [2]. The same characterization method applies to the human-specific *SLFN* genes with the absence of any subgroup I members and structural difference in *SLFN12L* (*SLFN12-LIKE*). Notably, *SLFN5* and *SLFN14* are the only *SLFN* family genes present in both human and mouse. The predicted transcript of *SLFN12L* is of a similar length to those in subgroup III; however, it also encompasses sequences which are unaccounted for and based on this structural analysis, has not yet been assigned a subgroup. According to length and homology, *SLFN14* belongs to subgroup III (Figure 1) [7]. The recently discovered crystal structure of rat *SLFN13* (related to human *SLFN13* and mouse *SLFN8*) N-domain gives a functional insight into its ability to cleave RNA acting as an endoribonuclease in inhibiting protein synthesis [8]. Importantly, *SLFN13* is the closest paralogue for *SLFN14* (44% identity and 61% similarity).

*SLFN14* in mice is located on chromosome 11 and comprised of five coding exons and in humans is on chromosome 17 with four coding exons [9]. *SLFN* genes are highly conserved throughout the mammalian species and are located in regions with other genes attributable to T-cell and macrophage development [1]. Early mouse knock-out studies investigating the role of *SLFN1* in the immune response showed no phenotype, suggesting potential functional redundancy or minimal contribution of some members in the process [9]. The precise mechanisms of how some of the specific members of the *SLFN* family cause human disease is still unclear.


### SLFN14 Mutations in Patients with Platelet Disorders and Bleeding

Platelet function disorders are associated with excessive bleeding and with the advent of next-generation sequencing (NGS) including whole exome sequencing (WES) the genetic impact of affected individuals can be evaluated. Initially, three extended families

Correspondence: Neil V. Morgan Institute for Cardiovascular Sciences, University of Birmingham, Institute of Biomedical Research, Edgbaston, Birmingham, B15 2TT, United Kingdom  
Email: [n.v.morgan@bham.ac.uk](mailto:n.v.morgan@bham.ac.uk)  
Color versions of one or more of the figures in the article can be found online at [www.tandfonline.com/ipt](http://www.tandfonline.com/ipt).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Figure 1. Schematic representation of the protein structure encoded by the *SLFN* family of genes, highlighting domains and regions in both *mus musculus* and *homo sapiens*. 'slfn' box is unique to SLFN proteins and function remains unknown. The AAA domain is responsible for DNA and RNA metabolism and the SWADL region, again believed to be SLFN specific, is a sequence flanked by SWA and DL amino acids. The helicase regions at the C-terminal end of the protein are known to mediate DNA and RNA metabolism. The involvement of mutations and thrombocytopenia within the AAA domain of the SLFN14 protein remains unclear.



were investigated in the Genotyping and Phenotyping of Platelets Study, with three heterozygous single nucleotide variations in *SLFN14* resulting in missense mutations of consecutive amino acids in the AAA domain [10]. The following year, an additional family was discovered with another missense mutation in the same region (R223W) and in 2019, Saes et al., reported a further patient with an alternative base change at nucleotide c.657 yet still resulting in the K219N mutation (Table I and Figure 1) [11,12]. All four mutations were inherited in an autosomal dominant fashion whereby amino acid changes are conserved across all mammalian species, except for V220D. The R223W variant is the only variant out of these families to be reported on gnomAD (genome aggregation database) with a frequency of  $6.5 \times 10^{-6}$ . This highlights variant rarity amongst different populations and confinement to the families described. All individuals had macrothrombocytopenia, characterized by a reduced platelet count (below  $150 \times 10^9/L$ ) and enlarged platelets suggesting this is a typical feature of *SLFN14* related thrombocytopenia. Patients from the three initial families showed a platelet function defect with impaired aggregation to ADP, protease-activated receptor-1 (PAR-1) and Collagen and reduced ATP secretion from dense granules after PAR-1 stimulation [10]. Platelet aggregation was not stated for the cases in family D or E but an unspecified secretion defect in the latter case was reported [12]. The proband in family A was 31 years of age at the time of measurement and had experienced severe cutaneous bruising, prolonged bleeding from wounds and menorrhagia [10]. These symptoms were consistent across the three generations and are all

typical of moderate thrombocytopenia. The proband in family B was 35 years old and had a history of spontaneous epistaxis; her mother was also recruited but had not experienced any prior bleeding tendencies [10]. Family C index case was 3 years old at the time, displayed a very low platelet count and reported easy bruising since birth [10]. Patients with variant R223W also reported bleeding symptoms after dental surgery and mild menorrhagia, typical of thrombocytopenia [11]. Individual history for the patient in family E was not detailed in publication however the patient was recruited to the study based on bleeding tendency so, therefore, is likely to be similar to those with the same mutation [12].

All patients in these reported families were heterozygous for the mutations and therefore phenotypically defined by platelet-type bleeding disorder 20 (BDPLT20) which is specific to heterozygous mutations in the *SLFN14* gene. The absence of homozygous patients in the families suggests the mutant allele in each variant group has significant influence over the wild type. Indeed, heterozygous *SLFN14* protein expression was reduced by 65%-80% when compared to controls and in platelet spreading assays there is a reduction of proplatelet extensions from developing megakaryocytes [10,11]. This apparent dominant negative effect on overall protein expression can impact the maturation of megakaryocytes, affecting differentiation and production of fully functional platelets [10,11]. In order to decipher the role of *SLFN14* in platelet biogenesis, it has been demonstrated that *SLFN14* acts as an endoribonuclease in controlling gene expression [13,14]. *SLFN14* is strongly overexpressed in rabbit

Table 1. SLFN14 variants reported to date in patients with inherited macrothrombocytopenia, platelet-type bleeding disorder 20 (BDPLT20).

Patient	Genomic Variant	Protein effect	Variant type	Inheritance	Platelet count ( $\times 10^9/L$ )	Mean platelet volume (MPV fl)	ISTH BAT Score	Aggregation/ Secretion Defect	Reference
A; III 2	c.659 T > A	p.V220D	Missense	Het	140	9.1	5	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
A; III 3	c.659 T > A	p.V220D	Missense	Het	74	10.4	10	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
A; IV 2	c.659 T > A	p.V220D	Missense	Het	110	9.3	13	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
A; IV 4*	c.659 T > A	p.V220D	Missense	Het	100	11.1	22	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
A; IV 5	c.659 T > A	p.V220D	Missense	Het	116	11.2	21	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
B; I 2	c.657 A > T	p.K219N	Missense	Het	83	11.9	13	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
B; II 3*	c.657 A > T	p.K219N	Missense	Het	68	11.9	20	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
C; II 2*	c.652 A > G	p. K218E	Missense	Het	89	13.0	NA	ADP, Collagen and PAR-1-activating peptide/ATP	[10]
D; II 1	c.667 C > T	p. R223W	Missense	Het	87	12.1	5	ADP, Collagen and PAR-1-activating peptide/ATP	[11]
D; II 2	c.667 C > T	p. R223W	Missense	Het	91	21.0	2	NA/NA	[11]
D; III 3*	c.667 C > T	p. R223W	Missense	Het	79	12.3	9	NA/NA	[11]
E; I 1*	c.657 A > C	p.K219N	Missense	Het	NR	NR	NR	NA/NR	[12]

All patients identified with variants in the *SLFN14* gene and affected by inherited bleeding. \*:Proband in family case; Het: Heterozygous inheritance pattern; International Society on Thrombosis and Haemostasis Bleeding Assessment Tool (BAT) score; NA: Not Available for *in vitro* study; NR: Not Reported in publication. Thrombocytopenia was defined as platelet count  $<150 \times 10^9/L$ .

reticulocyte lysate where it is one of the major ribosome-associated proteins [13]. SLFN14 also binds to ribosomes in different model cell lines [14]. Upon binding, SLFN14 cleaves ribosomal RNA and ribosome-associated messenger RNA resulting in ribosomal degradation and disrupting translation [13,14]. Endoribonucleases mediate effective mechanisms in regulating gene expression; however, mutant variants of ribosome-related proteins, such as SLFN14, may cause errors in ribosome homeostasis and subsequent hematopoietic lineage dysfunction [15,16]. For this reason, we can hypothesize that *SLFN14* contributes to improper platelet formation from the disrupted translation of their precursors, megakaryocytes, which then results in an inherited thrombocytopenia [14]. Given the structural similarity of *SLFN13* to other subgroup III genes (such as *SLFN14*) and the recent report on SLFN13's role as an endoribonuclease it seems plausible to hypothesize that SLFN14 may be impacting megakaryocyte differentiation and platelet synthesis by also acting as an endoribonuclease, inhibiting translation and preventing complete protein synthesis within the hematopoietic lineage [8].

It is only with the help of new genetic approaches over recent years, that inherited thrombocytopenias with previously unknown causes can now be attributed to single genetic defects. This not only allows for expanding knowledge within the field of genetics and hematological disorders but also provides patients with a clear diagnosis and a plan for tailored clinical management.

### Main Findings

- Mutations in *SLFN14* cause inherited macrothrombocytopenia characterized by enlarged platelets and moderate to severe bleeding phenotypes.
- Three out of the five families with *SLFN14* genetic variants show reduced aggregation to agonists ADP, PAR-1 and Collagen, decreased ATP secretion and dominant inheritance pattern.
- Endoribonucleolytic activity of SLFN14 contributes to platelet formation by regulating translation process during their maturation.
- The exact role of *SLFN14* in platelet biogenesis and how mutations within its AAA domain contribute to inherited thrombocytopenia remain to be explored.

### Declaration of interest

The authors report no conflict of interest.

### Funding

The work in these authors' laboratory is supported by grants provided by the British Heart Foundation (PG/16/103/32650; FS/18/11/33443) to N.V.M and also supported by a National Institutes of Health Grant GM097014 to A.V.P.

### ORCID

Neil V. Morgan  <http://orcid.org/0000-0001-6433-5692>

### References

1. Schwarz DA, Katayama CD, Hedrick SM. Schlafen, a new family of growth regulatory genes that affect thymocyte development. *Immunity* 1998;9(5):657–668.
2. Geserick P, Kaiser F, Klemm U, Kaufmann SHE, Zerrahn J. Modulation of T cell development and activation by novel members of the Schlafen (slfn) gene family harbouring an RNA helicase-like motif. *Int Immunol* 2004;16(10):1535–1548. doi:10.1093/intimm/dxh155
3. Neumann B, Zhao L, Murphy K, Gonda TJ. Subcellular localization of the Schlafen protein family. *Biochem Biophys Res Commun* 2008;370(1):62–66. doi:10.1016/j.bbrc.2008.03.032
4. Mavrommatis E, Arslan AD, Sassano A, Hua Y, Kroczyńska B, Plataniias LC. Expression and regulatory effects of murine Schlafen (Slfn) genes in malignant melanoma and renal cell carcinoma. *J Biol Chem* 2013;288(46):33006–33015. doi:10.1074/jbc.M113.460741
5. Hanson PI, Whiteheart SW. AAA+ proteins: have engine, will work. *Nat Rev Mol Cell Biol* 2005;6:519. doi:10.1038/nrm1684
6. Lupas AN, Martin J. AAA proteins. *Curr Opin Struct Biol* 2002;12(6):746–753.
7. Liu F, Zhou P, Wang Q, Zhang M, Li D. The Schlafen family: complex roles in different cell types and virus replication. *Cell Biol Int* 2018;42(1):2–8. doi:10.1002/cbin.10778
8. Yang J-Y, Deng X-Y, Li Y-S, Ma X-C, Feng J-X, Yu B, Chen Y, Luo Y-L, Wang X, Chen M-L, et al. Structure of Schlafen13 reveals a new class of tRNA/rRNA-targeting RNase engaged in translational control. *Nat Commun* 2018;9(1):1165. doi:10.1038/s41467-018-03544-x
9. Bustos O, Naik S, Ayers G, Casola C, Perez-Lamigueiro MA, Chippindale PT, Pritham EJ, de la Casa-Esperón E. Evolution of the Schlafen genes, a gene family associated with embryonic lethality, meiotic drive, immune processes and orthopoxvirus virulence. *Gene* 2009;447(1):1–11. doi:10.1016/j.gene.2009.07.006
10. Fletcher SJ, Johnson B, Lowe GC, Bem D, Drake S, Lordkipanidzé M, Guiú IS, Dawood B, Rivera J, Simpson MA, et al. SLFN14 mutations underlie thrombocytopenia with excessive bleeding and platelet secretion defects. *J Clin Invest* 2015;125(9):3600–3605. doi:10.1172/JCI80347
11. Marconi C, Di Buduo CA, Barozzi S, Palombo F, Pardini S, Zaninetti C, Pippucci T, Noris P, Balduini A, Seri M, et al. SLFN14-related thrombocytopenia: identification within a large series of patients with inherited thrombocytopenia. *Thromb Haemost* 2016;115(5):1076–1079. doi:10.1160/TH15-11-0884
12. Saes JL, Simons A, de Munnik SA, Nijziel MR, Blijlevens NMA, Jongmans MC, van der Reijden BA, Smit Y, Brons PP, van Heerde WL, et al. Whole exome sequencing in the diagnostic workup of patients with a bleeding diathesis. *Haemophilia* 2019;25(1):127–135. doi:10.1111/hae.13638
13. Pisareva VP, Muslimov IA, Tcherepanov A, Pisarev AV. Characterization of novel ribosome-associated endoribonuclease SLFN14 from rabbit reticulocytes. *Biochemistry* 2015;54(21):3286–3301. doi:10.1021/acs.biochem.5b00302
14. Fletcher SJ, Pisareva VP, Khan AO, Tcherepanov A, Morgan NV, Pisarev AV. Role of the novel endoribonuclease SLFN14 and its disease-causing mutations in ribosomal degradation. *Rna* 2018;24(7):939–949. doi:10.1261/rna.066415.118
15. Ricciardi S, Miluzio A, Brina D, Clarke K, Bonomo M, Aiolfi R, Guidotti LG, Falciani F, Biffo S. Eukaryotic translation initiation factor 6 is a novel regulator of reactive oxygen species-dependent megakaryocyte maturation. *J Thromb Haemost* 2015;13(11):2108–2118. doi:10.1111/jth.13150
16. Narla A, Ebert BL. Ribosomopathies: human disorders of ribosome dysfunction. *Blood* 2010;115(16):3196–3205. doi:10.1182/blood-2009-10-178129